

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/283329648>

Secondary Metabolite Constituents and Antibacterial Potency of *Physalis angulata* against some Clinical Isolates

Article · January 2014

CITATIONS

4

READS

561

4 authors:



[Marili Funmilayo Zubair](#)

University of Ilorin

16 PUBLICATIONS 223 CITATIONS

[SEE PROFILE](#)



[Ibikunle Ibitayo Anibijuwon](#)

University of Ilorin

36 PUBLICATIONS 480 CITATIONS

[SEE PROFILE](#)



[Mubarak Oloduowo Ameen](#)

University of Ilorin

89 PUBLICATIONS 1,156 CITATIONS

[SEE PROFILE](#)



[Halimat Abdulrahim](#)

University of Ilorin

36 PUBLICATIONS 195 CITATIONS

[SEE PROFILE](#)



Secondary Metabolite Constituents and Antibacterial Potency of *Physalis angulata* against some Clinical Isolates

*¹Zubair, M. F., ²Anibijuwon, I. I., ¹Ameen, O. M. and ³Abdulrahim, H. A.

¹Department of Chemistry, University of Ilorin, P.M.B. 1515, Ilorin, Nigeria.

²Department of Microbiology, Public Health Laboratory Unit, University of Ilorin, P.M.B. 1515, Ilorin, Nigeria.

³Department of Medical Biochemistry, University of Ilorin, P.M.B. 1515, Ilorin, Nigeria

Abstract: *Physalis angulata* is a specie of the Solanaceae family, in which edible fruit is used in several countries of tropical and subtropical regions of the world as medicinal and fruit-tree. In this study, the secondary metabolite screening of the crude extracts consisting of (ethanol, ethylacetate, chloroform and n-hexane) of the whole plant and antibacterial activity was investigated against some clinical isolates. The screening of the secondary metabolites revealed some active components such as alkaloids, tannins, terpenoids, phenolics and flavonoids. The antimicrobial screening of the ethanolic extract of the plant at concentrations of 25, 50, 100 and 125 mg/ml revealed various zones of inhibition from 8.2 to 17.2 mm for the different organisms. In conclusion, the extract of *Physalis angulata* exhibited antibacterial activity made possible by the presence of the secondary metabolite constituents and therefore could be used as antibiotic against the bacteria strains investigated in this study.

KEYWORDS: *Physalis angulata*, Solanaceae, Antibacterial, Clinical isolates

1.0 Introduction

The use of natural products in the treatment of a variety of diseases has increased due to the considerable number of medicinal plants with proven biological activities applicable to the treatment of some diseases.

Physalis angulata L., (family: Solanaceae), known in the Peruvian, Amazon as mullaca, is a much branched annual shrub distributed in many tropical countries and subtropical regions of the world. It can grow until it reaches 1.0 m. The flowers are bell shaped but the most distinctive feature is the fruiting calyx which enlarges to cover the fruit and hangs downward like a lantern. Each fruit is like a yellow pearl into a lantern shaped pod (Mejia and Rengifo, 2000).

Physalis angulata is used in many part of the world to treat several diseases such as cancer, bacteriemia, tumors, gonorrhea, diabetes

(macerated roots), infertility (fruits), malaria (Ruiz *et al.*, 2011), diuresis, inflammation of the bladder, asthma (leaf infusion), hepatitis (root infusion) Mejia and Rengifo, 2000), rheumatism, stomach ache, prostrate and kidney stones, (Coelho-Ferreira, 2009), anaemia, to prevent abortion, and in reducing fever. In Peru, the decoction of the leaves and fruits is used for "Tertian" due to postpartum infections (Jovel *et al.*, 1996). Their anti-inflammatory and antispasmodic properties are also known (Silva *et al.*, 2005).

Phytochemical studies on *P. angulata* revealed that it contains flavonoids, alkaloids and many different types of plant steroids. The main component are Physalins B, G, D, E, F, H, I, K, U and V isolated from root, stems and leaves of *P. angulata* (Damu *et al.*, 2007; Chen *et al.*, 2011); withanoloides A, B, C and D were obtained from the methanolic extract of the fresh leaves, stems and aerial parts of *P. angulata* (Shingu *et al.*, 1991, 1992; He *et al.*, 2007) and carotenoids (De Rosso and Mercadante, 2007).

***Corresponding Author**

Tel.: +2348030661412

E-mail: marilizub@unilorin.edu.ng

Its biological properties include antimicrobial, anticancerous, antitumorous, antimalarial, anti-inflammatory, antileishmanial, hypotensive and immunosuppressive activity (Helvacı *et al.*, 2010; He *et al.*, 2007; Ferreira *et al.*, 2006; Sa *et al.*, 2011; Pinto *et al.*, 2010; Guimaraes *et al.*, 2010; Yu *et al.*, 2010).

Due to the dearth of information on the antibacterial activity of the ethanolic extract of *Physalis angulata* whole plant, this study therefore reports on the secondary metabolite constituents of some solvent extracts of the whole plant and the effects of the ethanolic extract on some clinical isolates.

2.0 Materials and Methods

2.1 Plant Material

Fresh whole plant of *Physalis angulata* was collected from Odo-Isuti Egan, Lagos State, Nigeria in December, 2013. The plant was identified at the Plant Biology Department of the University of Ilorin, Nigeria.

2.2 Test Organisms

The test organisms used for this research, *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumonia* and *Pseudomonas aeruginosa* were gotten from the Microbiology and Parasitology Laboratory of the University of Ilorin Teaching Hospital, Ilorin, Nigeria.

2.3 Preparation of Ethanolic Extract

This was air-dried for two weeks and grounded into powder. Two litres of ethanol was added to 250 g of powdered *Physalis angulata* and was cold extracted for a week. This was filtered using Whatman No. 2 filter paper and concentrated using rotatory evaporation to obtain the crude ethanol extract. The crude ethanol extract was successively macerated with n-hexane, chloroform and ethylacetate respectively.

2.4 Screening of Secondary Metabolites

The crude extracts was screened for the presence of alkaloids, tannins, saponins,

steroids, terpenoids, phenolics, flavonoids and quinones according to the standard procedures (Harbone, 1978; Sofowora, 1993; Trease and Evans, 1989).

2.5 Antimicrobial Assay

2.5.1 Preparation of Media

All media were prepared according to research specifications following the manufacturers' instructions.

2.5.2 Maintenance of the Test Organisms

The organism was inoculated into nutrient agar slants, incubated and kept at 4°C prior to sub-culture.

2.5.3 Standardization of Inoculum

Colonies of each organism growing as a pure culture in the nutrient agar plate were inoculated into 10mls of peptone water in different test tubes respectively for each organism aseptically. The mouth of the tubes were covered with cotton wool and wrapped with aluminum foil followed by incubation at 37°C for 24hrs.

2.5.4 Dilution Methods

Colonies of each organism growing as a pure culture in the nutrient agar plate were inoculated into 10mls of peptone water in different test tubes respectively for each organism aseptically. The mouth of the tubes were covered with cotton wool and wrapped with aluminum foil followed by incubation at 37°C for 24hours.

2.5.5 Agar Dilution

Tests were carried out according to the Farmacopéia Brasileira (1988) and USP XXIII Pharmacopoeia (1995) procedures using impregnated disks of different concentrations. The plates were incubated at 35 °C for 18 h in an ambient air incubator. The zone diameters were measured.

3.0 Results and Discussion

The use of natural products to treat variety of diseases has increased due to large number of

medicinal plants that have shown biological activity.

The preliminary screening of the secondary metabolites in *Physalis angulata* whole plant revealed a wide array of phytoconstituents depending on the solvent used. For instance, among all chemical compounds like alkaloids, tannins, saponins, steroids, terpenoids, phenolics, flavonoids and quinones that were screened for, all except saponins were present in the ethanolic, ethyl acetate and n-hexane extracts (Table 1). Apart from saponins that were not detected in the chloroform extract, the extract did not also show detectable quantities of quinones (Table 1). Several of these secondary metabolites have been reported to be responsible for the diverse medicinal properties of these plants. Some of these compounds have also been associated with antibacterial activities and thus have curative properties against pathogens.

The ethanolic extract of the whole part of *Physalis angulata* inhibited the growth of all the bacterial species (*S. aureus*, *E. coli*, *K. pneumonia*, *P. aeruginosa*) investigated in the present study as evidenced by the absence of turbidity, although at different concentrations. The zones of inhibition produced by the ethanolic extract of the whole plant of *Physalis angulata* ranged from 8.2 mm to 17.2 mm and were concentration dependent (Table 2). Infectious disease of microbial origin caused by *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumonia* and *Pseudomonas aeruginosa* constitute major causes of morbidity and mortality in several countries of the world today (Kloos and Keim, 1993). Such microbial infection can result in diarrhoea, pneumonia, typhoid etc. The zones of inhibition and the absence of turbidity in the extract treated colonies of the bacteria confer antibacterial activity on the ethanolic extract of *Physalis angulata* whole plant. The finding is similar to

the report by Hwang *et al* (2004), where the bactericidal test showed that methanol extract of *P. angulata* conferred fast killing effect against *S. mutans* in 2 min at 50 mg/ml concentration.

Even though the exact antimicrobial mechanism of phenolic compounds is not clear, phenolic compounds are commonly known for their antimicrobial effects (Ciocan and Ioan, 2007; Tiwari *et al.*, 2009). The ability of phenolic compounds to alter microbial cell permeability, thereby permitting the loss of macromolecules from the cell interior, could help explain some of the antimicrobial activity (Bajpai *et al.*, 2008). In addition, the concentration dependent effects of the extract in the present study could also be attributed to the effect of phenolic compounds; at low concentration, phenols affect enzyme activity while at high concentrations they cause protein denaturation. Again, Flavonoid rich plant extracts from different species have been reported to possess antibacterial activity (Mishra *et al.*, 2011, Mishra *et al.*, 2013). Antibacterial flavonoids might be having multiple cellular targets, rather than one specific site of action. One of their molecular actions is to form complex with proteins through nonspecific forces such as hydrogen bonding and hydrophobic effects, as well as by covalent bond formation. Thus, their mode of antimicrobial action may be related to their ability to inactivate microbial adhesins, enzymes, cell envelope transport proteins, and so forth. Lipophilic flavonoids may also disrupt microbial membranes (Cowan, 1999; Mishra *et al.*, 2009). Therefore, the presence of phenolics and or flavonoids in the extract of the plant in this study may account for the antibacterial activity of *Physalis angulata*.

In conclusion, the extract has shown antibacterial activity against the test organisms. Therefore, it could be explored as an antibiotic.

Table 1: Secondary metabolite constituents of the crude extracts of whole plant of *Physalis angulata*

Secondary Metabolites	Ethanol	Ethyl acetate	n-henxane	Chloroform
Alkaloids	+	+	+	+
Tannins	+	+	+	+
Saponin	-	-	-	-
Steroids	+	+	+	+
Terpenoids	+	+	+	+
Phenolics	+	+	+	+
Flavonoids	+	+	+	+
Quinones	+	+	+	-

Key: + = Present

- = Absent

Table 2: Zones of inhibition of the bacteria by the ethanolic extract of the whole plant of *Physalis angulata***Concentration/Zones of Inhibition (mm)**

Bacterial Species	25mg/ml	50 mg/ml	100 mg/ml	125 mg/ml
<i>S. aureus</i>	8.2	9.6	10.6	11.6
<i>E. coli</i>	10	11.2	12.6	13.0
<i>K. pneumonia</i>	14.0	14.8	16.2	18.0
<i>P.aeruginosa</i>	12.0	15.0	16.5	17.2

References

- Bajpai, V. K., Rahman, A., Dung, N. T., Huh, M. K. and Kang, S. C. (2008) In vitro inhibition of food spoilage and foodborne pathogenic bacteria by essential oil and leaf extracts of *Magnolia liliflora* Desr. Journal of Food Science. Pg73:M314.
- Chen, L. X., He, H. and Qui, F. (2011). Natural withanolides:an overview. Natural Product Rep 28:705-740.
- Ciocan, D. and Ioan, B. (2007). Plant products as antimicrobial agents. Analele Stiintifice ale Universitatii" Alexandru Ioan Cuza" din Iasi Sec II a Genetica si Biologie Moleculara. pp. 8.
- Coelho-Ferreira, M. (2009). Medicinal knowledge and plant utilization in an Amazonian coastal community of Marudu para State (Brazil). Journal of Ethnopharmacology 126:159-175.
- Cowan, M. M. (1999). Plant products as antimicrobial agents. Clinical Microbiology Reviews 12(4): 564-582.
- Damu, A. G., Kuo, P., Su, C., Kuo, T., Chen, T., Bastow, K. F., Lee, K.. and Wu, T. (2007). Isolation, structures, and structure-cytotoxic activity relationships of withanolides and physalins from *Physalis angulata*. Journal of Natural Product 70:1146-1152.
- De Rosso, V. V. and Mercadante, A. Z. (2007). Identification and quantification de carotenoids, by HPLC-PDA-MS/MS, from Amazonian fruits. Journal of Agriculture Food Chemistry 55:5062-5072.
- Ferreira, H. L., Leopoldina, M., Rocha, M., Nergreiros, A. P., Loiola, O., Rocha, E., Veras, L., Odorico, M. and Pessoa, C. (2006). *In-vitro* and *in vivo* antitumor activity of physalins B and D from

- Physalis angulata*. Journal of Pharmacy and Pharmacology 58:235-241.
- Solanaceae, on the viability of leishmaniasis sp. Brazilian Journal of Pharmacognosy 20:945-949.
- Harbone, J. B. (1978). Phytochemical Methods. Chapman and Hall Ltd. London, pp. 49-188.
- He, Q. P., Ma, L., Luo, J. Y., He, F. Y., Lou, L.G., Hu, L. H. (2007). Cytotoxic withanolides from *Physalis angulata* L. Chemical Biodiversity 4:443-449.
- Helvacı, S., Kokdil, G., Kawai, M., Duran, N., Duran, G. and Guvenç, A. (2010). Antimicrobial activity of the extracts and physalin D from *Physalin alkekengi* and evaluation potential of antioxidant of physalin D. Pharmaceutical Biology 48:142-150.
- Hwang, J., Shim, J. and Chung, J. (2004). Anticariogenic activity of some tropical, medicinal plants against streptococcus mutans. Fitoterapia 75:596-598.
- Jovel, E., Towers, G. and Cabanillas, J. (1996). An ethnobotanical study of the traditional medicine of the Mestizo people of Suni Miraflores, Loreto, Peru. Journal of Ethnopharmacology 53: 149 -153.
- Kloss, H. and Zain, A. H. (1993). The ecology of health and diseases. Washington DC: West View Press Inc.
- Mejia, K. and Rengifo, E. (2000). Plants medicinales de uso popular en la Amazonia peruana. 2nd ed. Instituto de Investigaciones de la Amazonia Peruana, Iquitos, Peru.
- Mishra, A., Kumar, S., Bhargava, A., Sharma, B. and Pandey, A. K. (2011). Studies on *in vitro* antioxidant and antistaphylococcal activities of some important medicinal plants. Cellular and Molecular Biology 57(1): 16-25.
- Mishra, A., Kumar, S. and Pandey, A. K. (2013). Scientific validation of the medicinal efficacy of *Tinospora cordifolia*, The Scientific World Journal vol. 2013, Article ID 292934.
- Mishra, A. K., Mishra, A., Kehri, H. K., Sharma, B. and Pandey, A. K. (2009). Inhibitory activity of Indian spice plant *Cinnamomum zeylanicum* extracts against *Alternaria solani* and *Curvularia lunata*, the pathogenic dematiaceous moulds. Annals of Clinical Microbiology and Antimicrobials, Vol. 8, article 9.
- Pinto, N., Morais T., Carvilho K., Silva C., Andrade, G., Brito G., Veras M., Pessoa O., Rao, V. and Santo, F. (2010) Tropical anti-inflammatory potential of physalin E from *Physalin angulata* on experimental dermatitis in mice. Phytomedicine 17:740-743
- Ruiz, L., Maco, M., Cobos, M., Gutierrez-Choquevilca, A. and Roumy, V. (2011). Plants Guimaraes, E., Lima, M., Santos, L., Ribeiro, R., Dos Santos, W. and Soares, M. (2010). Effects of seco-steroids purified from *Physalin angulata* L. used by native Amazonian groups from the Naay River (Peru) for the treatment of malaria. Journal of Ethnopharmacology 133:917-921 .
- Sa, M. S., de Menezes, M. N., Kretli A. U, Ribeiro, I. M., Tomassini, T. C., Ribeiro dos Santos, R., de Azevedo, W. F. Jr. and Soares, M. B. (2011). Antimalarial activity of physalin B, D, F and G. Journal of Natural Product 74:2269-2272.
- Shingu, K., Marubayashi N., Ueda, I., Yahara, S. and Nohara T. (1991). Physagulin C: a new withanolide from *Physalis angulata* L. Chemical and Pharmaceutical Bulletin 39:1591-1593.
- Shingu, K., Yahara S., Nohara T. and Okabe H. (1992). Three new withanolides, physagulins A, B and D from *Physalis angulata* L. Chemical and Pharmaceutical Bulletin 40:2088-2091
- Silva, M., Simas, S., Batista, T., Cardarelli, P. and Tomassini T. (2005). Studies on antimicrobial activity, *in vitro*, of *Physalis angulata* L. (Solanaceae) fraction and physalin B bringing out the importance of assay determination. Memorial Institute, Oswaldo Cruz 100: 779 – 782.
- Sofowora, A. (1993). Medicinal Plants and Traditional Medicine in Africa. Spectrum Book Ltd., Ibadan, Nigeria, pp 191-289.
- Tiwari, B. K., Valdramidis, V. P., O'Donnell, C. P., Muthukumarappan, K., Bourke, P. and Cullen, P. (2009). Application of natural antimicrobials for food preservation. Journal of Agricultural and Food Chemistry 57:5987-5992.
- Trease, G. E. and Evans, W. C. (1989). Pharmacognosy, 11th Edition, Bailliere Tindall, London, pp. 45-50.
- Yu. Y., Sun, L., Ma, L., Li, J., Hu, L. and Liu, J. (2010). Investigation of the immunosuppressive activity of physalin H on T lymphocytes. International Immunopharmacology 10:290-297.