


Bis(acetonitrile- κN)dichlorido(η^4 -cycloocta-1,5-diene)ruthenium(II) acetonitrile monosolvate

Haleden Chiririwa, Reinout Meijboom, Samson O. Owalude, Uche B. Eke and Charmaine Arderne

Acta Cryst. (2011). **E67**, m1096

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The journal publishes over 4000 structures per year. The average publication time is less than one month.

Crystallography Journals **Online** is available from journals.iucr.org

Bis(acetonitrile- κ N)dichlorido(η^4 -cycloocta-1,5-diene)ruthenium(II) acetonitrile monosolvate

Haleden Chiririwa,^{a*} Reinout Meijboom,^a Samson O. Owalude,^b Uche B. Eke^b and Charmaine Arderne^a

^aDepartment of Chemistry, University of Johannesburg, PO Box 524, Auckland Park, Johannesburg 2006, South Africa, and ^bDepartment of Chemistry, University of Ilorin, P M B 1515, Ilorin, Nigeria

Correspondence e-mail: harrychiririwa@yahoo.com


Received 14 June 2011; accepted 11 July 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.022; wR factor = 0.054; data-to-parameter ratio = 22.4.

In the title Ru^{II} complex, $[\text{RuCl}_2(\text{C}_8\text{H}_{12})(\text{C}_2\text{H}_3\text{N})_2] \cdot \text{CH}_3\text{CN}$, the metal ion is coordinated to the centers of each of the double bonds of the cyclooctadiene ligand, to two chloride ions (in *cis* positions) and to two N-atom donors (from MeCN molecules) that complete the coordination sphere for the neutral complex. The coordination about the Ru^{II} atom can thus be considered to be octahedral with a slightly trigonal distortion. There is also one acetonitrile solvent molecule per molecule which is outside the coordination sphere of the ruthenium atom.

Related literature

For the structure of the water solvate complex, see: Ashworth *et al.* (1987).



Experimental

Crystal data

$[\text{RuCl}_2(\text{C}_8\text{H}_{12})(\text{C}_2\text{H}_3\text{N})_2] \cdot \text{C}_2\text{H}_3\text{N}$
 $M_r = 403.31$
 Monoclinic, $P2_1/c$
 $a = 8.7033$ (3) Å
 $b = 7.2434$ (3) Å
 $c = 26.4178$ (10) Å
 $\beta = 95.903$ (1)°

$V = 1656.59$ (11) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.26$ mm⁻¹
 $T = 100$ K
 $0.32 \times 0.20 \times 0.13$ mm

Data collection

Bruker APEXII CCD
 diffractometer
 Absorption correction: numerical
 (AXScale; Bruker, 2010)
 $T_{\min} = 0.688$, $T_{\max} = 0.853$

16175 measured reflections
 4125 independent reflections
 3806 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.022$
 $wR(F^2) = 0.054$
 $S = 1.07$
 4125 reflections

184 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.67$ e Å⁻³
 $\Delta\rho_{\min} = -0.61$ e Å⁻³

Table 1

Selected bond lengths (Å).

Ru1—N1	2.0303 (15)	Ru1—C4	2.2154 (17)
Ru1—N2	2.0418 (15)	Ru1—C5	2.2225 (17)
Ru1—Cl1	2.2082 (16)	Ru1—Cl1	2.4212 (4)
Ru1—C8	2.2116 (17)	Ru1—Cl2	2.4265 (4)

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2010); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We gratefully acknowledge the University of Johannesburg for funding this project and the University of Witwatersrand for award of a Research fellowship to SO. The authors also thank Dr E Singleton for fruitful discussions.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GO2018).

References

- Ashworth, T. V., Liles, D. C., Robinson, D. J., Singleton, E., Coville, N. J., Darling, E. & Markwell, J. (1987). *S. Afr. J. Chem.* **40**, 183–188.
- Bruker (2010). *APEX2*, *AXScale* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, m1096 [doi:10.1107/S1600536811027723]

Bis(acetonitrile- κ N)dichlorido(η^4 -cycloocta-1,5-diene)ruthenium(II) acetonitrile monosolvate

H. Chiririwa, R. Meijboom, S. O. Owalude, U. B. Eke and C. Arderne

Comment

The present ruthenium complex, Fig.1, has been synthesized earlier (Ashworth *et al.* 1987). The structure obtained by Ashworth *et al.* was a room temperature determination and with a water molecule as solvate. The current low temperature determination presents an acetonitrile molecule in the crystal lattice that is outside the coordination sphere of the ruthenium. To the best of our knowledge, there are no reports of other structures determined with organonitriles, obtained from the ruthenium [$\{\text{RuCl}_2(\text{COD})\}_x$] polymer. Organonitriles have been used over many years in synthetic inorganic chemistry and an interest in the chemistry of metal-nitrile complexes has prompted several reviews.

The two acetonitrile ligands are not *trans* to each other, as the N(1)—Ru—N(2) angle is 163.15 (6)°. This is due to repulsion by the alkene bonds of the COD ligand. It would seem that one of the acetonitrile ligands is slightly bent. The N(1)—C(9)—C(10) bond angle is 179.11 (19)°, whereas the same angle for the other acetonitrile is 178.3 (2)°. This is due to packing forces.

Experimental

A suspension of [$\{\text{RuCl}_2(\text{COD})\}_x$] (0.5 g) in acetonitrile (20 ml) was refluxed for 6 h. The orange solution was filtered hot and concentrated on a steam bath to half volume and cooled to 0° C overnight affording orange crystals suitable for X-ray diffraction studies.

Refinement

Hydrogen atoms could be identified from the difference Fourier map but once these atoms were refined, their distances from the parent atoms were found to be significantly shorter than the ideal distances for C—H and N—H respectively. The H-atoms were therefore geometrically positioned and refined in the riding-model approximation, with C—H = 0.97 Å, N—H = 0.89 Å, and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{N})$. For (I), the highest peak in the final difference map is 0.55 Å from H1 and the deepest hole is 0.22 Å from H5.

Figures




Fig. 1. A perspective view of $\text{C}_{12}\text{H}_{18}\text{Cl}_2\text{N}_2\text{Ru} \cdot \text{C}_2\text{H}_3\text{N}$ (1) showing the atomic numbering scheme.

Bis(acetonitrile- κ N)dichlorido(η^4 -cycloocta-1,5-diene)ruthenium(II) acetonitrile monosolvate

Crystal data

$[\text{RuCl}_2(\text{C}_8\text{H}_{12})(\text{C}_2\text{H}_3\text{N})_2] \cdot \text{C}_2\text{H}_3\text{N}$	$F(000) = 816$
$M_r = 403.31$	$D_x = 1.617 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 9251 reflections
$a = 8.7033 (3) \text{ \AA}$	$\theta = 2.7\text{--}28.3^\circ$
$b = 7.2434 (3) \text{ \AA}$	$\mu = 1.26 \text{ mm}^{-1}$
$c = 26.4178 (10) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 95.903 (1)^\circ$	Block, orange
$V = 1656.59 (11) \text{ \AA}^3$	$0.32 \times 0.20 \times 0.13 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD diffractometer	4125 independent reflections
Radiation source: fine-focus sealed tube	3806 reflections with $I > 2\sigma(I)$
graphite	$R_{\text{int}} = 0.030$
φ and ω scans	$\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 1.6^\circ$
Absorption correction: numerical (AXScale; Bruker, 2010)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.688$, $T_{\text{max}} = 0.853$	$k = -6 \rightarrow 9$
16175 measured reflections	$l = -35 \rightarrow 35$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.022$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.054$	H-atom parameters constrained
$S = 1.07$	$w = 1/[\sigma^2(F_o^2) + (0.0202P)^2 + 1.4236P]$
4125 reflections	where $P = (F_o^2 + 2F_c^2)/3$
184 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.67 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.61 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. The Following Model and Quality ALERTS were generated - (Acta-Mode) <<< Format: alert-number_ALERT_alert-type_alert-level text 912_ALERT_4_C Missing # of FCF Reflections Above STh/L= 0.600 5 Noted.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ru1	0.732203 (14)	0.145839 (19)	0.109901 (5)	0.01058 (5)
Cl1	0.77031 (5)	0.38454 (6)	0.048264 (16)	0.01597 (9)
Cl2	1.00549 (5)	0.09785 (6)	0.134734 (15)	0.01607 (9)
N1	0.76802 (16)	−0.0274 (2)	0.05184 (5)	0.0132 (3)
N2	0.75889 (16)	0.3467 (2)	0.16432 (6)	0.0145 (3)
N3	0.2246 (2)	0.6581 (3)	0.20027 (7)	0.0349 (5)
C1	0.49326 (19)	0.1021 (3)	0.07512 (7)	0.0152 (3)
H1	0.5348	0.1237	0.0438	0.018*
C2	0.4309 (2)	−0.0888 (3)	0.08525 (7)	0.0180 (4)
H2A	0.3862	−0.1427	0.0526	0.022*
H2B	0.3464	−0.0762	0.1074	0.022*
C3	0.5526 (2)	−0.2234 (3)	0.11060 (7)	0.0180 (4)
H3A	0.5015	−0.3090	0.1328	0.022*
H3B	0.5947	−0.2980	0.0838	0.022*
C4	0.6854 (2)	−0.1285 (2)	0.14209 (7)	0.0157 (3)
H4	0.7877	−0.1611	0.1361	0.019*
C5	0.6665 (2)	0.0024 (3)	0.17886 (6)	0.0159 (3)
H5	0.7561	0.0577	0.1961	0.019*
C6	0.5090 (2)	0.0627 (3)	0.19318 (7)	0.0183 (4)
H6A	0.5192	0.0962	0.2297	0.022*
H6B	0.4372	−0.0433	0.1885	0.022*
C7	0.4375 (2)	0.2278 (3)	0.16207 (7)	0.0181 (4)
H7A	0.3237	0.2144	0.1583	0.022*
H7B	0.4630	0.3429	0.1813	0.022*
C8	0.49209 (19)	0.2461 (3)	0.10961 (7)	0.0153 (3)
H8	0.5277	0.3635	0.0998	0.018*
C9	0.8070 (2)	−0.1171 (3)	0.02039 (6)	0.0145 (3)
C10	0.8594 (2)	−0.2322 (3)	−0.01961 (7)	0.0189 (4)
H10A	0.9666	−0.2698	−0.0100	0.028*
H10B	0.8531	−0.1621	−0.0515	0.028*
H10C	0.7938	−0.3421	−0.0242	0.028*
C11	0.79305 (19)	0.4484 (3)	0.19626 (6)	0.0150 (3)
C12	0.8397 (2)	0.5747 (3)	0.23806 (7)	0.0211 (4)
H12A	0.8546	0.5054	0.2700	0.032*
H12B	0.7591	0.6680	0.2403	0.032*
H12C	0.9366	0.6355	0.2319	0.032*

supplementary materials

C13	0.1734 (2)	0.6361 (3)	0.15954 (8)	0.0217 (4)
C14	0.1070 (3)	0.6105 (3)	0.10732 (8)	0.0288 (5)
H14A	0.0412	0.7163	0.0968	0.043*
H14B	0.0451	0.4973	0.1048	0.043*
H14C	0.1901	0.6006	0.0851	0.043*

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ru1	0.01023 (7)	0.00908 (8)	0.01235 (7)	−0.00028 (5)	0.00080 (5)	−0.00063 (5)
Cl1	0.01604 (19)	0.0128 (2)	0.01920 (19)	−0.00040 (15)	0.00236 (15)	0.00325 (16)
Cl2	0.01206 (18)	0.0203 (2)	0.01555 (18)	0.00113 (16)	−0.00022 (14)	0.00109 (16)
N1	0.0121 (6)	0.0121 (7)	0.0151 (6)	−0.0007 (6)	0.0004 (5)	0.0006 (6)
N2	0.0123 (6)	0.0142 (8)	0.0171 (7)	−0.0003 (6)	0.0020 (5)	0.0001 (6)
N3	0.0387 (11)	0.0370 (12)	0.0282 (9)	0.0033 (9)	−0.0009 (8)	0.0035 (8)
C1	0.0102 (7)	0.0168 (9)	0.0185 (8)	−0.0008 (7)	0.0004 (6)	0.0010 (7)
C2	0.0173 (8)	0.0149 (9)	0.0219 (8)	−0.0046 (7)	0.0021 (7)	−0.0027 (7)
C3	0.0211 (9)	0.0113 (9)	0.0223 (8)	−0.0023 (7)	0.0058 (7)	−0.0005 (7)
C4	0.0176 (8)	0.0112 (9)	0.0189 (8)	−0.0008 (7)	0.0049 (6)	0.0039 (7)
C5	0.0163 (8)	0.0147 (9)	0.0170 (8)	−0.0016 (7)	0.0034 (6)	0.0032 (7)
C6	0.0181 (8)	0.0191 (10)	0.0186 (8)	−0.0017 (7)	0.0056 (7)	−0.0023 (7)
C7	0.0141 (8)	0.0170 (9)	0.0237 (9)	−0.0012 (7)	0.0044 (7)	−0.0043 (7)
C8	0.0101 (7)	0.0124 (9)	0.0233 (8)	−0.0008 (6)	0.0011 (6)	−0.0007 (7)
C9	0.0134 (7)	0.0136 (9)	0.0163 (8)	−0.0010 (7)	0.0002 (6)	0.0028 (7)
C10	0.0226 (9)	0.0182 (10)	0.0168 (8)	0.0030 (8)	0.0059 (7)	−0.0013 (7)
C11	0.0140 (7)	0.0157 (9)	0.0156 (8)	0.0005 (7)	0.0029 (6)	0.0032 (7)
C12	0.0258 (9)	0.0201 (10)	0.0172 (8)	−0.0038 (8)	0.0018 (7)	−0.0045 (7)
C13	0.0192 (9)	0.0161 (10)	0.0304 (10)	0.0005 (7)	0.0059 (7)	−0.0002 (8)
C14	0.0274 (10)	0.0304 (12)	0.0281 (10)	−0.0060 (9)	0.0001 (8)	−0.0094 (9)

Geometric parameters ($\text{\AA}$, $^\circ$)

Ru1—N1	2.0303 (15)	C5—C6	1.523 (2)
Ru1—N2	2.0418 (15)	C5—H5	0.9500
Ru1—C1	2.2082 (16)	C6—C7	1.545 (3)
Ru1—C8	2.2116 (17)	C6—H6A	0.9900
Ru1—C4	2.2154 (17)	C6—H6B	0.9900
Ru1—C5	2.2225 (17)	C7—C8	1.517 (2)
Ru1—Cl1	2.4212 (4)	C7—H7A	0.9900
Ru1—Cl2	2.4265 (4)	C7—H7B	0.9900
N1—C9	1.134 (2)	C8—H8	0.9500
N2—C11	1.136 (2)	C9—C10	1.455 (2)
N3—C13	1.133 (3)	C10—H10A	0.9800
C1—C8	1.385 (3)	C10—H10B	0.9800
C1—C2	1.520 (3)	C10—H10C	0.9800
C1—H1	0.9500	C11—C12	1.459 (3)
C2—C3	1.541 (3)	C12—H12A	0.9800
C2—H2A	0.9900	C12—H12B	0.9800
C2—H2B	0.9900	C12—H12C	0.9800

C3—C4	1.517 (2)	C13—C14	1.452 (3)
C3—H3A	0.9900	C14—H14A	0.9800
C3—H3B	0.9900	C14—H14B	0.9800
C4—C5	1.379 (3)	C14—H14C	0.9800
C4—H4	0.9500		
N1—Ru1—N2	163.15 (6)	C3—C4—Ru1	110.84 (12)
N1—Ru1—C1	78.95 (6)	C5—C4—H4	118.0
N2—Ru1—C1	115.42 (6)	C3—C4—H4	118.0
N1—Ru1—C8	114.75 (6)	Ru1—C4—H4	87.0
N2—Ru1—C8	78.91 (6)	C4—C5—C6	123.22 (16)
C1—Ru1—C8	36.53 (7)	C4—C5—Ru1	71.62 (10)
N1—Ru1—C4	77.55 (6)	C6—C5—Ru1	112.56 (12)
N2—Ru1—C4	112.38 (6)	C4—C5—H5	118.4
C1—Ru1—C4	80.14 (7)	C6—C5—H5	118.4
C8—Ru1—C4	94.86 (7)	Ru1—C5—H5	85.9
N1—Ru1—C5	113.76 (7)	C5—C6—C7	114.42 (15)
N2—Ru1—C5	77.06 (6)	C5—C6—H6A	108.7
C1—Ru1—C5	87.90 (7)	C7—C6—H6A	108.7
C8—Ru1—C5	80.42 (7)	C5—C6—H6B	108.7
C4—Ru1—C5	36.21 (7)	C7—C6—H6B	108.7
N1—Ru1—C11	83.74 (4)	H6A—C6—H6B	107.6
N2—Ru1—C11	87.20 (4)	C8—C7—C6	114.01 (15)
C1—Ru1—C11	90.62 (5)	C8—C7—H7A	108.8
C8—Ru1—C11	87.59 (5)	C6—C7—H7A	108.8
C4—Ru1—C11	160.38 (5)	C8—C7—H7B	108.8
C5—Ru1—C11	161.74 (5)	C6—C7—H7B	108.8
N1—Ru1—C12	83.88 (4)	H7A—C7—H7B	107.6
N2—Ru1—C12	82.78 (4)	C1—C8—C7	124.02 (17)
C1—Ru1—C12	161.26 (5)	C1—C8—Ru1	71.60 (10)
C8—Ru1—C12	161.37 (5)	C7—C8—Ru1	110.66 (11)
C4—Ru1—C12	88.94 (5)	C1—C8—H8	118.0
C5—Ru1—C12	92.25 (5)	C7—C8—H8	118.0
C11—Ru1—C12	94.926 (15)	Ru1—C8—H8	87.7
C9—N1—Ru1	171.31 (14)	N1—C9—C10	179.11 (19)
C11—N2—Ru1	170.70 (15)	C9—C10—H10A	109.5
C8—C1—C2	122.84 (16)	C9—C10—H10B	109.5
C8—C1—Ru1	71.87 (10)	H10A—C10—H10B	109.5
C2—C1—Ru1	113.29 (12)	C9—C10—H10C	109.5
C8—C1—H1	118.6	H10A—C10—H10C	109.5
C2—C1—H1	118.6	H10B—C10—H10C	109.5
Ru1—C1—H1	85.0	N2—C11—C12	178.3 (2)
C1—C2—C3	114.21 (15)	C11—C12—H12A	109.5
C1—C2—H2A	108.7	C11—C12—H12B	109.5
C3—C2—H2A	108.7	H12A—C12—H12B	109.5
C1—C2—H2B	108.7	C11—C12—H12C	109.5
C3—C2—H2B	108.7	H12A—C12—H12C	109.5
H2A—C2—H2B	107.6	H12B—C12—H12C	109.5
C4—C3—C2	113.72 (15)	N3—C13—C14	179.2 (2)
C4—C3—H3A	108.8	C13—C14—H14A	109.5

supplementary materials

C2—C3—H3A	108.8	C13—C14—H14B	109.5
C4—C3—H3B	108.8	H14A—C14—H14B	109.5
C2—C3—H3B	108.8	C13—C14—H14C	109.5
H3A—C3—H3B	107.7	H14A—C14—H14C	109.5
C5—C4—C3	123.94 (16)	H14B—C14—H14C	109.5
C5—C4—Ru1	72.17 (10)		
N1—Ru1—C1—C8	−168.61 (12)	N1—Ru1—C5—C4	−0.84 (12)
N2—Ru1—C1—C8	2.09 (13)	N2—Ru1—C5—C4	−167.20 (11)
C4—Ru1—C1—C8	112.29 (12)	C1—Ru1—C5—C4	76.11 (11)
C5—Ru1—C1—C8	76.69 (11)	C8—Ru1—C5—C4	112.09 (11)
Cl1—Ru1—C1—C8	−85.11 (10)	Cl1—Ru1—C5—C4	161.71 (12)
Cl2—Ru1—C1—C8	167.48 (12)	Cl2—Ru1—C5—C4	−85.14 (10)
N1—Ru1—C1—C2	72.69 (13)	N1—Ru1—C5—C6	−119.99 (13)
N2—Ru1—C1—C2	−116.61 (13)	N2—Ru1—C5—C6	73.65 (13)
C8—Ru1—C1—C2	−118.70 (18)	C1—Ru1—C5—C6	−43.04 (13)
C4—Ru1—C1—C2	−6.41 (13)	C8—Ru1—C5—C6	−7.06 (13)
C5—Ru1—C1—C2	−42.01 (13)	C4—Ru1—C5—C6	−119.15 (18)
Cl1—Ru1—C1—C2	156.19 (12)	Cl1—Ru1—C5—C6	42.6 (2)
Cl2—Ru1—C1—C2	48.8 (2)	Cl2—Ru1—C5—C6	155.72 (12)
C8—C1—C2—C3	−91.4 (2)	C4—C5—C6—C7	−90.1 (2)
Ru1—C1—C2—C3	−8.64 (19)	Ru1—C5—C6—C7	−7.95 (19)
C1—C2—C3—C4	26.6 (2)	C5—C6—C7—C8	26.4 (2)
C2—C3—C4—C5	50.8 (2)	C2—C1—C8—C7	3.5 (3)
C2—C3—C4—Ru1	−31.22 (18)	Ru1—C1—C8—C7	−102.95 (16)
N1—Ru1—C4—C5	179.21 (11)	C2—C1—C8—Ru1	106.48 (16)
N2—Ru1—C4—C5	13.50 (12)	C6—C7—C8—C1	49.7 (2)
C1—Ru1—C4—C5	−100.05 (11)	C6—C7—C8—Ru1	−31.57 (18)
C8—Ru1—C4—C5	−66.49 (11)	N1—Ru1—C8—C1	12.33 (13)
Cl1—Ru1—C4—C5	−162.97 (11)	N2—Ru1—C8—C1	−178.08 (12)
Cl2—Ru1—C4—C5	95.25 (10)	C4—Ru1—C8—C1	−66.19 (11)
N1—Ru1—C4—C3	−60.44 (12)	C5—Ru1—C8—C1	−99.52 (12)
N2—Ru1—C4—C3	133.85 (12)	Cl1—Ru1—C8—C1	94.30 (10)
C1—Ru1—C4—C3	20.30 (12)	Cl2—Ru1—C8—C1	−167.41 (12)
C8—Ru1—C4—C3	53.86 (13)	N1—Ru1—C8—C7	132.64 (12)
C5—Ru1—C4—C3	120.34 (17)	N2—Ru1—C8—C7	−57.76 (13)
Cl1—Ru1—C4—C3	−42.6 (2)	C1—Ru1—C8—C7	120.32 (18)
Cl2—Ru1—C4—C3	−144.41 (12)	C4—Ru1—C8—C7	54.12 (13)
C3—C4—C5—C6	1.9 (3)	C5—Ru1—C8—C7	20.80 (13)
Ru1—C4—C5—C6	105.41 (17)	Cl1—Ru1—C8—C7	−145.38 (12)
C3—C4—C5—Ru1	−103.53 (17)	Cl2—Ru1—C8—C7	−47.1 (2)

Fig. 1

