

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/341026976>

Cardiorenal Effects of Pharmaceutical Plant Effluent in Mice (*Mus musculus*)

Article in *Nigerian journal of physiological sciences: official publication of the Physiological Society of Nigeria* · December 2019

CITATIONS

0

READS

29

5 authors, including:



Adam Olaitan Abdulkareem
University of Ilorin

13 PUBLICATIONS 31 CITATIONS

[SEE PROFILE](#)



Segun Olayinka Oladipo
Kwara State University

36 PUBLICATIONS 80 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Leachate [View project](#)



Pollution Assessment [View project](#)

Cardiorenal Effects of Pharmaceutical Plant Effluent in Mice (*Mus musculus*)

Abdulkareem A. O.^{1*}, Olafimihaan T. F.², Busari S. A.¹, Garuba A.³, Oladipo S. O.⁴

¹Animal Physiology Unit, Department of Zoology, University of Ilorin, Ilorin, Nigeria. ²Ecology and Environmental Biology Unit, Department of Zoology, University of Ilorin, Ilorin, Nigeria. ³Department of Zoology, University of Ilorin, Ilorin, Nigeria. ⁴Zoology Unit, Department of Bioscience and Biotechnology, Kwara State University, Malete, Kwara State, Nigeria

Summary: Many pharmaceutical industries carelessly handle their effluents and indiscriminately release same to aquatic environment. These effluents often find their ways into surface and ground waters, contaminating public water and thus, serving as a potential threat to animals and human health. In this study, we investigated the cardiorenal effects of chronic oral exposure to pharmaceutical effluent in mice. Thirty male mice (*Mus musculus*) were randomly divided into groups A-F and treated with 0.2 mLs 0.0 %, 2.5 %, 5.0%, 10.0%, 20.0% and 40% concentration (v/v, effluent/distilled water) of the effluent for 28 days, respectively. At the end of the experiment, the animals were sacrificed by cervical dislocation. Activities of alanine aminotransferase (ALT), aspartate aminotransferase (AST) and alkaline phosphatase (ALP) were determined in serum and heart homogenate, while uric acid, creatinine and electrolytes (sodium, potassium, bicarbonate and chloride ions) were determined in serum only. Data were expressed as Means $\pm$ standard error of mean and values were considered significant at $p < 0.05$. Results showed that, oral exposure to pharmaceutical effluent reduced ($p < 0.05$) cardiac ALP, AST and ALT activities as well as serum ALT activity. However, serum activities of ALP, creatinine and uric acid were elevated ($p < 0.05$). Similarly, there was derangement of electrolytes (potassium, chloride, bicarbonate and sodium ions) in the exposed mice, compared with control. This study has demonstrated that poorly treated pharmaceutical effluent disrupted cardiac and serum enzyme activities, caused electrolytes imbalance and elevated serum uric acid level, suggesting that, drinking water contaminated with pharmaceutical effluent may impair kidney and cardiac functions. Further study, investigating the histology of the kidney and heart of the pharmaceutical effluent-exposed animals as well as mechanism(s) of cardiorenal toxicity of the effluent, should be carried out to exploit its roles in pathogenesis of cardiorenal diseases.

Keywords: cardiorenal, pharmaceutical effluent; electrolytes; enzymes; *Mus musculus*

©Physiological Society of Nigeria

*Address for correspondence: abdulkareem.ao@unilorin.edu.ng. Phone number: +234(80)66528548

Manuscript Accepted: November, 2019

INTRODUCTION

Extensive industrialization, due to increasing population density and high urbanized societies, has resulted to global waste water management problem (Akpore et al., 2014). Many industries indiscriminately discharge contaminated and untreated waste products in to aquatic environment, causing pollution (Abu, 2012; Oyeniyi and Latunji, 2012). Inlands and estuary waters are both vulnerable to pollutants from industrial effluent discharge, affecting quality of water supply for domestic use (Osibanjo et al., 2011). This, therefore, causes global decrease in quality of drinking water below WHO standard and accounts for 3.1% death increase, globally (Khan et al., 2013; Pawari et al., 2015). It has also been reported that, heavy metals in water pollutants induce formation of reactive oxygen species (ROS), leading to organ damage in animals (Bando et al., 2005; Adegbesan and Adenuga, 2007).

Pharmaceuticals are among the important pollutants commonly released into environment (Stackelberg et

al., 2004). Pharmaceutical industries generally produce large quantity of wastes during manufacturing and maintenance operations, which are subsequently released into water bodies (Chander et al., 2014). These wastes often consist of toxic biodegradable and non-biodegradable substances, and therefore remain public health concern. Although, concentration of pharmaceuticals discharged into environment is drastically reduced by various physical and biological processes occurring in aquatic ecosystem, trace concentrations of human and veterinary pharmaceutical compounds as well as their metabolites have been detected in water bodies like surface, ground and drinking waters (Kolpin et al., 2002; Benotti and Brownawell, 2009; Bruce et al., 2010). Exposure to trace amount of pharmaceuticals for a long time can result in considerable adverse effects, such as tissue damages and inhibition of cell proliferation, in humans and aquatic life (Brooks et al., 2003; Pomati et al., 2006).

In Nigeria, there is continuous increase in pharmaceutical industries, mostly located in riverside

areas. The discharge practices of these industries are too crude, putting society in danger, especially in the industrialized parts of the country (Ijeoma and Achi, 2011). The industries carelessly handle their wastes and indiscriminately discharge their effluents into water bodies, promoting aquatic pollution and affecting human and environmental health (Osaigbovo and Orhue, 2006; Idris et al., 2013). This is largely due to their ignorance of harmful effects of xenobiotics and microbes present in the effluent on both aquatic and terrestrial life as well as poor enforcement of stringent regulations prohibiting illegal discharge of effluents (Lateef et al., 2007; Adeoye et al., 2015).

Previous studies have revealed the mutagenicity, genotoxicity and hepatotoxicity of pharmaceutical effluent (Zhao et al., 2007; Akintonwa et al., 2009; Bakare et al., 2009; Adeoye et al., 2015) but there is paucity of information on its cardio-renal toxicity. Recently, we showed that oral exposure to pharmaceutical effluent impairs cardiac Na^+/K^+ -ATPase activity and reduced cardiac weight index in mice (Abdulkareem et al., 2019). In this present study, we evaluated the effects of chronic oral exposure to pharmaceutical effluent on both renal and cardiac functions in mice.

MATERIALS AND METHODS

Effluent collection: The raw effluent from a pharmaceutical plant (which produces analgesics, anti-malarials, anesthetics, multivitamins and antibiotics) in Ilorin, Kwara State, Nigeria was collected in a 5L transparent plastic container (from the point of discharge into the environment). The collected effluent was filtered; the pH was taken and it was stored at 4°C until use.

Physico-chemical properties and heavy metal analysis: The effluent was analyzed for a number of standard physico-chemical properties, including: chemical oxygen demand (COD), total dissolved solids (TDS), alkalinity, biochemical oxygen demand (BOD), chlorides, nitrates, ammonia, and phosphates, according to methods described by APHA (1998). Eight metals which include: cadmium (Cd), chromium (Cr), iron (Fe), zinc (Zn), nickel (Ni), manganese (Mn), copper (Cu), and lead (Pb) were analyzed in the effluent sample according to standard analytical methods as previously reported (Bakare et al., 2009). Concentration of the metals were estimated by using an Atomic Absorption Spectrophotometer (Perkin Elmer E., Analyst, 2000, USA).

Animals and experimental design: Thirty male mice (*Mus musculus*) of 8-10 weeks old were obtained from Central Research Laboratory, University of Ilorin, Ilorin, Nigeria. They were kept in a clean, frequently disinfected and well-ventilated animal house of the Department of Zoology, University of Ilorin, for 2

weeks in order to acclimatize. All mice were maintained under standard condition (12h light: 12h dark) and were exposed to standard feed and drinking water *ad libitum*. Handling of animals was kept in accordance with the regulation of University of Ilorin Ethical committee and in conformity to the NIH Guidelines on the care and use of laboratory animals. The mice were divided into 6 groups. Group A (control) mice received 0.2ml distilled water, while groups B-F were treated with 0.2 mLs 2.5%, 5.0%, 10.0%, 20.0% and 40% concentration (v/v, effluent/distilled water) of the effluent, respectively. All the treatments were administered orally and lasted for 28 days.

Tissue preparation: On 29th post-treatment day, the animals were sacrificed by cervical dislocation. The heart was quickly excised, cleared of connective tissues, and transferred into the ice-cold 0.25M sucrose solution. The heart was homogenized in ice-cold 0.25M sucrose solution (1:5, w=v) as previously reported (Olatunji et al., 2006). The resulting homogenates were kept frozen overnight and centrifuged before use.

Enzymes assays: Method of Reitman and Frankel's (1957) was employed in estimating alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activities. Activity of alkaline phosphatase (ALP) was determined, using Babson et al. (1966) method.

Determination of serum uric acid and creatinine concentrations:

Concentrations of serum uric acid and creatinine were determined, using the method of Tietz (1994) as outlined in Randox kits, UK.

Determination of serum electrolytes: Potassium, sodium and chloride were determined by the method of Tietz et al. (1996), while serum bicarbonate level was assessed by the method of Roth and Chan (2001).

Statistical analysis

All data were expressed as means $\pm$ standard error of mean (SEM). Statistical analysis was performed with Graphpad Prism 5 (GraphPad Software, USA). Mean values of variables among the groups was compared by One-way analysis of variance (ANOVA), following Bonferroni Post-hoc test. Values were considered significant at $p < 0.05$.

RESULTS

Physicochemical properties of the raw pharmaceutical effluent: Details of the physicochemical properties of the effluent have been provided in our previous study (Abdulkareem et al., 2019). The pH value (6.40) of the effluent falls within national permissible limit (NESREA). Concentrations

of Cadmium (Cd), Chromium (Cr), Zinc (Zn), and Copper (Cu), phosphate, alkalinity and TDS were lower than international (USEPA) and national (NESREA) recommended limits, while iron (Fe), manganese (Mn) and ammonia (NH₃) in the sample were higher than the limits. Lead (Pb) and nickel (Ni) were below detectable limits.

Cardiac and serum activities of alanine aminotransferase (ALT), aspartate aminotransferase (AST) and alkaline phosphatase (ALP): As shown in figures 1, 2 and 3), analyses of enzyme activities showed that, 28 day oral administration of pharmaceutical effluent significantly decreased ($p < 0.05$) cardiac activities of ALP, AST and ALT. Similarly, there was a significant decrease ($p < 0.05$) in serum activity of ALT in the exposed

mice. In contrast, there was an increase ($p < 0.05$) in serum activity of ALP while serum activity of AST was not significantly changed ($p > 0.05$), when compared with control.

Serum creatinine and uric acid: Figure 4 shows that, chronic oral exposure to pharmaceutical effluent at 5 %, 10 %, 20 % and 40 % concentrations significantly ($p < 0.05$) elevated serum uric acid and creatinine levels, as compared with control.

Serum electrolytes: There was a significant ($p < 0.05$) concentration independent increase in serum potassium, chloride and bicarbonate ions values, meanwhile values of sodium ion decreased (concentration independent) significantly ($p < 0.05$) as compared with control (Table 1).



Fig. 1: Effect of pharmaceutical plant effluent on cardiac and serum alkaline phosphatase (ALP) activities in mice. Oral exposure to the effluent significantly decreased cardiac ALP activity, but increased the activity in the serum (* $p < 0.05$ vs control).



Fig. 2: Effect of pharmaceutical plant effluent on cardiac and serum aspartate aminotransferase (AST) activities in mice. Oral exposure to the effluent significantly decreased cardiac AST activity, whereas, there was no significant increase in serum activity of the enzyme (* $p < 0.05$ vs control).




Fig. 3: Effect of pharmaceutical plant effluent on cardiac and serum alanine aminotransferase (ALT) activities in mice. Oral exposure to the effluent significantly decreased cardiac ALT activity, and elevated serum activity of the enzyme (* $p < 0.05$ vs control).




Fig. 4: Effect of pharmaceutical plant effluent on serum uric acid and creatinine in mice. There was an increase in serum uric acid and creatinine levels at concentrations 5 %, 10 %, 20 % and 40 % (* $p < 0.05$ vs control).

Table 1:

Effect of pharmaceutical effluent on electrolytes

Treatment Groups	POTASSIUM	SODIUM	CHLORIDE	BICARBONATE
Control	4.63±0.89 _a	73.38±3.82 _a	13.07±0.46 _a	3.71±0.92 _a
2.5 %	9.60±0.72 _b	73.17±0.35 _a	15.31±2.87 _{ab}	17.94±9.32 _b
5 %	6.12±1.18 _{ab}	60.19±3.87 _b	21.82±2.70 _{bc}	17.66±3.50 _b
10 %	6.45±1.50 _{ab}	57.90±0.35 _b	24.51±2.25 _c	10.08±3.02 _{ab}
20 %	6.86±1.05 _{ab}	47.21±2.43 _c	16.81±1.92 _{ab}	11.72±7.24 _b
40 %	4.30±0.81 _a	58.35±2.01 _b	15.44±1.67 _{ab}	6.70±3.24 _a

Values along the same column with different superscripts are significantly different ($p < 0.05$)

DISCUSSION

The results of this study show that, chronic oral exposure to pharmaceutical effluent can induce cardiac and renal dysfunctions by disrupting enzyme activities, raising serum uric acid and by causing electrolytes imbalance. Activities of ALP, AST and ALT in both tissues and serum are important biomarkers in determining organ integrity and function (Arise et al., 2012; Shahjahan et al., 2004). More so, correlation of ALP with C-reactive protein, inflammation, obesity, and atherosclerosis makes it a potential diagnostic marker or predictor of cardiovascular disease (Webber et al., 2010). During organ damage, these enzymes are leaked into the blood, causing increase in their serum levels, with subsequent decreased activities in the respective organs (Arise et al., 2012). Therefore, decrease in cardiac ALP, AST and ALT with concomitant elevated level of serum ALP indicates that, heart membrane was damaged, after chronic exposure to pharmaceutical effluent, and probably resulting in cardiac dysfunction. Decrease in cardiac AST and ALT activities was equally followed by a decrease in serum activity of the enzymes. This may mean that, the decrease in cardiac enzymes activities observed, was due to antagonistic effect of the effluent on AST and ALT production (Akanji et al., 1993). Previous studies have associated elevated serum ALP level with coronary artery disease (CAD), as it promotes vascular calcification through pyrophosphate pathway (Johnson et al., 2006; Schoppet and Shanahan, 2008). Furthermore, high level of serum ALP increases risk

of mortality and unfavourable prognosis in coronary artery disease (Park et al., 2013; Wannamethee et al., 2013) as well as other cardiovascular diseases (CVDs) such as peripheral artery disease (Cheung et al., 2009), left ventricular hypertrophy (Nasri et al., 2004), secondary cardiac failure and diastolic dysfunction (Salgueira et al., 2005). Increase in serum ALP level observed in our study, therefore, suggests that, prolonged drinking of pharmaceutical effluent-contaminated water may promote the incidence of CVDs. Our finding is in consistence with our previous study (Abdulkareem et al., 2019) and the report of Karabulut et al. (2014).

Uric acid (UA) is a double bonded organic compound which, due to its antioxidant property, protects against oxidative stress (Sautin and Johnson, 2008). Elevated level of this compound however, may serve as an independent risk factor for coronary artery disease as it is frequently observed in patients with heart failure (Tian et al., 2012). Our results show that, in conjunction with elevated serum ALP level, oral exposure to pharmaceutical effluent at highest concentration (40 % v/v) elevated serum uric acid (SUA). This proposes further that, water contamination with pharmaceutical effluent may predispose an individual to cardiovascular diseases. Similarly, previous studies have reported increased SUA as an independent predictor of renal dysfunction in diseases such as rheumatoid arthritis and congestive heart failure (Daoussis et al., 2009; Tian et al., 2012). Therefore, the elevated SUA in our study may be a consequence of impaired renal function, which resulted to reduced excretion of UA. This thought is

supported further by concomitant increase in serum creatinine and electrolyte imbalance in our results (Figure 1a and Table 1).

Increase in serum creatinine in the exposed groups may be an indication of acute kidney injury, chronic kidney disease and renal dysfunction. Once creatinine is produced, it is immediately removed from the body by the kidney via urine. Since elimination of this molecule is solely by kidney, serum creatinine is widely used as a marker for renal function and readily employed in assessing acute kidney injury and chronic kidney disease (Winnett, et al., 2011; Baumgarten, 2011). Although, serum creatinine as a sole marker of kidney function has limitations; the observed electrolyte imbalance is a further confirmation of kidney dysfunction. The kidneys play a fundamental role in the regulation of body fluids and electrolytes; hence, impairment of kidney function results in derangement in electrolytes (Dhondup and Qian, 2017). Importantly, the observed hyperkalemia, hyperchloremia, hyponatraemia and elevated serum bicarbonate in the exposed animals have been earlier reported as predictors of chronic kidney disease and end-stage renal diseases (Dobre et al., 2015; Suetrong et al., 2016; Lim et al., 2016; Dhondup and Qian, 2017). Therefore, drinking water, contaminated with pharmaceutical effluent, may lead to development of chronic kidney disease and end-stage renal diseases. Interestingly, renal dysfunction has been noted to be a strong independent predictor of cardiovascular outcomes and mortality in the general population (Go et al., 2004), after myocardial infarction (Anavekar et al., 2004) and heart failure (Hillege et al., 2000). It can therefore be inferred from this study that, water contamination with pharmaceutical effluent may perhaps promote the pathogenesis of cardiovascular diseases via kidney function impairment.

The cardiorenal effects of pharmaceutical effluent observed in this study may be attributed to the combined effects of the chemical constituents. Chemicals such as Fe, Mn and NH₃ were found to be above permissible limit in our effluent sample. Meanwhile, iron overload has been reported to increase the fragility and density of renal lysosomes, thus, affecting their activities (Dimitriou et al., 2000). Since renal lysosomes play an important role in mediating kidney function of maintaining water-electrolyte homeostasis (Surendran et al., 2014), the disruption of serum electrolytes balance observed in this study may be due to renal iron accumulation. Similarly, cardiotoxicity and tissue damaging effect of Mn and NH₃, respectively, have been previously reported (Millera et al., 2006; McDaniel et al., 2016), hence, presence of these chemicals in the effluent probably resulted to impairment in cardiac enzymes activities.

The results of this study demonstrated that, drinking water, contaminated with pharmaceutical effluent,

may disrupt enzyme activities, cause electrolytes imbalance and elevate serum uric acid. Therefore, such water may independently predispose individuals to cardiorenal syndrome. Further study, investigating the histology of concerned organs and mechanism(s) of cardiorenal toxicity of pharmaceutical effluent, should be carried out to exploit its roles in pathogenesis of kidney and cardiovascular diseases.

REFERENCES

- Abdulkareem, A.O., Olafimihan, T.F., Akinbobola, O.O., Busari, S.A., Olatunji, L.A. (2019). Effect of pharmaceutical effluent on cardiac Na⁺-K⁺ ATPase and Ca²⁺-Mg²⁺-ATPase activities in mice (*Mus musculus*). *Toxicology reports*. 6:439-443
- Abu, N.E. (2012). Cytogenotoxicity effects of industrial effluents on *Allium cepa* root meristem: A review on positive results and problems of effective compliance to environmental legislations; the Nigeria perspective. *J. Toxicol. Environ. Health Sci.* 4 (10), 162-170.
- Adegbesan, B.O., Adenuga, G.A. (2007). Effect of lead exposure on liver lipid peroxidative and antioxidant defense systems of protein-undernourished rats. *Biol. Trace Element Res.* 116, 219–225.
- Adeoye, G.O., Alimba, C.G., Oyeleke, O.B. (2015). The genotoxicity and systemic toxicity of a pharmaceutical effluent in Wistar rats may involve oxidative stress induction. *Toxicol Rep* 2, 1265–1272.
- Akanji, M.A., Olagoke, O.A., Oloyede, O.B. (1993). Effect of chronic consumption of metabisulphite on the integrity of rat liver cellular system. *Toxicol.* 81, 173-179.
- Akintonwa, A., Awodele, O., Olofinnade, A.T., Anyakora, C., Afolayan, G.O., Coker, H.A.B. (2009). Assessment of the Mutagenicity of Some Pharmaceutical Effluents. *Am J Pharmacol Toxicol.* 4 (4), 144-150.
- Akpor, O.B., Okolomike, U.F., Olaolu, T.D., Aderiye, B.I. (2014). Remediation of polluted wastewater effluents Hydrocarbon Removal. *Trends Appl Sci Res.* 9, 160-173.
- Anavekar, N.S., McMurray, J.J., Velazquez, E.J., Solomon, S.D., Kober, L., Rouleau, J.L., White, H.D., Nordlander, R., Maggioni, A., Dickstein, K., Zelenkofske, S., Leimberger, J.D., Califf, R.M., Pfeffer, M.A. (2004). Relation between renal dysfunction and cardiovascular outcomes after myocardial infarction. *N Engl J Med.* 351, 1285–1295.
- Arise, R.O., Malomo, S.O., Lawal, M.M. (2012). *Comparative antimalarial and toxicological effects of artemisinin with methanolic extract of Carica papaya leaves and bark of Alstoniabroonai in animal models.* *Adv. Natl. Applied Sci.* 6, 116-123.
- Babson, L.A., Greeley, S.J., Coleman, C.M., Philips, G.D. (1966). Serum alkaline phosphatase determination. *Clin. Chem.* 12, 482-490.
- Bakare, A.A., Okunola, A.A., Adetunji, O.A., Jenmi, H.B. (2009). Genotoxicity assessment of a pharmaceutical effluent using four bioassays. *Genet Mol Biol.* 32 (2), 373-381.

- Bando, I., Reus, M.I., Andres, D., Cascales, M. (2005). Endogenous antioxidant defence system in rat liver following mercury chloride oral intoxication. *J. Biochem. Mol. Toxicol.* 19, 154–161.
- Baumgarten, M. (2011). Chronic Kidney Disease: Detection and Evaluation. *Am Fam Physician.* 84(10), 1138–1148.
- Benotti, M.J., Brownawell, B.J. (2009). Microbial degradation of pharmaceuticals in estuarine and coastal seawater *Environ Pollut.* 157 (3), 994–1002.
- Brooks, B.W., Turner, P.K., Stanley, J.K., Weston, J.J., Glidewell, E.A., Foran, C.M., Slattery, M., LaPoint, T.W., Huggett, D.B. (2003). Waterborne and sediment toxicity of fluoxetine to select organisms. *Chemosphere.* 52, 135–142.
- Bruce, G.M., Pleus, R.C., Snyder, S.A. (2010). Toxicological Relevance of Pharmaceuticals in Drinking water. *Environ. Sci. Technol.* 44, 5619–5626.
- Chander, M.P., Sachithanandam, V., Vijayachari, P. (2014). Antimicrobial and haemolytic activity of seaweed *Padina gymnospora* from South Andaman, Andaman and Nicobar Islands of India. *Int J Curr Microbiol Appl Sci.* 3, 364–69.
- Cheung, B.M., Ong, K.L., Wong, L.Y., 2009. Elevated serum alkaline phosphatase and peripheral arterial disease in the United States National Health and Nutrition Examination Survey 1999–2004. *Int J Cardiol.* 135, 156–61.
- Daoussis, D., Panoulas, V., Toms, T., John, H., Antonopoulos, I., Nightingale, P., Douglas, K.M.J., Klocke, R., Kitas, G.D. (2009). Uric acid is a strong independent predictor of renal dysfunction in patients with rheumatoid arthritis. *Arthritis Res Ther.* 11, R116.
- Dhondup, T., Qian, Q. (2017). Electrolyte and Acid–Base Disorders in Chronic Kidney Disease and End-Stage Kidney Failure. *Blood Purif.* 43, 179–188.
- Dimitriou, E., Kairis, M., Saraçdou, J., Michelakakis, H. (2000). Iron overload and kidney lysosomes. *Biochimica et Biophysica Acta.* 1501, 138–148.
- Dobre, M., Yang, W., Pan, Q.M., Appel, L., Bellovich, K., Chen, J., Feldman, H., Fischer, M.J., Ham, L.L., Hostetter, T., Jaar, G.B., Kallem, R.R., Rosas, S.E., Scialla, J.J., Wolf, M., Rahman, M. (2015). Persistent High Serum Bicarbonate and the Risk of Heart Failure in Patients With Chronic Kidney Disease (CKD): A Report From the Chronic Renal Insufficiency Cohort (CRIC) Study. *J Am Heart Assoc.* 4:e001599 doi: 10.1161/JAHA.114.001599.
- Go, A.S., Chertow, G.M., Fan, D., McCulloch, C.E., Hsu, C.Y. (2004). Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med.* 351, 1296–1305.
- Hillege, H.L., Girbes, A.R.J., de Kam, P.J., Boomsma, F., de Zeeuw, D., Charlesworth, A., Hampton, J.R., van Veldhuisen, D.J. (2000). Renal function, neurohormonal activation, and survival in patients with chronic heart failure. *Circulation.* 102, 203–210.
- Idris, M.A., Kolo, B.G., Garba, S.T., Ismail, M.A. (2013). Physico-chemical analysis of pharmaceutical effluent and surface water of River Gorax in Minna, Niger State, Nigeria. *Bull. Environ. Pharmacol. Life Sci.* 2, 45–49.
- Ijeoma, K., Achi, O.K. (2011). Industrial Effluents And Their Impact On Water Quality Of Receiving Rivers In Nigeria. *JATES,* 1 (1), 75–86.
- Johnson, R.C., Leopold, J.A., Loscalzo, J. (2006). Vascular calcification: pathobiological mechanisms and clinical implications. *Circ Res.* 99, 1044–1059.
- Karabulut, A., Sahin, I., Avci, I.I., Okuyan, E., Dogan, Z., Uzunlar, B., Satilmis, S. (2014). Impact of serum alkaline phosphatase level on coronary collateral circulation. *Kardiol Pol.* 72 (12), 1388–1393.
- Khan, N., Hussain, S.T., Saboor, A., Jamila, N., Kim, K.S. (2013). Physiochemical investigation of the drinking water sources from Mardan, Khyber Pakhtunkhwa, Pakistan. *Int. J. Phys. Sci.* 8 (33), 1661–1671.
- Kolpin, D.W., Furlong, E.T., Meyer, M.T., Thurman, E.M., Zaugg, S.D., Barber, L.B., Buxton, H.T. (2002). Pharmaceuticals, Hormones, and other Organic Wastewater Contaminants in U.S. Streams, 1999–2000: A National Reconnaissance. *Environ. Sci. Technol.* 36, 1202–1211.
- Lateef, A., Ufuoma, P.E., Yekeen, T.A. (2007). Bacteriology and genotoxicity of some pharmaceutical wastewaters in Nigeria. *Int. J. Environ. Health.* 1, 551–562.
- Lim, L.M., Tsai, N.C., Lin, M.Y., Hwang, D.Y., Lin, H.Y.H., Lee, J.J., Hwang, S.J., Hung, C.C., Chen, H.C. (2016). Hyponatremia is Associated with Fluid Imbalance and Adverse Renal Outcome in Chronic Kidney Disease Patients Treated with Diuretics. *Sci. Rep.* 6, 1–9.
- Mansour, H.B., Dellai, A., Ayed, Y., Shahjahan, M., Sabitha, K.E., Mallika J., Shyamala-Devi C.S. (2004). Effect of *Solanum trilobatum* against carbon tetrachloride induced hepatic damage in albino rats. *Indian J. Med Res.* 120, 194–198.
- McDaniel, J., Davuluri, G., Hill, E.A., Moyer, M., Runkana, A., Prayson, R., Lunteren, E.V., Srinivasan, D. (2016). Hyperammonemia results in reduced muscle function independent of muscle mass. *Am J. Physiol/Gastrointest Liver Physiol.* 310 (3), G163–G170.
- Millera, K.B., Catona, J.S., Finley, J.W. (2006). Manganese depresses rat heart muscle respiration. *Bio Factors.* 28, 33–46.
- Nasri, H., Baradaran, A., Naderi, A.S., 2004. Close association between parathyroid hormone and left ventricular function and structure in end-stage renal failure patients under maintenance hemodialysis. *Acta Med Austriaca.* 31, 67–72.
- Olatunji, L.A., Adebayo, A.O., Adesokan, A.A., Olatunji, V.A., Soladoye, A.O. (2006). Chronic administration of aqueous extract of *Hibiscus sabdariffa* enhances Na⁺-K⁺-ATPase and Ca²⁺-Mg²⁺-ATPase activities of rat heart. *Pharmaceutical biology,* 44 (3), 213–216.
- Osaigbovo, A.U., Orhue, E.R. (2006). Influence of pharmaceutical effluent on some soil chemical properties and early growth of maize (*Zea mays* L.). *Afr. J. Biotechnol.* 5, 1612–1617.
- Osibanjo, O., Daso, A.P., Gbadebo, A.M. (2011). The impact of industries on surface water quality of River Ona

- and River Alaro in Oluyole Industrial Estate, Ibadan, Nigeria. *Afr. J. Biotechnol.* 10 (4), 696-702.
- Oyeniyi, T., Latunji, C. (2012). Industrial effluents induced abnormal sperm cells in mice (*Mus musculus*). *N Y Sci J.* 5 (6), 60-64.
- Park, J.B., Kang, D.Y., Yang, H.M., Cho, H.J., Park, K.W., Lee, H.Y., Kang, H.J., Koo, B.K., Kim, H.S. (2013). Serum alkaline phosphatase is a predictor of mortality, myocardial infarction, or stent thrombosis after implantation of coronary drug-eluting stent. *Eur Heart J.* 34, 920-931.
- Pawari, M.J., Gawande, S. (2015). Ground water pollution & its consequence. *Int. j. eng. res. gen. sci.* 3 (4), 773-776.
- Pomati, F., Castiglioni, S., Zuccato, E., Fanelli, R., Vigetti, D., Rossetti, C., Calamari, D. (2006). Effects of a complex mixture of therapeutic drugs at environmental levels on human embryonic cells. *Environ. Sci. Technol.* 40 (7), 2442-2447.
- Reitman, S., Frankel, S. (1957). Determination of serum transaminases. *Amer J. Clin path.* 28, 56-59.
- Roth, K.S., Chan, J.C. (2001). Renal tubular acidosis: a new look at an old problem. *Clin Pediatr (Phila).* 40 (10), 533-543.
- Salgueira, M., Milan, J.A., Moreno, A.R., Amor, J., Aresté, N, Jiménez E, et al. Cardiac failure and diastolic dysfunction in hemodialysis patients: associated factors. *Nefrologia.* 25, 668-77.
- Sautin, Y.Y., Johnson, R.J. (2008). Uric acid: the oxidant-antioxidant paradox. *Nucleosides Nucleotides Nucleic Acids.* 27, 608-619.
- Schoppet, M., Shanahan, C.M. (2008). Role for alkaline phosphatase as an inducer of vascular calcification in renal failure. *Kidney Int.* 73, 989-991.
- Shahjahan M., Sabitha K.E., Jainu, M., Shyamala Devi, C.S. (2004). Effect of *Solanum trilobatum* against carbon tetrachloride induced hepatic damage in albino rats. *Indian J Med Res* 120 (3), 194-198.
- Stackelberg, P.E., Furlong, E.T., Meyer, M.T., Zaugg, S.D., Henderson, A.K., Reissman, D.B. (2004). Persistence of pharmaceutical compounds and other organic wastewater contaminants in a conventional drinking-water-treatment plant. *Sci Total Environ.* 329 (1-3), 99-113.
- Suetrong, B., Pisitsak, C., Boyd, J.H., Russell, J.A., Walley, K.R. (2016). Hyperchloremia and moderate increase in serum chloride are associated with acute kidney injury in severe sepsis and septic shock patients. *Crit Care.* 20 (315), 1-8.
- Surendran, K., Vitiello, S.P., Pearce1, D.A. (2014). Lysosome dysfunction in the pathogenesis of kidney diseases. *Pediatr Nephrol.* 29 (12), 2253-2261.
- Tian, Y., Chen, Y., Deng, B., Lui, G., Ji, Z.G., Zhao, Q.Z., Zhen, Y.Z., Gao, Y.Q., Tian, L., Wang, L., Ji, L.S., Ma, G.P., Liu, K.S., Liu, C. (2012). Serum uric acid as an index of impaired renal function in congestive heart failure. *J Geriatr Cardiol.* 9, 137-142.
- Tietz, N., Pruden, L.E., Andersen, S. (1996). Electrolytes. In: *Tietz Fundamentals of Clinical Chemistry*, 2nd Ed. WB Saunders Company, U.S.A., pp: 721-738.
- Tietz, N.W. (1994). *Textbook of Clinical Chemistry*. 2nd Edn. Burtis CA, Ashwood ER, W.B. Saunders Company, Philadelphia, pp. 751.
- Wannamethee, S.G., Sattar, N., Papcosta, O., Lennon, L., Whincup, P.H. (2013). Alkaline phosphatase, serum phosphate, and incident cardiovascular disease and total mortality in older men. *Arterioscler Thromb Vasc Biol.* 33, 1070-1076.
- Webber, M., Krishnan, A.S., Thomas, N.G., Cheung, B.M.Y., 2010. Association between serum alkaline phosphatase and C-reactive protein in the United States National Health and Nutrition Examination Survey 2005-2006. *Clin Chem Lab Med.* 48(2), 167-173.
- Winnett, G., Cranfield, L., Almond, M. (2011). Apparent renal disease due to elevated creatinine levels associated with the use of boldenone. *Nephrol Dial Transplant.* 26, 744-747.
- Zhao, D., Zhu, C., Sun, S., Yu, H., Zhang, L., Pan, W., Zhang, X., Yu, H., Gu, J., Cheng, S. (2007). Toxicity of pharmaceutical wastewater on male reproductive system of *Mus musculus*. *Toxicol Ind Health.* 23: 47-54.