

Volume 6, Numbers 1 & 2, December, 2019



**A Journal of Research
and Development**



Produced with the Support of
Tertiary Education Trust Fund, Nigeria

© A. Uwaherem



ISSN: 2276-9595

✓

*ASUU JOURNAL OF SCIENCE,
A Journal of Research and Development*

Prof. S.S. Noma

Editor –in- Chief
Usmanu Danfodiyo University, Sokoto, Nigeria
E-mail: samailanoma@gmail.com,
+234 7037777273

Prof. Joseph Onch Abu, FNIST

Managing Editor
Federal University of Agriculture, Makurdi, Nigeria
E-mail: joabuza@yahoo.co.uk, +2348056602892

Prof. C.O. Ofor

Business Editor
Micheal Okpara University of Agriculture,
Umudike, Nigeria
E-mail: chukofofor@gmail.com, +2348035472301

Prof. Francisca Bassey

Associate Editor
University of Calabar, Calabar, Nigeria
fibassey@yahoo.ca, +2348036567632

Consulting Editors

Prof. K. Abdullahi

Federal University, Gusau, Zamfara
State, Nigeria

Prof. Tunde Ochende-Bello

University of Cape Town, Cape
Town, South Africa

Prof. U. A. Ali

Federal University, Dutse, Jigawa
State, Nigeria

Prof. E. Okunade

ObafemiAwolowo University, Ile-
Ife, Nigeria

Dr. A. Singh

The University of Nottingham, Malaysia Campus

Prof. A.R. Isara

University of Benin, Benin, Edo State, Nigeria

Editorial Advisers

Prof. Blodun Ogunyemi

Olabisi Onabanjo University, Ago-Iwoye, Ogun
State, Nigeria

Dr. Nasir F. Isa

Prof. A.A. Ahmed

Prof. Ikenna Onyido

Bayero University, Kano, Nigeria
Ahmadu Bello University, Zaria, Nigeria
Nnamdi Azikiwe University, Awka,
Anambra State, Nigeria

sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or PowerPoint before pasting in the Microsoft Word manuscript file. Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by *et al.* except where newer versions of APA referencing styles are adopted in line with global best practices. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

SUBMISSIONA covering letter should include the corresponding author's full address, e-mail and telephone and WhatsApp enabled number(s). Softcopies of the manuscripts should be sent to following email address: thesciencejournal@gmail.com and copied to: thesciencejournal@asuu.org.ng

For further enquiries please call:

Professor S.S. Noma, FSSSN
Editor-in-Chief
Usmanu Danfodiyo University, Sokoto
E-mail: samailanoma@gmail.com
Phone: +234 7037777273

Prof. Joseph Oneh Abu, FNIFST
Managing Editor
Federal University of Agriculture, Makurdi,
Nigeria, +2348056602892
E-mail: joabuza@yahoo.co.uk

Contents

1. The Use of Progressive Operating Characteristic Values for Gamma and Weibull Distributions on Designing Acceptance Sampling Plan
Zakari, Y. and Muhammad, I. 1
2. Exponentially Fitted Collocation Method for Solving Singular Multi-order Fractional Integro Differential Equations
Owolanke A.O. , Uweheren O.A. , And Taiwo O.A. 10
3. Acceptance Sampling Plan Based on Percentiles for Tsallis q-Exponential Distribution
Zoramawa, A. B. and S. Bhanu Prakash 24
4. Hepatic Antioxidant Status and Organ indices of *Oreochromis niloticus* Exposed to Sub-lethal Concentrations of *Anacardium occidentale* Linn; Aqueous Bark Extract
Ugbor, O. N., Oke, A. O., Okpokwu, U. J. and Ogbuagu, C. U. Anosike, G. E. 35
5. Impact of Aqueous Leaf Extract of Coffee Weed (*Senna occidentalis*) on the Blood Profile of Broiler Chickens
Omoikhoje, S.O, Uwaya, J.I, Adamu, I.A. and Idahor, E.F. 51

6. *In Vitro* Comparative Inhibitory activities of Extracts of *Annona muricata* on α -amylase and α -glucosidase for possible anti-diabetic remedy
Agu K. C., Eluehike N., Aparo P., Ima-Obakpolor S., T. Adetula 68
7. Effects of Cooking Methods on the Quality Attributes of Ripe Plantain Fruit
Egwujeh, S. I. D. and Yusufu A. 90
8. Safety Assessment of Drinking Water in Anyigba Town, Kogi State, Nigeria
Yusufu, P. A and Egwujeh, S.I.D. 105
9. Examining the Determinants of Perceived Agricultural-Land Degradation Among Farming Households in Sokoto Plain, North-Western Nigeria
Abdullahi, A.N, Abdulsalam, Z., Rahman, S. A, Atala, T.K. and Oladimeji, Y. U 118
10. Effect of Harvesting Methods and Fruit Size on Physiological quality of Mango (*Mangifera indica* L.) Fruits During Storage
Ahmed, M.,Gungula, D.T.,Tame, V.T. 143

Exponentially Fitted Collocation Method For Solving Singular Multi-order Fractional Integro Differential Equations

Owolanke, A. O.,¹ Uweheren, O. A.² and Taiwo, O. A.¹

Abstract

This work considered the construction of canonical polynomials and used as basis functions for the approximation of singular multi-order fractional integro-differential equations. The idea is that the singular multi-order problem is slightly perturbed with shifted Chebyshev polynomials, and the resulting equation is collocated at equally spaced interior points. The conditions are exponentially fitted with one tau-parameter along with the unknown constants. This results into a system of linear algebraic equations which are then solved using Gaussian elimination method to obtain the unknown parameters involved. Some examples are solved to demonstrate the effectiveness of the method.

Keywords : Canonical Polynomials, Perturbed Collocation Method, Fractional Integro-Differential Equations.

Owolanke, A. O.,¹ Uweheren, O. A.² and Taiwo, O. A.¹
¹Department of Mathematics, Olusegun Agagu University of Science and Technology, Okitipupa, PMB 353, Ondo State, Nigeria.
²Department of Mathematics, University of Ilorin, Ilorin, Kwara State, Nigeria.
Corresponding author Email: owolankeayo@yahoo.com

highly efficient method of solution. For instance, Taiwo (2000) investigated the numerical solution of two point boundary value problems by the application of the Cubic Spline Collocation Method using exponential fitting. Among other authors that have also adopted the technique as method of solution are Raptis (1982), Simos (1999), Simos (2006), and Daele and Berghe (2007). Thus, in this work, the exponential fitting with collocation method is introduced to solve equation (1), implementing the constructed canonical polynomials as basis functions.

Definition of Relevant Terms

Definition 1.1: Riemann-Liouville Fractional derivative

Riemann Liouville fractional derivative denoted by $D_{-0^+}^q f(t)$ is defined as Follows

$$D_{-0^+}^q f(t) = D_{-0^+}^m D_{-0^+}^{q-m} f(t) = D_{-0^+}^m I_{-0^+}^{m-q} f(t); \quad q > 0$$

$$= \frac{1}{\Gamma(m-q)} \frac{d^m}{dx^m} \int_0^x (x-\tau)^{m-q-1} f(\tau) d\tau, \quad q > 0, m-1 < q < m \quad (2)$$

where $D^m = \frac{d^m}{dx^m}$ is the derivative part and $D_{-0^+}^{q-m} f(x) = I_{-0^+}^{m-q} f(x)$ is the integral part.

Definition 1.2: Caputo Fractional Derivative

Let $m-1 < q < m$, and $q < 0$. Caputo fractional derivative denoted by

$D_{-0^+}^q f(x)$ is defined as follows

$$D_{-0^+}^q f(x) = D_{-0^+}^{q-m} [D^m f(x)] = I_{-0^+}^{m-q} [D^m f(x)]; \quad m-1 < q < m$$

$$\begin{cases} \frac{1}{\Gamma(m-q)} \int_0^x (x-\tau)^{m-q-1} [D^m f(\tau)] d\tau, & m-1 < q < m \\ \frac{d^m}{dx^m} f(x); & q = m, \end{cases} \quad m \in \mathbb{N} \quad (3)$$

Definition 1.3: Chebyshev Polynomials

The Chebyshev polynomials belong to a family of orthogonal polynomials in the interval $[-1, 1]$. They are widely utilized for their good properties in approximating problems. The Chebyshev polynomial of the first kind of degree n denoted by $T_n(x)$ and valid in $[-1, 1]$ is defined as

$$T_n(x) = \cos(ncos^{n-1}x)$$

$$\text{Let } \theta = \cos^{-1}x \quad (4)$$

where $x = \cos\theta$, and the recurrence relation is given as

$$T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x), \quad n \geq 1 \quad (5)$$

But for the purpose of this work, Chebyshev polynomials valid in interval $[a, b]$ is defined as

$$T_n(x) = \cos \left\{ n \cos^{-1} \frac{2x - (a+b)}{b-a} \right\}, \quad n \geq 0$$

$$T_n(x) = \cos \left\{ n \cos^{-1} \frac{2x - (a+b)}{b-a} \right\} T_n(x) - T_{n-1}(x) \quad (6)$$

Methodology

In this section, a general singular multi-order Volterra fractional integro-differential equation of the form:

$$D^\alpha y(x) + \sum_{i=0}^n p_i y^{(i)}(x) + \lambda \int_0^x \frac{y(t)}{\sqrt{x-t}} dt = f(x), \quad a \leq x \leq b \quad (7)$$

$$y^{(k)}(0) = \alpha_k, \quad k = 0, 1, \dots, n-1 \quad (8)$$

Equation (7) is equivalently written as

$$D^\alpha y(x) + \sum_{i=0}^n p_i y^{(i)}(x) = f(x) - \lambda \int_0^x \frac{y(t)}{\sqrt{x-t}} dt, \quad a \leq x \leq b \quad (9)$$

and equation (9) is re-written as

$$D^\alpha y(x) + \sum_{i=0}^n p_i y^{(i)}(x) = g(x) \quad (10)$$

where,

$$g(x) = f(x) - \lambda \int_0^x \frac{y(t)}{\sqrt{x-t}} dt$$

λ and p_i ($i \geq 0$) are constant parameters. Also, $y(x)$ and $g(x)$ are functions of x , where x and t are independent variables.

Construction of Canonical Polynomials

Canonical polynomials basis function constructed from equation (7) is obtained as follows:

Expanding equation (10) derived from the general singular and multi-order fractional order integro-differential equations stated in equations (7) and (8), gives

$$(D^\alpha + p_0 + p_1 \frac{d}{dx} + p_2 \frac{d^2}{dx^2} + \dots + p_n \frac{d^n}{dx^n})y(x) = g(x) \quad (11)$$

such that an operator denoted by L is defined as follows.

$$L = D^\alpha + p_0 + p_1 \frac{d}{dx} + p_2 \frac{d^2}{dx^2} + \dots + p_n \frac{d^n}{dx^n} \quad (12)$$

or simply:

$$L \equiv D^\alpha + \sum_{i=0}^n p_i \frac{d^i}{dx^i} \quad (13)$$

Following Lanczos (1956), we defined canonical polynomials $Q_j(x)$, $j \geq 0$, which are uniquely associated with the operator L by

$$LQ_j(x) = x^j, j = 0, 1, 2, 3, \dots \quad (14)$$

From equation (12), it follows that:

$$Lx^j = \frac{\Gamma(j+1)}{\Gamma(j+1-\alpha)} x^{j-\alpha} + p_0 x^j + p_1 j x^{j-1} + p_2 j(j-1) x^{j-2} + \dots + p_n j(j-1)(j-2) \dots (j-n+1) x^{j-n} \quad (15)$$

This implies,

$$L[LQ_j(x)] = \frac{\Gamma(j+1)}{\Gamma(j+1-\alpha)} LQ_{j-\alpha}(x) + p_0 LQ_j(x) + p_1 j LQ_{j-1}(x) + p_2 j(j-1) LQ_{j-2}(x) + \dots + p_n j(j-1)(j-2) \dots (j-n+1) LQ_{j-n}(x) \quad (16)$$

Hence, taking L^{-1} of both sides of equation (16), leads to

$$L^{-1}L[LQ_j(x)] = \frac{\Gamma(j+1)}{\Gamma(j+1-\alpha)} L^{-1}LQ_{j-\alpha}(x) + p_0 L^{-1}LQ_j(x) + p_1 j L^{-1}LQ_{j-1}(x) + p_2 j(j-1) L^{-1}LQ_{j-2}(x) + \dots + p_n j(j-1)(j-2) \dots (j-n+1) L^{-1}LQ_{j-n}(x) \quad (17)$$

which implies,

$$x^j = \frac{\Gamma(j+1)}{\Gamma(j+1-\alpha)} Q_{j-\alpha}(x) + p_0 Q_j(x) + p_1 j Q_{j-1}(x) + p_2 j(j-1) Q_{j-2}(x) + \dots + p_n j(j-1)(j-2) \dots (j-n+1) Q_{j-n}(x) \quad (18)$$

Thus,

$$Q_j(x) = \frac{1}{p_0} \left[x^j - \frac{\Gamma(j+1)}{\Gamma(j+1-\alpha)} Q_{j-\alpha}(x) - p_0 Q_j(x) + p_1 j Q_{j-1}(x) - p_2 j(j-1) Q_{j-2}(x) - \dots - p_n j(j-1)(j-2) \dots (j-n+1) Q_{j-n}(x) \right], j \geq 0 \quad (19)$$

Method of Solution

The perturbed Collocation Method with exponential fitting is considered in this section, as a new method to handle singular and multi-order fractional integro-differential equations.

First, an assumed approximate solution of the form:

$$y(x) \approx y_N(x) = \sum_{j=0}^N a_j Q_j(x), \quad j \geq 0 \quad (20)$$

is considered, where, $a_j (j \geq 0)$ are constants to be determined, $Q_j (j \geq 0)$ are canonical polynomials constructed in equation (19) and N is the degree of the approximant. Thus, equation (20) is substituted into a slightly perturbed equation (7) to give

$$D^\alpha y_N(x) + \sum_{i=0}^n p_i y_N^{(i)}(x) + \lambda \int_a^x \frac{y_N(t)}{\sqrt{x-t}} dt = f(x) + H_N(x), \quad a \leq x \leq b \quad (21)$$

where $H_N(x)$ is the given perturbation term

$$H_N(x) = \sum_{i=0}^{n-1} \tau_i T_{N-i+1}(x) \quad (22)$$

and $T_{N-i+1}(x)$ is the shifted Chebyshev polynomial defined in equation (6). Thus, equation (22) is substituted into equation (21) and further simplification gives

$$\begin{aligned} D^\alpha \left(\sum_{j=0}^N a_j Q_j(x) \right) + p_0 \sum_{j=0}^N a_j Q_j(x) + p_1 \sum_{j=0}^N a_j Q_j'(x) + p_2 \sum_{j=0}^N a_j Q_j''(x) + \dots \\ + p_n \sum_{j=0}^N a_j Q_j^{(n)}(x) + \lambda \int_a^x \frac{y_N(t)}{\sqrt{x-t}} dt = f(x) + \tau_1 T_N(x) + \tau_2 T_{N-1}(x) + \\ \tau_3 T_{N-2}(x) + \dots + \tau_{n-1} T_{N-i+2}(x) \end{aligned} \quad (23)$$

where, N is the degree of approximation and n is the highest order singular and multi-order fractional integro differential equation, $\tau_{i,i=1(1)n-1}$ are the free Tau parameters to be determined. Hence, equation (23) is further simplified to give

$$\begin{aligned}
& \{D^a Q_0(x) + p_0 Q_0(x) + p_1 Q'_0(x) + \dots + p_n Q_0^{(n)}(x)\} a_0 + \\
& \{D^a Q_1(x) + p_0 Q_1(x) + p_1 Q'_1(x) + \dots + p_n Q_1^{(n)}(x)\} a_1 + \\
& \vdots \\
& \{D^a Q_N(x) + p_0 Q_N(x) + p_1 Q'_N(x) + \dots + p_n Q_N^{(n)}(x)\} a_N + \\
& \lambda \int_0^x \frac{y(t)}{\sqrt{x-t}} dt = f(x) + \tau_1 T_N(x) + \tau_2 T_{N-1}(x) + \tau_3 T_{N-2}(x) \\
& + \dots + \tau_{n-1} T_{N-l+2}(x)
\end{aligned}
\tag{24}$$

The $y(t)$ in equation (24) is evaluated by adopting a Taylor series expansion of appropriate order.

$$\begin{aligned}
& \{D^a Q_0(x) + p_0 Q_0(x) + p_1 Q'_0(x) + \dots + p_n Q_0^{(n)}(x)\} a_0 + \\
& \{D^a Q_1(x) + p_0 Q_1(x) + p_1 Q'_1(x) + \dots + p_n Q_1^{(n)}(x)\} a_1 + \\
& \vdots \\
& \{D^a Q_N(x) + p_0 Q_N(x) + p_1 Q'_N(x) + \dots + p_n Q_N^{(n)}(x)\} a_N + \\
& \lambda \int_0^x \frac{1}{\sqrt{x-t}} \left\{ y_N(x) + (t-x) y'_N(x) + \frac{(t-x)^2}{2!} y''_N(x) + \frac{(t-x)^n}{n!} y_N^{(n)}(x) \right\} dt = \\
& + \tau_1 T_N(x) + \tau_2 T_{N-1}(x) + \tau_3 T_{N-2}(x) + \dots + \tau_{n-1} T_{N-l+2}(x)
\end{aligned}
\tag{25}$$

where

$$\begin{aligned}
 y_N(x) &= u_0 Q_0(x) + u_1 Q_1(x) + \dots + u_N Q_N(x) \\
 y'_N(x) &= u_0 Q'_0(x) + u_1 Q'_1(x) + \dots + u_N Q'_N(x) \\
 &\vdots \\
 y_N^{(n)} &= u_0 Q_N^{(n)}(x) + u_1 Q_N^{(n)}(x) + \dots + u_N Q_N^{(n)}(x)
 \end{aligned}
 \quad (26)$$

Hence, substituting equation (26) into equation (25) and simplifying gives

$$\begin{aligned}
 &[D^\alpha Q_0(x) - p_0 Q_0(x) + p_1 Q'_0(x) + \dots + p_N Q'_N(x) + 2\lambda x^{\frac{3}{2}} Q_0(x) - \\
 &\frac{2}{3}\lambda x^{\frac{3}{2}} Q'_0(x) + \frac{2}{10}\lambda x^{\frac{3}{2}} Q''_0(x) - \frac{2}{42}\lambda x^{\frac{3}{2}} Q'''_0(x)]u_0 + [D^\alpha Q_1(x) + p_1 Q_1(x) + \\
 &p_1 Q'_1(x) + \dots + p_N Q'_N(x) + 2\lambda x^{\frac{3}{2}} Q_1(x) - \frac{2}{3}\lambda x^{\frac{3}{2}} Q'_1(x) + \frac{2}{10}\lambda x^{\frac{3}{2}} Q''_1(x) - \\
 &\frac{2}{42}\lambda x^{\frac{3}{2}} Q'''_1(x)]u_1 + \dots + [D^\alpha Q_N(x) + p_0 Q_N(x) + p_1 Q'_N(x) + \dots + \\
 &p_N Q'_N(x) + 2\lambda x^{\frac{3}{2}} Q_N(x) - \frac{2}{3}\lambda x^{\frac{3}{2}} Q'_N(x) + \frac{2}{10}\lambda x^{\frac{3}{2}} Q''_N(x) + \frac{2}{42}\lambda x^{\frac{3}{2}} Q'''_N(x)]u_N \\
 &f(x) + \tau_1 T_N(x) + \tau_2 T_{N-1}(x) + \tau_3 T_{N-2}(x) + \dots + \tau_{N-1} T_{N-1+2}(x)
 \end{aligned}
 \quad (27)$$

Equation (27) is collocated at point $x = x_i$ to obtain,

$$\begin{aligned}
 &[D^\alpha Q_0(x_i) + p_0 Q_0(x_i) + p_1 Q'_0(x_i) + \dots + p_N Q'_N(x_i) + 2\lambda x_i^{\frac{3}{2}} Q_0(x_i) - \\
 &\frac{2}{3}\lambda x_i^{\frac{3}{2}} Q'_0(x_i) + \frac{2}{10}\lambda x_i^{\frac{3}{2}} Q''_0(x_i) - \frac{2}{42}\lambda x_i^{\frac{3}{2}} Q'''_0(x_i)]u_0 + [D^\alpha Q_1(x_i) + p_0 Q_1(x_i) + \\
 &p_1 Q'_1(x_i) + \dots + p_N Q'_N(x_i) + 2\lambda x_i^{\frac{3}{2}} Q_1(x_i) - \frac{2}{3}\lambda x_i^{\frac{3}{2}} Q'_1(x_i) + \frac{2}{10}\lambda x_i^{\frac{3}{2}} Q''_1(x_i) - \\
 &\frac{2}{42}\lambda x_i^{\frac{3}{2}} Q'''_1(x_i)]u_1 + \dots + [D^\alpha Q_N(x_i) + p_0 Q_N(x_i) + p_1 Q'_N(x_i) + \dots + \\
 &p_N Q'_N(x_i) + 2\lambda x_i^{\frac{3}{2}} Q_N(x_i) - \frac{2}{3}\lambda x_i^{\frac{3}{2}} Q'_N(x_i) + \frac{2}{10}\lambda x_i^{\frac{3}{2}} Q''_N(x_i) + \frac{2}{42}\lambda x_i^{\frac{3}{2}} Q'''_N(x_i)]u_N - \\
 &f(x_i) + \tau_1 T_N(x_i) + \tau_2 T_{N-1}(x_i) + \tau_3 T_{N-2}(x_i) + \dots + \tau_{N-1} T_{N-1+2}(x_i)
 \end{aligned}
 \quad (28)$$

where,

$$x_i = a + \frac{(b-a)i}{N+2}, \quad i=1, 2, \dots, N+1$$

the perturbed collocation method with exponential fitting, an approximation to the conditions is sought by adding $\tau_n e^a$ to equation (8); that is

$$y_N'' = \tau_n e^a \quad (29)$$

and

$$y_N^{(k)}(a) + \tau_n e^a = \alpha_k; \quad k = 0, 1, \dots, n-1 \quad (30)$$

is obtained. Thus, equations (28) and (30) gives $(N+n+1)$ algebraic equations in $(N+n+1)$ unknown constants $a_0, a_1, a_2, \dots, a_N, \tau_1, \tau_2, \dots, \tau_n$ to obtain a single polynomial approximation

$$y(x) \approx y_N(x) + y_N''(x); \quad a \leq x \leq b, \quad (31)$$

Therefore, equation (31) is the required approximation solution.

Numerical Experiments and Discussion

To illustrate the performance of the presented method, two numerical examples are considered in this section.

Problem 1: Consider a singular multi-order fractional integro-differential equation of the form

Problem 1: Consider a singular multi-order fractional integro-differential equation of the form

$$y'' + \frac{d^{1.5} y(x)}{dx^{1.5}} - \int_0^x \frac{y(t)}{\sqrt{x-t}} dt = 2.2256758334x^{0.5} + 7.221636669x^{1.5} - \frac{16}{15}x^{\frac{5}{2}} - \frac{256}{315}x^{\frac{7}{2}} + 2 + 12x^3, \quad 0 \leq x \leq 1 \quad (32)$$

subject to the conditions:

$$y(0) = 0, y'(0) = 0 \quad (33)$$

The exact solution is given as $y(x) = x^4 + x^2$

Problem 2: Consider a singular multi-order fractional integro-differential equation of the form

$$y''' + \frac{d^{2.75}y(x)}{dx^{2.75}} + xy(x) + \int_0^x \frac{y(t)}{\sqrt{x-t}} dt = 4 + 12x^2 + \frac{32}{35}x^2 + 4.419575909x^{0.75},$$

$$0 \leq x \leq 1 \quad (34)$$

subject to the conditions

$$y(0) = 0, y'(0) = 0, y''(0) = 0 \quad (35)$$

The exact solution is $y(x) = x^3$.

Tables of Results and Graphical Representations

Table (i) :

Absolute Errors of Problem 1 for case $N = 8$

x	Exact Solution	Approximate Solution	Absolute Error
0.000	0.00000000	0.000000000	0.0000000
0.100	0.01010000	0.010099998	.1291E -09
0.200	0.04160000	0.041599998	2.0010E -09
0.300	0.09810000	0.098099973	2.6614E -08
0.400	0.18560000	0.185599959	4.0101E -08
0.500	0.31250000	0.312499957	4.2119E -08
0.600	0.48960000	0.489599950	4.9991E -08
0.700	0.73010000	0.730099950	5.0010E -08
0.800	1.04960000	1.049599581	4.1935E -07
0.900	1.46610000	1.466099500	4.9989E -07
1.000	2.00000000	1.999999437	5.6292E -07

Table (ii):
Absolute Errors of Problem 2 for case $N = 10$

x	Exact Solution	Approximate Solution	Absolute Error
0.000	0.0000000	0.000000000	0.0000000
0.100	0.0010000	0.000999999	5.8350E - 11
0.200	0.0080000	0.007999999	5.9828E - 10
0.300	0.0270000	0.026999999	6.1135E - 10
0.400	0.0640000	0.063999999	6.3596E - 10
0.500	0.1250000	0.124999992	7.5421E - 09
0.600	0.2160000	0.215999992	7.9915E - 09
0.700	0.3430000	0.342999991	8.3044E - 09
0.800	0.5120000	0.511999997	2.3153E - 09
0.900	0.7290000	0.728999998	1.8460E - 09
1.000	1.0000000	0.999999962	3.7380E - 08




Figure 1: The Graph of the Approximate and Exact Solution for Problem 1




Figure 2: The Graph of the Approximate and Exact Solution for Problem 2

Discussion of Results

It is required to obtain approximate solutions of singular and multiorder fractional integro-differential equations due to difficulties in obtaining the analytical solutions. Thus, exponentially fitted collocation method by canonical polynomials is considered as the method of solution in this work, and it is revealed that it's highly efficient and effective in terms of convergence as it's clearly seen that the approximate solution converges rapidly to the exact solution as the degree of the approximant N increases.

References

- Avipsita, C., Uma, B. & Manda, B.N., (2017). Numerical solution of Volterra type fractional order integro-differential equations in Bernstein polynomials. *Journal of Applied Mathematical Sciences* 11(6); 249-264.
- Cheney, E. W & Kincaid, D. R., (2008). Numerical mathematics and computing. *Cengage Learning, Boston*.
- Daele, M., & Berghe, G. (2007). P-stable Obrechhoff methods of arbitrary order for second-order differential equations. *J. of Numer. Algorithms*, 44; 115 – 131
- Kumar, M. & Singh, N. (2010). Modified Adomian decomposition method and computer implementation for solving singular boundary value problems arising in various physical problems. *Journal of Comput Chem Eng*; 34(11): 1750 – 1760
- Lanczos, C. (1956). *Applied analysis* (New Jersey: Prentice Hall)
- Raptis, A. D. (1982). Two-step methods for the numerical solution of the Schrodinger equation, *J. Computing* 28; 373 - 378.
- Simos, T. E., (1999). An exponentially Fitted Eighth-Order Method for the Numerical Solution of the Schrodinger Equation. *Journal of Computational and Applied Mathematics*, 108; 177-194
- Simos, T. E. (2006). A Four-Step Exponentially Fitted Method for the Numerical Solution of the Schrodinger Equation. *Journal of Mathematical Chemistry*, 40(3); 305318

- Taiwo, O. A. (2000). Exponential Fitting for the Solution of Two point Boundary Value Problems with Cubic Spline Collocation Tau Method. *International Journal of Computer Math*, 79(3), 299 – 306.
- Taiwo, O.A., Jimoh, A.K. & Bello, A.K., (2014). Comparison of some numerical methods for the solution of first and second orders linear integro-differential equations. *American Journal of Engineering Research*, 3(1); 245-250.
- Wang, Y. & Li, Z. (2017). Solving nonlinear Volterra integro-differential equations of fractional orders by Euler Wavelet method. *Open Access Journal*, 2017: 27.
- Wang, Y., Li, Z. and Zhi, W. (2018). Fractional order Euler functions for solving fractional integro-differential equations with weakly singular kernel. *Open Access Journal*, 2018: 254
- Yahaya, Q. H., & Liu, M. Z. (2008). Modified Adomian Decomposition Method for Singular Initial Value Problems. *Journal of Surv. Math. Appl.*, 3; 183 - 193.
- Yasir, K. & Zdenek, S. (2012). Solving Certain Classes of LaneEmden Type Equations Using the Differential Transformation Method. *Advances in difference equations*, 174; 1687-1699.