

3/16/2021

Ribarstvo

Croatian Journal of Fisheries

SCIENTIFIC AND TECHNICAL JOURNAL IN FISHERIES

since 1938

Croatian Journal of Fisheries Volume: 72 Issue: 1

ISSN 1848-05

(online)

ISSN 1330-06

(print)

E-mail address

Password

☐ Stay signed in

sign in

[Home](#) [Search](#) [Volumes](#) [Publishers](#) [Links](#) [Users](#)


search

search




Volumes

2021	Volume 79
	Issue 1
2020	Volume 78
	Issue 4
	Issue 3
	Issue 2
	Issue 1
2019	Volume 77
	Issue 4
	Issue 3
	Issue 2
	Issue 1
2018	Volume 76
	Issue 4
	Issue 3
	Issue 2
	Issue 1
2017	Volume 75
	Issue 4
	Issue 3
	Issue 2
	Issue 1
2016	Volume 74
	Issue 4
	Issue 3
	Issue 2
	Issue 1
2015	Volume 73
	Issue 4
	Issue 3
	Issue 2
	Issue 1
2014	Volume 72
	Issue 4
	Issue 3
	Issue 2
	Issue 1
2013	Volume 71
	Issue 4
	Issue 3

Croatian Journal of Fisheries Volume: 72 Issue: 1

Volume: 72 Issue: 1

Zagreb, 2014

[show article ...](#)

EDITORIAL

Editorial 2014

Piria, M., Pranjić, M.

ORIGINAL SCIENTIFIC PAPER

[show article ...](#)

GENETIC VARIABILITY IN CULTURED AND WILD POPULATIONS OF *Clarias gariepinus* (Osteichthys: Clariidae) USING RANDOM AMPLIFIED POLYMORPHIC DNA (RAPD) MARKER

Popoola, O. M., Fasakin, E. A., Awopetu, J. I.

ORIGINAL SCIENTIFIC PAPER

[show article ...](#)

AGE AND GROWTH OF THE SILVER CARP (*Hypophthalmichthys molitrix* Val.) IN TUDAKUL RESERVOIR, UZBEKISTAN

Kamilov, B. G.

ORIGINAL SCIENTIFIC PAPER

[show article ...](#)

PARASITOFUNA OF FIVE FRESHWATER FISHES IN A NIGERIAN FRESHWATER ECOSYSTEM

Chikwendu Ejere, V., Aguzie, O. I., Ivoke, N., Ekeh, F. N., Ezenwaji, N. E., Onoja, U. S., Eyo, J. E.

ORIGINAL SCIENTIFIC PAPER

[show article ...](#)

OCCURRENCE AND INTENSITY OF PARASITES IN EUROPEAN CATFISH, *Silurus glanis* L., 1758 FROM THE ANZALI WETLAND, SOUTHWEST OF THE CASPIAN SEA, IRAN

Daghigh Roohi, J., Sattari, M., Asgharnia, M., Rofchaei, R.

SHORT COMMUNICATION

[show article ...](#)

LENGTH WEIGHT RELATIONSHIPS OF THE EUROPEAN EEL *Anguilla anguilla* (Linnaeus, 1758) FROM SIX KARST CATCHMENTS OF THE ADRIATIC BASIN, CROATIA

Piria, M., Šprem, N., Tomljanović, T., Slišković, M., Jelić Mrčelić, G., Treer, T.

SHORT COMMUNICATION

[show article ...](#)

LIPID PEROXIDATION IN SMOKE-DRIED AFRICAN CATFISH TREATED WITH *Moringa oleifera* MARINADE, SALT OR BUTYLATED HYDROXYL ANISOLE

Dauda Adeyemi, K., Ahmed El-Imam, A. M., Oluwatunbo Olorunsanya, A., Solajo, F. E., Olatunji, O. S., Okukpe, K. M., Dosunmu, O. O., Shittu, R. M., Ramota Karim, O., Adewale Fatai, I.

SHORT COMMUNICATION

[show article ...](#)
<https://ribarstvo.agr.hr/volumes.php?lang=en&search=Vol%3A72&Num%3A1&field=title>

2021

Issue 2
Issue 1

2012 Volume 70

Issue Supplement 1
Issue 4
Issue 3
Issue 2
Issue 1

2011 Volume 69

Issue 4
Issue 3
Issue 2
Issue 1

2010 Volume 68

Issue 4
Issue 3
Issue 2
Issue 1

2009 Volume 67

Issue 4
Issue 3
Issue 2
Issue 1

2008 Volume 66

Issue 4
Issue 3
Issue 2
Issue 1

2007 Volume 65

Issue 4
Issue 3
Issue 2
Issue 1

2006 Volume 64

Issue 4
Issue 3
Issue 2
Issue 1

2005 Volume 63

Issue 4
Issue 3
Issue 2
Issue 1

2004 Volume 62

Issue 4
Issue 3
Issue 2
Issue 1

2003 Volume 61

Issue 4
Issue 3
Issue 2
Issue 1

2002 Volume 60

Issue 4

Croatian Journal of Fisheries Volume: 72 Issue: 1

**BLACKSPOTTED CROAKER, *Protonibea diacanthus* (Lacepède, 1802): A
NEW DIMENSION TO THE FISHING PATTERN IN WEST BENGAL, INDIA**
Dutta, S., Giri, S., Dutta, J., Hazra, S.

SUPPLEMENT TO FISHING PROFESSION

show article ...

**MANAGING IN FISHERIES BY USING INDIVIDUAL TRANSFERABLE
QUOTAS**

Ukić, H., Zubanović, V., Bečić, M., Slišković, M., Jelić Mrčelić, G.

Issue 3
Issue 2
Issue 1

2001 **Volume 59**

Issue 4
Issue 3
Issue 2
Issue 1

2000 **Volume 58**

Issue 4
Issue 3
Issue 2
Issue 1

1999 **Volume 57**

Issue 4
Issue 3
Issue 2
Issue 1

1998 **Volume 56**

Issue 4
Issue 3
Issue 2
Issue 1

1997 **Volume 55**

Issue 4
Issue 3
Issue 2
Issue 1

1996 **Volume 54**

Issue 4
Issue 3
Issue 2
Issue 1

1995 **Volume 53**

Issue 4
Issue 3
Issue 2
Issue 1

1994 **Volume 52**

Issue 4
Issue 3
Issue 2
Issue 1

[privacy policy](#)

Copyright Croatian Journal of Fisheries® 2021

LIPID PEROXIDATION IN SMOKE-DRIED AFRICAN CATFISH TREATED WITH *Moringa oleifera* MARINADE, SALT OR BUTYLATED HYDROXYL ANISOLE

Kazeem Dauda Adeyemi^{1*}, Aminat Mustapha Ahmed El-Imam², Ayotunde Oluwatunbo Olorunsanya³, Foluke Eunice Sola-Ojo¹, Olaife Salome Olatunji¹, Kehinde Mathias Okukpe¹, Olusegun Oyeshina Dosunmu³, Rafiat Morolayo Shittu¹, Olayinka Ramota Karim⁴, Ismail Adewale Fatai¹

¹ Department of Animal Production, University of Ilorin, Ilorin, Nigeria

² Department of Microbiology, University of Ilorin, Ilorin, Nigeria

³ Department of Animal Science, Ibrahim Badamosi Babangida University, Lapai, Nigeria

⁴ Department of Home-Economics and Food Science, University of Ilorin, Ilorin, Nigeria

⁵ Godbet Homestead Fish Farm, Ilorin, Nigeria

* Corresponding Author, E-mail: kazyadeyemi@gmail.com

ARTICLE INFO

Received: 25 August 2013

Received in revised form:

16 December 2013

Accepted: 3 December 2013

Available online: 17 December 2013

ABSTRACT

Smoke-dried fish is vulnerable to lipid peroxidation, which can reduce product quality and pose health risks to consumers. The study examined the antioxidant potency of *Moringa oleifera* marinade on oxidative stability of smoke-dried catfish in comparison with salt and Butylated hydroxyl anisole (BHA), a synthetic antioxidant. Seventy-two catfish (208±6 g) were processed, randomly assigned to six antioxidant treatment groups and hot smoked. The treatments are the control (0%), 1%, 2% and 3% (w/v) *Moringa oleifera* marinade (MOM), 5% Brine (w/v) and 0.2% BHA (w/v). The smoke-dried fish were stored at room temperature (35±1°C) for 8 weeks. Lipid peroxidation was monitored weekly using Thiobarbituric acid (TBA) assay. The results showed that *Moringa oleifera* marinade and BHA decreased lipid peroxidation more than ($p < 0.05$) the control (0.94 mg/MDA/kg) and salt (0.92 mg/MDA/kg) treated samples. This was shown by the lower Thiobarbituric acid reactive substance (TBARS) values of 1% MOM (0.44 mg/MDA/kg), 2% MOM (0.88 mg/MDA/kg), 3% MOM (0.85 mg/MDA/kg) and BHA (0.80 mg/MDA/kg) treated samples. A general increase in oxidative spoilage was observed for all treatments as storage time progressed. However, the increment was more ($p < 0.05$) intense in control and salt treated samples. No significant ($p > 0.05$) difference was observed among all *Moringa* treated samples and BHA. *Moringa oleifera* marinade could be used as an alternative to BHA in suppressing lipid peroxidation in smoke-dried African catfish stored for 8 weeks.

Keywords:

Antioxidant

BHA

Salt

Moringa

Lipid peroxidation

INTRODUCTION

Fish is an important source of food in many regions of the world. It is a good source of animal protein, which is indispensable to a balanced diet, and it is valued in many cultural culinary traditions. Fish is rich in beneficial unsaturated fats, particularly omega-3 fatty acids, that help protect human against diabetes, atherosclerosis and other cardiovascular diseases (Root et al., 2013). However, unsaturated fats are readily susceptible to lipid peroxidation. Peroxidation of lipids can instigate loss of nutritional and quality attributes of foods (Adeyemi et al., 2012) and could predispose consum-

ers to various diseases such as cancer and atherosclerosis (Martin et al., 2013). Due to the susceptibility of fresh fish to spoilage, various processing techniques are employed to curb deterioration, add value and maintain product quality. Some of the processing techniques are smoking, chilling, freezing, salting, canning and drying (Kumolu-Johnson et al., 2010). Among the aforementioned processing methods, smoking is the most common in many developing countries (Bako, 2005). Smoking enhances fish flavor, increases utilization, reduces waste, prolongs the shelf life of fish and increases protein availability (Jallow, 1995; Olley et al., 2000). Nonetheless, despite the effectiveness and the numerous

advantages derived from smoking, smoke-dried fish is still liable to oxidative damage resulting in partial or total loss of nutrients, oxidative rancidity, off-color and warmed-over flavor (Yasemen, 2007). Synthetic antioxidants such as butylated hydroxyl anisole (BHA), butylated hydroxyl toluene (BHT) are effective in curbing lipid peroxidation but are expensive and scarce, especially in developing countries (Adeyemi and Olorunsanya, 2012), and are carcinogenic (Nabavi et al., 2010; Dolatabadi and Kashanian, 2010). There are concerns about the toxicity and safety of synthetic antioxidants in relation to their accumulation and metabolism in the body tissues and organs (Nabavi et al., 2010; Barku et al., 2013). The need to find a lasting solution to the aforementioned problem necessitates the use of natural antioxidants. Phenolic extracts from legumes (Adeyemi et al., 2011), cereals, spices and herbs have been reported to retard lipid peroxidation in oils and fatty foods (Velioglu et al., 1998). *Moringa oleifera* is a natural antioxidant being researched into recently. *M. oleifera* contains myriad phytochemicals that are antioxidants. Kaempferol and Quercetin are recognized as the most effective antioxidants in *Moringa* leaves (Fahey, 2005). Siddhuraju and Becker (2003) reported that antioxidant properties of Kaempferol and Quercetin are higher than that of conventional antioxidants like Vitamin C, which is also abundant in *Moringa* leaves. *M. oleifera* leaf is cheap and readily available in many homes in Nigeria. Given the antioxidant properties of *M. oleifera*, it was hypothesized that *M. oleifera* marinade could reduce oxidative spoilage in smoke-dried catfish. The objective of the study was to determine the antioxidant effect of *M. oleifera* marinade on oxidative stability of smoke-dried catfish in comparison with salt, a common preservative used in fish smoking, and butylated hydroxyl anisole, a synthetic antioxidant.

MATERIALS AND METHODS

Preparation of *Moringa oleifera* Marinade

Fresh *Moringa* leaves were obtained from Olayinka Village Igbaja Oke-ode, Kwara state, Nigeria. The leaves were air-dried for 4 days and ground into powder using a food blender (Starlite, Model No: SL-999 CHINA). *Moringa oleifera* marinade (MOM) was prepared by adding separately specific quantity (10 g, 20 g or 30 g) of *M. oleifera* leaf powder to 1000 ml of water. 50 g salt or 2 g BHA were also added to a separate 1000 ml of water. The percentage concentration of the marinade was estimated using the formulae:

$$\% \text{ Concentration} = \frac{\text{Weight of additive (g)}}{\text{Volume of water (ml)}} \times 100$$

Salt was used as a treatment in this trial because it is a common preservative for smoke-dried fish.

Preparation and smoking of fish

The processing and smoking of the fish were carried out at Godbet Homestead Fish Farm, Basin road Ilorin, Nigeria. The total and average weights of the fish were 15 kg and 208±6 g, respectively. The fish were gutted using a sharp knife by cutting laterally from the end of the gill cover through the belly portion to the anus. Thereafter, they were washed and rinsed. The total and average weights of the fish after gutting were 13.3 kg and 200±7 g, respectively. The fish were treated and smoked in a metallic smoking chamber.

Experimental design

The fish were randomly assigned to six antioxidant treatments. The treatments were the control (0%), 1%, 2% and 3% *M. oleifera* marinade (MOM), 5% Brine and 0.2% BHA. Each treatment was replicated thrice with 4 fish/replicate. The fish were soaked in the marinade for 2 h. Thereafter, the fish were set in the smoking chamber consisting of five-twin tiers and subjected to hot smoking for 18 h with charcoal as the heat source. The tiers were interchanged every 6 h to ensure uniform heat distribution and drying. After smoking, the fish were stored in air-free netted boxes to prevent contamination by flies and to enhance flow-through ventilation throughout the storage period on laboratory shelves at room temperature (35±1°C) for 8 weeks.

Lipid Peroxidation in Smoked Catfish

Lipid peroxidation in the smoke-dried catfish samples was evaluated by the 2-thiobarbituric acid (TBA) test. The Thiobarbituric acid reactive substance (TBARS) values were measured in triplicate 10 g samples using the distillation method of Tarladgrs et al. (1964) as described by Adeyemi and Olorunsanya (2012). 10 g of the smoked catfish sample were homogenized with 47.5 ml of distilled water in a specimen bottle using a glass pestle. The homogenized mixture was rinsed with 50 ml of distilled water into a round bottom flask and, thereafter, 2.5 ml of dilute Hydrochloric acid (1:2 solutions) were added and the mixture was distilled through a condensing assembly to collect about 20 ml of the distillate. 5 ml of the distillate were mixed with 5 ml of the Thiobarbituric (TBA) acid (0.02 M) and boiled for 35 min in water. Afterwards, the mixture was cooled for 10 min in a water bath for color development. The absorbance readings were measured at a wavelength of 538 nm against a blank of 5 ml of Thiobarbituric acid reagent and 5 ml of Hydrochloric acid using a spectrophotometer (CECILL-2000). The absorbance values were multiplied by factor 7.8 (Tarladgrs et al., 1964) to obtain the TBARS value in milligram per malondialdehyde per kilogram sample (mg/MDA/kg).

Table 1. Main effect of antioxidant and storage time on oxidative stability of smoke-dried catfish

Factor	TBARS (mg/MDA/Kg)						
	control	1% MOM	2% MOM	3% MOM	5% Brine	0.2% BHA	
Antioxidant	0.94 ^b	0.84 ^a	0.88 ^a	0.85 ^a	0.92 ^b	0.80 ^a	
Storage weeks	1	2	3	4	5	6	7
	0.74 ^a	0.82 ^a	0.82 ^a	0.90 ^b	0.90 ^b	0.92 ^b	0.93 ^b
							1.25 ^c

a, b, c means having different superscript along the same row are significantly different ($p < 0.05$);
MOM = *Moringa oleifera* marinade

Statistical Analysis

The experiment followed a completely randomized design (CRD) in a 6 (antioxidant treatments) $\times$ 8 (storage weeks) factorial arrangement. The data obtained were analyzed by the Generalized Linear Model procedures of SAS Version 9.2 software (Statistical Analysis System, SAS Institute Inc, Cary, NC, USA). Mean differences were considered significant at $p < 0.05$ and separated by Duncan multiple range test.

RESULTS AND DISCUSSION

There were significant differences ($p < 0.05$) among the antioxidant treatments (Table 1). The control samples had the highest TBARS value that was not significantly different ($p > 0.05$) from salt treated samples. There was no significant difference among the Moringa and BHA treated samples. The potency of all Moringa treated samples could have resulted from various phytochemicals present in Moringa leaves. Fahey (2005) reported that kaempferol and quercetin were the most effective antioxidants in Moringa leaves in addition to other flavonoid antioxidants such as isoquercitrin, kaempferitrin and rhamnetin. Moringa leaves also contain high amount of vitamin C and E, which are potent antioxidants (Grubben and Denton, 2004). However, this observation was contrary to the report given by Adeyemi et al. (2012) in which *M. oleifera* powder was not effective in cooked broiler meat samples but was effective in raw broiler meat samples. This suggests that antioxidant compounds in *M. oleifera* leaves are heat labile. The discrepancy between the earlier report and the present study could also be due to the nature of heat used. Furthermore, the oxidative stability exhibited by Moringa treated samples in the present study could be due to the synergistic effect of the antioxidant compounds liberated during the smoking process from the charcoal and the inherent antioxidants present in *M. oleifera* leaves. Smoking deposits certain preservative compounds into the smoked products (Ekpenyong and Ibok, 2012). The amount and type of compounds deposited depend on the type of wood or charcoal used (Ekpenyong and Ibok, 2012).

A general increase in TBARS values was observed as stor-

age progressed (Table 1). There were no significant differences among the TBARS values recorded in the first, second and third week of storage. These values were significantly different from TBARS values observed from the fourth to the seventh week. The eighth week incurred the highest TBARS value that was significantly different ($p < 0.05$) from those observed for other weeks. This observation was in line with the report given by Yasemen (2007) who observed that peroxide and thiobarbituric acid values of hot smoke-dried catfish stored in refrigerator increased, whilst the sensory scores declined as storage time continued. The author concluded that in order to improve the shelf life of hot smoked catfish, a suitable preservation and packaging technique should be used or developed.

There was no significant difference among the antioxidant treatments in the first, third, fourth and sixth week of storage (Table 2). In the second week, brine (salt) treated samples had the highest TBARS value that was significantly different ($p < 0.05$) from other treatments. In the seventh week, there was no significant difference ($p > 0.05$) between the control and BHA treated samples. This was unexpected because BHA has been reported to be a potent synthetic antioxidant (Mariod et al., 2012; Laghari et al., 2011). There was no significant difference among all levels of Moringa and salt treated samples in the seventh week. In the eighth week, BHA and all levels of Moringa marinade performed equally, while the control and salt treated samples were not significantly different ($p > 0.05$). There was a steady increase in TBARS values of the control samples from week one to eight. The pattern of increase in TBARS values does not differ for 1%, 2%, 3% MOM and BHA. This clearly attests the preservative capability of *M. oleifera* marinade (MOM) which lends credence to our earlier study (Adeyemi et al., 2013) in which MOM maintained the chemical and microbiological quality of smoke-dried catfish. The antioxidant effect of *M. oleifera* marinade could be attributed to the presence of phenolic compounds, which donate hydrogen atoms from their hydroxyl groups thus reducing the formation of hydroperoxides, the first product of lipid peroxidation (Velasco and Williams, 2011). Phenolic compounds exert their antioxidant properties by three main processes (Velasco and Williams, 2011). These include one

Table 2. Combined effect of antioxidant and storage time on oxidative stability of smoke-dried African catfish

Antioxidant	TBARS (mg/MDA/kg)							
	Storage week							
	1	2	3	4	5	6	7	8
Control	0.53 ^a	0.74 ^x	0.78 ^b	0.85 ^b	0.99 ^y	0.99 ^c	1.20 ^y	1.27 ^d
1% MOM	0.58 ^a	0.64 ^x	0.75 ^a	0.82 ^b	0.76 ^x	0.90 ^b	0.89 ^x	0.90 ^b
2% MOM	0.52 ^a	0.70 ^x	0.66 ^a	0.81 ^b	0.90 ^x	0.95 ^b	0.80 ^x	0.98 ^b
3% MOM	0.54 ^a	0.63 ^x	0.76 ^a	0.82 ^b	0.86 ^x	0.81 ^b	0.81 ^x	0.91 ^b
5% Salt	0.67 ^a	0.87 ^y	0.76 ^a	0.78 ^a	0.85 ^x	0.82 ^a	0.90 ^x	1.145 ^y
0.2% BHA	0.56 ^a	0.69 ^x	0.74 ^a	0.82 ^b	0.85 ^x	0.89 ^b	1.02 ^x	1.03 ^x

a, b, c, d means having different superscript along the same row are significantly different ($p < 0.05$); x, y means having different subscript along the same column are significantly different ($p < 0.05$)

or more transition metal-chelating activities (Andjelković et al., 2006), singlet-oxygen quenching capacity (Mukai et al., 2005) and free radical scavenging activity (Zheng et al., 2009). *M. oleifera* leaves contain ample amount of vitamin C (ascorbic acid) and vitamin E (α -tocopherol) (Grubben and Denton, 2004) which exhibited high radical scavenging properties as reported by Kulisic et al. (2004). In addition, Yeum et al. (2009) reported synergistic properties between α -tocopherol and ascorbic acid against peroxidation.

CONCLUSION

M. oleifera marinade and BHA were more effective than control and salt in protecting smoke-dried catfish against lipid peroxidation. There was an increment in oxidative spoilage as storage time progressed. However, the increment was more intense in the control and salt treated samples. *M. oleifera* marinade and BHA were equally potent as antioxidants. *M. oleifera* marinade could be used to retard lipid peroxidation in smoke-dried catfish for two months. Since salt is a common preservative used in fish smoke drying, further study should be carried out to determine the combined effect of *M. oleifera* marinade and salt on lipid peroxidation in smoke-dried catfish.

Sažetak

LIPIDNA PEROKSIDACIJA KOD DIMLJENOG AFRIČKOG SOMA TRETIRANOG MARINADOM *Moringa oleifera*, SOLJU I BUTIL HIDROKSI ANISOLOM

Sušena dimljena riba izložena je lipidnoj peroksidaciji, što može smanjiti kvalitetu krajnjeg proizvoda, ali i ugroziti zdravlje potrošača. Ovim istraživanjem ispitano je potencijalno antioksidantsko djelovanje mari-

nade *Moringa oleifera* na stabilnost oksidacije kod dimljenog soma te je ono uspoređeno s djelovanjem soli i butil hidroksi anisola (BHA), sintetičkog antioksidanta. Obrađeno je 72 primjeraka soma (208 ± 6 g), a ravnomjerno su raspoređeni u 6 skupina tretiranih antioksidansima te izloženih vrućem dimu. Skupine s obzirom na vrstu tretmana i postotak izloženosti bile su sljedeće: kontrolna skupina (0%), 3 skupine somova tretiranih marinadom *Moringa oleifera* (MOM) (1%, 2% i 3%), somovi u rasolu (w/v) (5%) i somovi u butil hidroksi anisolu (w/v) (0,2%). Dimljena riba bila je pohranjena na sobnoj temperaturi ($35 \pm 1^\circ\text{C}$) 8 tjedana. Peroksidacija lipida tjedno je praćena pomoću ispitivanja tiobarbiturnom kiselinom (TBA). Rezultati su pokazali smanjenje lipidne peroksidacije kod skupina somova u marinadi *Moringa oleifera* i u butil hidroksi anisolu ($p < 0,05$), za razliku od rezultata povećane lipidne peroksidacije kod somova iz kontrolne skupine (0,94 mg/MDA/kg) i onih tretiranih solju (0,92 mg/MDA/kg). Dakle, reaktivne promjene prilikom ispitivanja tiobarbiturnom kiselinom (TBARS) su kod pojedinih skupina somova bile manje: uzorci tretirani 1% marinadom *Moringa oleifera* (0,44 mg/MDA/kg), uzorci tretirani 2% i 3% istom marinadom (0,88 mg/MDA/kg; 0,85 mg/MDA/kg) i uzorci tretirani BHA-om (0,80 mg/MDA/kg). Tijekom izloženosti riba prije navedenim uvjetima, praćen je proces kvarenja riba uzrokovan oksidacijom. Prirast je bio intenzivniji u kontrolnoj i skupini riba tretiranih solju ($p < 0,05$). Nema značajne razlike kod skupina riba tretiranih marinadom *Moringa oleifera* i onih tretiranih butil hidroksi anisolom. Marinada *Moringa oleifera* mogla bi se u osmotjednom periodu izloženosti dimljenog afričkog soma koristiti kao alternativa za BHA u suzbijanju lipidne peroksidacije.

Ključne riječi: antioksidant, BHA, sol, *Moringa*, lipidna peroksidacija

REFERENCES

- Adeyemi, K. D., Ahmed El-Imam, A. M., Olorunsanya, A. O., Sola-Ojo, F. E., Okukpe, K. M., Dosunmu, O. O., Shittu, R. M., Idris, J. T. (2013a): Effect of *Moringa oleifera* marinade on proximate composition and sensory characteristics of smoke dried African catfish (*Clarias gariepinus*). *Croatian Journal of Fisheries*, 71, 11–18.
- Adeyemi, K. D., Ahmed El-Imam, A. M., Dosunmu, O. O., Lawal, O. K. (2013b): Effect of *Moringa oleifera* marinade on microbial stability of smoke-dried African catfish (*Clarias gariepinus*). *Ethiopian Journal of Environmental Studies and Management*, 6, 1, 104–109.
- Adeyemi, K. D., Olorunsanya, A. O., Akanji, M. A. (2011): Effect of watermelon (*Citrus lantus*) extracts on oxidative stability of broiler meat. *African Journal of General Agriculture*, 7, 3, 103–108.
- Adeyemi, K. D. and Olorunsanya, O. A. (2012): Comparative analysis of phenolic composition and antioxidant effect of four Cowpea (*Vigna unguiculata*) varieties on broiler meat. *Iranian Journal of Applied Animal Science*, 2, 4, 343–349.
- Adeyemi, K. D., Olorunsanya, A. O., Sola-Ojo, F. E., Okukpe, K. M., Malomo, G. A., Amira, O. J., Badmus, A. H. A., Ahmed El-Imam, A. M., Shittu, R. M., Aboyeji, O. W. (2012): Effect of *Moringa oleifera* powder on oxidative stability of cooked and raw broiler meat. *Journal of Applied Agricultural Research*, 4, 2, 175–181.
- Andjelković, M., Van Camp, J., De Meulenaer, B., Depaemelaere, G., Socaciu, C., Verloo, M., Verhe, R. (2006): Iron-chelation properties of phenolic acids bearing catechol and galloyl groups. *Food Chemistry*, 98, 23–31.
- Bako, W. S. (2005): The role of women in fish processing: Handling and Marketing in Kainji lake basin. *Proceedings of the 19th Annual Conference of the Fisheries society of Nigeria (FISON)*, Nov 29–Dec 3, 2004, Ilorin, Nigeria.
- Barku, V., Opoku-Boahen, Y., Owusu-Ansah, E., Mensah, E. (2013): Antioxidant activity and the estimation of total phenolic and flavonoid contents of the root extract of *Amaranthus spinosus*. *Asian Journal of Plant Science and Research*, 3, 1, 69–74.
- Dolatabadi, J. E. N. and Kashanian, S. (2010): A review on DNA interaction with synthetic phenolic food additives. *Food Research International* 43, 5, 1223–1230.
- Ekpenyong, C. O. and Ibok, C. O. (2012): Effect of smoking, salting and frozen storage on the nutrient composition of the African catfish (*Clarias gariepinus*). *Journal of Food Agriculture and Environment*, 10, 1, 12–16.
- Fahey, J. W. (2005): *Moringa oleifera*: A review of the medical evidence for its nutritional, therapeutic and prophylactic properties. Part 1. *Trees Life Journal*.
- Grubben, G. J. H. and Denton, O. A. (2004): *Plant Resources of Tropical Africa 2. Vegetables*. PROTA Foundation, Wageningen, Netherlands/ Backhuys Publishers, Leiden, Netherlands/ CTA, Wageningen, Netherlands.
- Jallow, A. M. (1995): Contribution of improved chorkor oven to artisanal fish smoking in the Gambia. *Proceedings of the workshop on seeking improvements in Fish Technology in West Africa*. Nov, 7–9, 1994, Congo, pp. 22–28.
- Kulisić, T., Radonić, A., Katalinić, V., Milos, M. (2004): Use of different methods for testing antioxidative activity of oregano essential oil. *Food Chemistry*, 85, 4, 633–640.
- Kumolu-Johnson, N. F., Aladetohun, C. A., Ndimele, A. (2010): The effect of smoking on the nutrient composition of the African catfish (*Clarias gariepinus*). *African Journal of Biotechnology*, 9, 73–76.
- Laghari, A. H., Memon, S., Nelofar, A., Khan, K. M., Yasmin, A. (2011): Determination of free phenolic acids and antioxidant activity of methanolic extracts obtained from fruits and leaves of chenopodium album. *Food Chemistry*, 126, 4, 1850–1855.
- Mariod, A., Ibrahim, R., Ismail, M., Ismail, N. (2012): Antioxidant activity of phenolic extracts from kenaf (*Hibiscus cannabinus*) seed-cake. *Grasa y Aceites*, 63, 2, 167–174.
- Martin, C., Zhang, Y., Tonelli, C., Petroni, K. (2013): Plants, diet, and health. *Ann. Rev. Plant Biology*, 64, 19–46.
- Mukai, K., Nagai, S., Ohara, K. (2005): Kinetic study of the quenching reaction of singlet oxygen by tea catechins in ethanol solution. *Free Radical Biology and Medicine*, 39, 752–761.
- Nabavi, F. S., Ebrahimzadeh, M. A., Nabavi, M. S., Eslami, B. (2010): Antioxidant activity of flower, stem and leaf extracts of ferula Gummosa boiss. *Grasas Aceites*, 61, 3, 244–250.
- Olley, J., Doe, P. E., Heruwat, E. S. (2000): The influence of drying and smoking on the nutritional properties of Fish. An introductory overview. In: *International Union of Nutritional Science Conference*, Malaysia. Pp 1–6.
- Root, M., Collier, S. R., Zwetsloot, K. A., West, K. L., McGinn, M. C. (2013): A randomized trial of fish oil omega-3 fatty acids on arterial health, inflammation, and metabolic syndrome in a young healthy population. *Nutrition Journal*, 12, 40–47.
- Siddhuraju, P. and Becker, K. (2003): Antioxidant properties of various solvent extracts of total phenolic constituents from three different agro-climatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *Journal of Agricultural and Food chemistry*, 51, 2144–2155.
- Tarlalgrs, B. G., Pearson, A. M., Dungan, L. R. (1964): Chemistry of 2-TBA test for determination of oxidative Rancidity in foods and formation of TBA malondialdehyde complex without acid heat treatment. *Journal of Science of Food and Agric*, 15, 607–620.
- Velasco, V. and Williams, P. (2011): Improving meat quality through natural antioxidants. *Chilean Journal of Agricultural Research*, 71, 2, 313–322.
- Velioglu, Y. S., Mazza, G., Gao, L., Oomah, B. D. (1998): Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *Journal of Agricultural and Food chemistry*, 46, 4113–4117.
- Vasemen, Y. (2007): Quality changes of hot smoked catfish (*Clarias gariepinus*) during refrigerated storage. *Journal of Muscle Foods*, 18, 4, 391–400.
- Yeum, K., Beretta, G., Krinsky, N. I., Russell, R. M., Aldini, G. (2009): Synergistic interactions of antioxidant nutrients in a biological model system. *Nutrition*, 25, 7, 839–846.
- Zheng, G., Xu, L., Wu, P., Xie, H., Jiang, J., Chen, F., Wei, X. (2009): Polyphenols from Longan seeds and their radical-scavenging activity. *Food Chemistry*, 116, 433–436.