

Impact of Malnutrition on Serum Levels of Pro-inflammatory Cytokines among *Plasmodium falciparum* Malaria Subjects in Nigeria

Olusola Ajibaye^{1,2}, Victoria Olawunmi Adaramoye³, Shittu Olalere⁴, Agatha N. Ibekpobaoku⁵, Mayowa A. Olasope^{2,6}, Gabriel O. Eniafe¹, Judith C. Okanazu¹, Bassey A. Orok¹, Sunday Ojo⁷, Yetunde Adeola Olukosi¹, Emmanuel Oluwadare Balogun^{8,9}

¹Malaria Genomics Research and Training Centre, Biochemistry and Nutrition Division, Nigerian Institute of Medical Research, Yaba, ²Department of Medical Biochemistry, Eko University of Medicine and Health Sciences, Ojo, ³Department of Zoology and Environmental Biology, Lagos State University, Lagos, ⁴Department of Obstetrics and Gynaecology, Lagos University Teaching Hospital, Surulere, ⁵Department of Zoology, Parasitology and Bioinformatics Unit, University of Lagos, Akoka, Lagos, ⁶Department of Zoology, Parasitology Unit, University of Ilorin, Ilorin, ⁷Department of Zoology, Kwara State University, Molete, Ilorin, Kwara State, ⁸Department of Biochemistry, Ahmadu Bello University, Zaria, Nigeria, ⁹Department of Biomedical Chemistry, Graduate School of Medicine, The University of Tokyo, Bunkyo-ku, Tokyo, Japan

Abstract

Background: Understanding the impact of malnutrition on innate immune response in *Plasmodium falciparum* (Pf)-infected subjects is critical for malaria control. **Aims and Objectives:** This study aims to investigate the nutritional status and innate immune response of Pf-infected subjects in Lagos, Nigeria. **Materials and Methods:** A total of 1183 patients with a history of fever or axillary temperature $\geq 37^\circ\text{C}$ were screened microscopically for Pf at Ijeda General Hospital, Lagos, Nigeria. Malnutrition was determined according to the U.S National Center for Health Statistics (NCHS) as stunting, wasting, or underweight when the Z-score is < -2 in the participants aged < 20 years. Serum levels of tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β), and IL-12 were determined by capture ELISA while hematological parameters were measured using an automated hematology system. **Results:** A total of 384 volunteers were positive for Pf, of which 114 were < 20 years with a median age of 10 years. Overall malaria prevalence was 20.89%. The malnutrition rate was 89.5%; 24 (21.05%) were stunted, 30 (26.32%) were underweight, and 48 (42.11%) were wasted. Pro-inflammatory cytokine responses were not affected by the type of malaria. TNF- α was higher in participants < 5 years ($P = 0.001$) and in malnourished patients ($P < 0.05$). **Conclusion:** Together, it could be deduced that nutritional status influences *Plasmodium falciparum* malaria outcomes and progression pattern.

Keywords: Cytokines, IL-12, TNF- α malnutrition, necrosis, plasmodium

Résumé

Contexte: Comprendre l'impact de la malnutrition sur la réponse immunitaire innée chez les sujets infectés par *Plasmodium falciparum* (Pf) est essentiel pour la lutte contre le paludisme. **Buts et objectifs:** Cette étude vise à étudier l'état nutritionnel et la réponse immunitaire innée des sujets infectés par Pf à Lagos, au Nigeria. **Matériels et méthodes:** Un total de 1183 patients ayant des antécédents de fièvre ou une température axillaire $\geq 37^\circ\text{C}$ ont fait l'objet d'un dépistage microscopique de Pf à l'hôpital général Ijeda, Lagos, Nigeria. La malnutrition a été déterminée selon le National Center for Health Statistics (NCHS) des États-Unis comme un retard de croissance, une émaciation ou une insuffisance pondérale lorsque le score Z est < -2 chez les participants âgés de moins de 20 ans. Les taux sériques de facteur de nécrose tumorale alpha (TNF- α), d'interleukine-1 β (IL-1 β) et d'IL-12 ont été déterminés par capture ELISA, tandis que les paramètres hématologiques ont été mesurés à l'aide d'un système d'hématologie automatisé. **Résultats:** Au total, 384 volontaires étaient

Address for correspondence: Dr. Olusola Ajibaye, Malaria Genomics Research and Training Centre, Nigerian Institute of Medical Research, 6 Edmond Crescent, Off Murtala Mohammed Way, P.M.B 2013, Yaba, Lagos, Nigeria. E-mail: sajibaye@yahoo.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

Access this article online

Quick Response Code:



Website:
www.annalsafmed.org

DOI:
10.4103/aam.aam_209_23

How to cite this article: Ajibaye O, Adaramoye VO, Olalere S, Ibekpobaoku AN, Olasope MA, Eniafe GO, *et al.* Impact of malnutrition on serum levels of pro-inflammatory cytokines among *plasmodium falciparum* malaria subjects in Nigeria. Ann Afr Med 2024;23:379-84.

Submitted: 18-Dec-2023 **Revised:** 24-Jan-2024

Accepted: 02-Feb-2024 **Published:** 08-Jun-2024

positifs pour le Pf, dont 114 étaient âgés de moins de 20 ans avec un âge médian de 10 ans. La prévalence globale du paludisme était de 20,89 %. Le taux de malnutrition était de 89,5 % ; 24 (21,05 %) souffraient d'un retard de croissance, 30 (26,32 %) d'une insuffisance pondérale et 48 (42,11 %) d'émaciation. Les réponses des cytokines pro-inflammatoires n'ont pas été affectées par le type de paludisme. Le TNF- α était plus élevé chez les participants de moins de 5 ans ($P = 0,001$) et chez les patients souffrant de malnutrition ($P < 0,05$). **Conclusion:** On peut en déduire que l'état nutritionnel peut influencer les résultats et le schéma de progression du paludisme.

Mots-clés: Cytokines, IL-12, TNF- α , malnutrition, necrosis, plasmodium

INTRODUCTION

Malaria is still a leading global health concern. With 3.4 billion people at risk and 198 million clinical cases, sub-Saharan African countries have the highest burden.^[1] *Plasmodium* species associated with human infections are *Plasmodium falciparum* (Pf), *Plasmodium malariae*, *Plasmodium vivax*, *Plasmodium ovale*, and *Plasmodium knowlesi*.^[2] Pf is the most virulent species of *Plasmodium* and is responsible for the adverse consequences of malaria.

Pf malaria and malnutrition are major causes of child morbidity and mortality with common consequences ranging from cognitive impairment to decreased school performance,^[3-6] yet their precise interaction remains underreported. Moreover, undernutrition is highly prevalent in developing countries and is considered to be the underlying cause of more than 50% of all childhood deaths in the world.^[7,8] In sub-Saharan Africa, 38% of under-five children suffer from chronic malnutrition or stunting (height-for-age [HAZ] z-score below -2 of the international growth reference), and acute malnutrition or wasting (weight-for-height [WHZ] z-score below -2) affects 9% of preschool children.^[7]

Malnutrition is estimated to cause about half of the world's 12 million annual deaths in children <5 years of age as well as substantial proportions of infectious disease morbidity.^[9,10] Although many studies have shown that Pf infection can result in acute weight loss, improvements in growth and other anthropometric indices have been described in children protected from malaria.^[11-13] Nutritional status hence is considered to be one of the major determinants of host resistance to infection.^[14] In sub-Saharan Africa, the impact of chronic malnutrition or stunting (HAZ z-score below -2 of an international growth reference) and acute malnutrition or wasting (WHZ z-score below -2) is debilitating, especially in children.^[14] In Nigeria, stunting or chronic malnutrition has been reported in about 10 million children resulting in increased susceptibility to infections, more commonly malaria.^[15] This is because malaria is one of the most common parasitic infections occurring in comorbidity.^[16]

Malnutrition is the primary cause of immunodeficiency worldwide; however, there have been conflicting reports on the influence of malnutrition on the immune response of malaria patients. Malnutrition could influence the development of the protective immune response acquired during infection and regulate antimalarial immunity. Cytokines play both protective and pathological roles in malaria; however, an early

and effective inflammatory response is crucial in the control of parasitemia and resolution of malaria infection through the mechanism of tumor necrosis factor-alpha (TNF- α).^[17] In response to infection, the immune system first executes innate and then subsequently acquired host defense functions of high diversity. Both processes involve the activation and propagation of immune cells and synthesis of an array of molecules requiring DNA replication, RNA expression, and protein synthesis and secretion and therefore consume additional anabolic energy.

Pro-inflammatory cytokines, which are mediators of inflammation further, increase the catabolic response acclaiming much necessity for protein-energy and nutritional status of infected hosts. Interleukin-1beta (IL-1 β), TNF- α , and IL-12 are pro-inflammatory cytokines crucial in malaria immunity.^[7] However, a cohort study reported the downregulation of malaria-specific immunity in contrast to the report from a study in Papua New Guinea, which indicated that malnutrition protected children from Pf^[18] while Müller *et al.*^[19] and Müller *et al.*^[20] reported no association.

There is a need for better understanding of the impact of malnutrition on innate immune response in Pf malaria-infected individuals though it is widely recognized that malnutrition compromises the immune function, resulting in a higher risk of infection. However, there is paucity of data on the relationship between malaria, malnutrition, and antimalarial immunity in Nigeria. It is therefore pertinent to investigate their inter-relationship.

METHODOLOGY

Study site and enrollment

In this cross-sectional study, participants were recruited at Ijede General Hospital (IJGH), Ikorodu Local Government Area of Lagos State, Nigeria [Figure 1]. Ijede is a low socio-economic settlement with ecological features favoring malaria. IJGH is one of the centers characterized in 2011 as sentinel sites for malaria research in Nigeria; on average, there are 400 outpatients per week. Patients were recruited at the peak of malaria transmission period (i.e. May–July) in 2021. Patients presenting at the IJGH with symptoms suggestive of malaria (axillary temperature $\geq 37.5^\circ\text{C}$ or history of fever in the last 48 h) and with Pf mono-infection who consented were enrolled in the study.

Parasitemia assessment

Finger-pricked blood samples were collected from each participant. Parasite densities were estimated in thick




Figure 1: Map showing the location of the study site (Ijode General Hospital), Ikorodu Local Government Area of Lagos State

blood films, assuming an average white blood cell count of 8000/ μ L. All slides were Giemsa stained and read by two certified microscopists, and ties were broken by a third expert microscopist. Venous blood samples of positive patients and 40 uninfected controls were collected into EDTA bottles for ELISA and hematological assays.

Anthropometric measurements

Anthropometric data were collected in accordance with internationally recommended procedures.^[4,5,9,21] Weight measurements were taken using a scale (Téfal, Paris, France) precise to the nearest 100 g. Recumbent length measurements were taken for children under 2 years of age, while standing height was measured for older participants using a stadiometer (SECA, Hamburg, Germany). Body mass index (BMI) (kg/m^2) was calculated and used as an index of nutritional status for the participants; those within the range of 18.5 – 24.9 were grouped as normal, while those less than or above this range were categorized malnourished or overweight, respectively.

Evaluation of cytokine response

Levels of TNF- α , IL-12, and IL-1 β were assessed in sera of included participants using antigen capture ELISA.

Statistical analyses

The nutritional indicators HAZ, weight-for-age (WAZ), and WHZ were computed in z-scores of the WHO/National Center for Health Statistics (NCHS) reference using Epi Info software V.6 for participants below 20 years after BMI computation. Stunting, underweight, and wasting were defined for values below – 2 for HAZ, WAZ, and WHZ, respectively. SPSS V.20

Table 1: Nutritional status and cytokine levels of participants

Parameters	Uncomplicated malaria		
	Infected, <i>n</i> (%)	Uninfected, <i>n</i> (%)	<i>P</i>
Age (years), mean $\pm$ SEM (years)	13 $\pm$ 12	15.5 $\pm$ 2.7	
<5	37 (32.8)	14 (35.00)	
>5	77 (67.2)	26 (65.00)	
Mean parasitemia/ μ L (years)			
<5	136,267 $\pm$ 1449	-	0.04
$\geq$ 5	32,912 $\pm$ 1256	-	
Gender			
Female	55 (48.44)	20 (50.00)	
Male	59 (51.56)	20 (50.00)	
Mean parasitemia/ μ L			
Female	37,203 $\pm$ 4751	-	0.43
Male	44,239 $\pm$ 2985	-	
Nutritional status (participants <20 years)			
Normal	12 (10.56)	14 (35.00)	
Stunted (HAZ)	24 (21.05)	2 (5.00)	
Wasted (WHZ)	48 (42.11)	14 (35.00)	
Underweight (WAZ)	30 (26.32)	10 (25.00)	
Cytokines (ng/L)			
TNF- α	362.5 $\pm$ 40.00	97.5 $\pm$ 10.80	0.00016
IL-12	83.44 $\pm$ 2.14	57.25 $\pm$ 2.80	0.003
IL-1 β	23.14 $\pm$ 2.30	10.47 $\pm$ 0.30	0.02

HAZ=Height-for-age, WHZ=Weight-for-height, WAZ=Weight-for-age, SEM=Standard error of mean, IL-12=Interleukin-12

IBM, New York, USA was used for analysis of variance, multivariate regression, and correlation analysis. Differences were considered significant at $P < 0.05$.

Ethics

Approval was obtained from the NIMR Institutional Ethics Review Board for clearance to carry out the research.

RESULTS

A total of 384 participants were included in this cross-sectional study meeting the inclusion criteria from the 1838 screened. Malaria prevalence was 20.89%. Of these, 252 (65.6%) were malnourished, 96 (25%) normal, and 36 (9.4%) overweight. In the control group (uninfected participants), 50% were malnourished [Table 1]. A total of 114 infected participants were <20 years. Using the WHO/NCHS reference, 48 (42.11%) of participants <20 years were wasted, 24 (21.05%) were stunted, and 30 (26.32%) of

them were underweight [Table 1]. There was no significant difference in infection intensity by parasitemia between the male and female participants ($P = 0.43$); however, malaria parasite density (MPD) was significantly higher ($P < 0.05$) children <5 years than in older subjects (<20 years) [Table 1].

Correlation of tumor necrosis factor-alpha, interleukin-12, and interleukin-1beta levels with intensity of *Plasmodium falciparum* infection

It has been shown that Th1 responses directed to wide ranges of *Pf* antigens were associated with *Pf* infection.^[22-24] In the present study, levels of IL-12 decreased with increasing intensity of infection (from ≤ 1000 to $\leq 600,000$ pRBCs/ μ L of blood) while TNF- α increased with increasing intensity of infection although not significantly ($P > 0.05$) [Table 2].

TNF- α levels were significantly higher in malnourished patients ($P < 0.0001$) whereas there was no significant difference in the levels of IL-12 and IL-1 β in all the nutritional status and the control ($P < 0.05$) [Table 3]. From Table 2, with increasing parasitemia, IL-1 β response reduced significantly but not so with IL-12. Moreover, all the pro-inflammatory cytokines measured were significantly higher in infected than uninfected participants irrespective of the nutritional status ($P < 0.05$) [Table 3].

Hematological parameters

The frequency of anemia was 9.5% in the malnourished group and 0% in the normal and control groups [Table 4]. Approximately 26% of the malnourished infected participants had platelet counts $<150 \times 10^9/L$ (altered platelet count) as

Table 2: Effect of malaria parasite density on cytokine response of participants

Cytokines (μ/L)	Malaria parasite density groups (parasites/ μ L), mean $\pm$ SD				
	≤ 1000	$\leq 100,000$	$\leq 600,000$	P	n
IL-12	90 $\pm$ 0.6	83.5 $\pm$ 3.1	82.7 $\pm$ 2.1	0.85	114
IL-1 β	50 $\pm$ 1.2	21.34 $\pm$ 2.4	23.81 $\pm$ 2.3	0.08	114
TNF- α	250 $\pm$ 21.1	365 $\pm$ 53.3	368.2 $\pm$ 67.5	0.88	114

SD=Standard deviation, TNF- α =Tumor necrosis factor-alpha, IL-1 β =Interleukin-1 beta, IL-12=Interleukin-12

Table 3: Cytokine levels with respect to nutritional status of participants

Nutritional status	Mean $\pm$ SEM (test/CTRL)		
	TNF- α (μ/L)	IL-1 β (μ/L)	IL-12 (μ/L)
Stunted			
Infected ^{a,c}	446.70 $\pm$ 52.59	28.88 $\pm$ 2.97	86.76 $\pm$ 2.97
Uninfected	111.00 $\pm$ 18.51	10.15 $\pm$ 0.43	50.50 $\pm$ 6.43
P	0.001*	0.12	0.06
Wasted			
Infected ^b	445.75 $\pm$ 52.12	26.14 $\pm$ 3.22	88.77 $\pm$ 15.25
Uninfected	100.05 $\pm$ 38.22	9.44 $\pm$ 0.88	51.02 $\pm$ 10.10
P	0.03*	0.04*	0.05
Normal			
Infected ^{a,b}	356.25 $\pm$ 79.3	26.85 $\pm$ 6.29	81.25 $\pm$ 2.95
Uninfected	91.25 $\pm$ 11.94	11.19 $\pm$ 0.42	63.75 $\pm$ 5.49
P	0.002*	0.3	0.82
Underweight			
Infected ^c	351.0 $\pm$ 115.47	23.33 $\pm$ 5.83	80 $\pm$ 5.77
Uninfected	188 $\pm$ 10.44	11.23 $\pm$ 1.33	38 $\pm$ 7.89
P	0.008*	0.05	0.05

t-test, ANOVA: *Significant at $P < 0.05$ (stunted vs. wasted and normal), ^{a,b,c}Tukey's. SEM=Standard error of mean, TNF- α =Tumor necrosis factor-alpha, IL-1 β =Interleukin-1 beta, IL-12=Interleukin-12, CTRL=Uninfected controls

Table 4: Anemia and altered platelet counts in study participants

Parameter frequency	Malnutrition (HAZ, WHZ, and WAZ) (%)	Normal, n (%)	Control, n (%)
Anemia (Hb <7 g/dL)	24 (9.5)	0	0
Fever (auxiliary temperature $\geq 37.5^\circ C$)	80 (31.7)	15 (15.6)	0
Altered platelet count $<150 \times 10^9/L$	65 (25.8)	40 (41.6)	20 (50)

HAZ=Height-for-age, WHZ=Weight-for-height, WAZ=Weight-for-age, Hb=Hemoglobin

Table 5: Cytokine levels with respect to malaria infection status (i.e., asymptomatic or symptomatic)

Cytokines	Malaria infection status (mean $\pm$ SEM)		P
	Asymptomatic	Symptomatic	
TNF- α	395.11 $\pm$ 17.31	417.74 $\pm$ 24.74	0.15
IL-1 β	67.88 $\pm$ 29.26	25.77 $\pm$ 1.69	0.08
IL-12	87.52 $\pm$ 1.57	88.39 $\pm$ 2.22	0.22

Values are mean levels of pro-inflammatory cytokines $\pm$ SEM (t-test: $P > 0.05$). SEM=Standard error of mean, TNF- α =Tumor necrosis factor-alpha, IL-1 β =Interleukin-1 beta, IL-12=Interleukin-12

compared to 5 (12.5%) uninfected controls having similar platelet counts. Only the malnourished infected participants 80 (31.7%) had fever (temperature $\geq 37.5^{\circ}\text{C}$) in the study.

Pearson's correlation showed that IL-12 was negatively correlated ($r = -0.327$, $P = 0.03$) with platelet counts in the test participants with a multiple regression coefficient of -2.48 and $P = 0.0018$ [Table 5].

DISCUSSION

This cross-sectional study conducted at the onset of the rainy season in 2021 set out to investigate whether nutritional status affects the cellular immune response of malaria patients in Lagos, Nigeria. The findings showed that the malaria burden is relatively high in rural areas of Nigeria similar to what was reported by Afolabi *et al.*^[25] This may not be unconnected with the noncompliance of people to the National Malaria Control Program guidelines for malaria control in Nigeria. The geometric mean parasite density was significantly higher in males than females due to higher immunity in females and body exposure by males consistent with Akanbi *et al.*^[26] and Mandel and Whit.^[27]

TNF- α was higher in <5 in response to slowly reducing MPD due to low immunity and increased TNF- α response at this age. Malnourished patients presented high levels of TNF- α , a potent protective immune component, of which (low) leptin-induced high levels have been reported in malnourished individuals^[28] and associated with cachexia seen in malnourished malaria subjects. However, no association was found between malnutrition and IL-1 β although IL-12 levels have been reported to substantiate levels of TNF- α in severe malaria.^[29-31]

Infection intensity was found to be nonuniformly related to the pro-cytokines studied signifying that after induction, recruitment of these immune components is independent of parasite factors.

With negative socio-economic indices, the poverty level is high in Nigeria negatively affecting the standard of living, nutritional status, and susceptibility to infections resulting in a high rate of wasting in children, especially of rural dwellers as found in this study.

Altered platelet (platelet count $<150 \times 10^9/\text{L}$) and anemia (Hb <7 g/dL) were predominant in infected subjects, resulting from cytokine inhibition of erythropoiesis. Analysis of the hematological indices showed that IL-12 was negatively correlated with mean cell hemoglobin, granulocyte, and platelet in the test participants as IL-12 plays a significant role in the induction of Th1 immunity while IL-1 β has an early protective role in malaria.^[32] Platelets play a major role as first-line inflammatory mediators. This is evident from their capability to recruit early inflammatory cells such as neutrophils, granulocytes, and monocytes and even to exhibit direct antimicrobial activity, hence they might have delayed the immune functions of pro-inflammatory cytokines in malaria-infected subjects. However, cytotoxic

properties of blood platelets are induced by cytokines including interferon-gamma and TNF- α .^[33-36] Moreover, numerous studies have also demonstrated that nutritional status could modulate the immune response directed to malaria antigens^[7,37-39] and therefore to vaccine candidates in Nigeria.^[40-44]

The main findings from this study indicated that malnutrition may modulate Th1 response in *Pf* malaria, especially affecting the TNF- α production. However, IL-12 levels were also raised and have been linked to significantly higher levels of TNF- α . Susceptibility to malaria and clinical episodes is therefore associated with elevated TNF- α , of which high values of IL-12 are a major contributor. This should also be taken into consideration in designing vaccine trials in Nigeria as malnutrition influences patients' immunity to malaria and disease progression possibly via TNF- α .

CONCLUSION

Nutritional status is a key determinant of outcomes during infection with *Plasmodium falciparum*.

Limitation of study

The sample space was limited to one study site as funding was limited.

Acknowledgment

We acknowledge all the members of the staff of Ijede General Hospital for their cooperation during sample collection. Dr Faneye Dayo, at the Department of Microbiology, Nigerian Institute of Medical Research for the ELISA equipment support.

Financial support and sponsorship

The study was self-sponsored.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. World Health Organization. Global Health Statistics. Geneva: World Health Organization; 2013. p. 167.
2. Ajibaye O, Osuntoki AA, Balogun EO, Olukosi YA, Iwalokun BA, Oyeola KM, *et al.* Genetic polymorphisms in malaria vaccine candidate *Plasmodium falciparum* reticulocyte-binding protein homologue-5 among populations in Lagos, Nigeria. (BMC) Malar J 2020. doi:10.1186/s12936-019-3096-0.
3. Martorell R, Rivera J, Kaplowitz H, Pollitt E. Long-term consequences of growth retardation during early childhood. In: Hernandez AJ, editor. Human Growth: Basic and Clinical Aspects. Amsterdam: Elsevier Science; 1992. p. 143-9.
4. Fernando SD, Gunawardena DM, Bandara MR, De Silva D, Carter R, Mendis KN, *et al.* The impact of repeated malaria attacks on the school performance of children. Am J Trop Med Hyg 2003;69:582-8.
5. Branca F, Ferrari M. Impact of micronutrient deficiencies on growth: The stunting syndrome. Ann Nutr Metab 2002;46 Suppl 1:8-17.
6. Sumbele IU, Kimbi HK, Ndamukong Nyanga JL, Nweboho M, Anchang Kimbi JK, Lum E, *et al.* Malarial anaemia and anaemia severity in apparently healthy primary school children in urban and rural settings in the Mount Cameroon area: Cross sectional survey. PLoS One

- 2015;10:e0123549.
7. Fillol F, Sarr JB, Boulanger D, Cisse B, Sokhna C, Riveau G, *et al.* Impact of child malnutrition on the specific anti-*Plasmodium falciparum* antibody response. *Malar J* 2009;8:116.
8. Hassen HY, Ali JH. The association between chronic undernutrition and malaria among Ethiopian children aged 6-59 months: A facility-based case-control study. *S Afr J Child Health* 2016;10:63-7.
9. Pinstrup Andersen P, Burger S, Habicht JP. Protein-energy malnutrition. In: Jamison DT, Measham AR, Bobadilla JL, editors. *Disease Control Priorities in Developing Countries*. New York: Oxford University Press; 1993. p. 391-420.
10. Rice AL, Sacco L, Hyder A, Black RE. Malnutrition as an underlying cause of childhood deaths associated with infectious diseases in developing countries. *Bull World Health Organ* 2000;78:1207-21.
11. McGregor IA. Malaria: Nutritional implications. *Rev Infect Dis* 1982;4:798-804.
12. McGregor IA, Gilles HM, Walters JH, Davies AH, Pearson FA. Effects of heavy and repeated malarial infections on Gambian infants and children; effects of erythrocytic parasitization. *Br Med J* 1956;2:686-92.
13. McGregor IA. Malaria and nutrition. In: McGregor WW, editor. *Malaria – Principles and Practice of Malariology*. London: Churchill Livingstone; 1982. p. 753-68.
14. Gershwin M, Beach RS, Hurley L, *HandBook of Nutrition and Immunity*. Totowa, New Jersey: Humana Press; 2004.
15. Global Health Initiative: 2011-World Health Organization Report; 2011.
16. Ehrhardt S, Burchard GD, Mantel C, Cramer JP, Kaiser S, Kubo M, *et al.* Malaria, anemia, and malnutrition in African children – Defining intervention priorities. *J Infect Dis* 2006;194:108-14.
17. Wroczynska A, Nahorski A, Bakowska A, Pietkiewicz H. Cytokines and clinical manifestations of malaria in adults with severe and uncomplicated disease. *Int Marit Health* 2005;56:103-14.
18. Genton B, Al Yaman F, Ginny M, Taraika J, Alpers MP. Relation of anthropometry to malaria morbidity and immunity in Papua New Guinean children. *Am J Clin Nutr* 1998;68:734-41.
19. Müller O, Garenne M, Kouyaté B, Becher H. The association between protein-energy malnutrition, malaria morbidity and all-cause mortality in West African children. *Trop Med Int Health* 2003;8:507-11.
20. Müller O, Traoré C, Jahn A, Becher H. Severe anaemia in West African children: Malaria or malnutrition? *Lancet* 2003;361:86-7.
21. World Health Organization. Physical Status: The Use and Interpretation of Anthropometry. Report of WHO Expert Committee. WHO Technical Report Series, Geneva: World Health Organization; 1995. p. 321-44. Available from <https://apps.who.int/iris/handle/10665/37003>. [Last accessed on 2023 Oct 21].
22. Omosun YO, Anumudu CI, Adoro S, Odaibo AB, Sodeinde O, Holder AA, *et al.* Variation in the relationship between anti-MSP-1 (19) antibody response and age in children infected with *Plasmodium falciparum* during the dry and rainy seasons. *Acta Trop* 2005;95:233-47.
23. Sarr JB, Remoué F, Samb B, Dia I, Guindo S, Sow C, *et al.* Evaluation of antibody response to *Plasmodium falciparum* in children according to exposure of *Anopheles gambiae* s.l or *Anopheles funestus* vectors. *Malar J* 2007;6:117.
24. Jakobsen PH, Kurtzhals JA, Riley EM, Hviid L, Theander TG, Morris Jones S, *et al.* Antibody responses to rophtry-associated protein-1 (RAP-1) of *Plasmodium falciparum* parasites in humans from areas of different malaria endemicity. *Parasite Immunol* 1997;19:387-93.
25. Afolabi BM, Salako LA, Mafe AG, Ovwigho UB, Rabiou KA, Sanyaolu NO, *et al.* Malaria in the first 6 months of life in urban African infants with anemia. *Am J Trop Med Hyg* 2001;65:822-7.
26. Akanbi OM, Odaibo AB, Ademowo OG. Effect of antimalarial drugs and malaria infection on oxidative stress in pregnant women. *Afr J Reprod Health* 2010;14:209-12.
27. Mandel BK, Whit MR. *Lecture Note on the Infectious Diseases*. 4th ed. London: Blackwell Scientific Publications; 1984.
28. Zarkesh Esfahani H, Pockley AG, Wu Z, Hellewell PG, Weetman AP, Ross RJ. Leptin indirectly activates human neutrophils via induction of TNF-alpha. *J Immunol* 2004;172:1809-14.
29. Lyke KE, Burges R, Cissoko Y, Sangare L, Dao M, Diarra I, *et al.* Serum levels of the proinflammatory cytokines interleukin-1 beta (IL-1beta), IL-6, IL-8, IL-10, tumor necrosis factor alpha, and IL-12(p70) in Malian children with severe *Plasmodium falciparum* malaria and matched uncomplicated malaria or healthy controls. *Infect Immun* 2004;72:5630-7.
30. Kordes M, Matuschewski K, Hafalla JC. Caspase-1 activation of interleukin-1 β (IL-1 β) and IL-18 is dispensable for induction of experimental cerebral malaria. *Infect Immun* 2011;79:3633-41.
31. Ismaili J, van der Sande M, Holland MJ, Sambou I, Keita S, Allsopp C, *et al.* *Plasmodium falciparum* infection of the placenta affects newborn immune responses. *Clin Exp Immunol* 2003;133:414-21.
32. Mbugi EV, Meijerink M, Veenemans J, Jeurink PV, McCall M, Olomi RM, *et al.* Effect of nutrient deficiencies on *in vitro* Th1 and Th2 cytokine response of peripheral blood mononuclear cells to *Plasmodium falciparum* infection. *Malar J* 2010;9:162.
33. Lang A, Lahav M, Sakhnini E, Barshack I, Fidler HH, Avidan B, *et al.* Allicin inhibits spontaneous and TNF-alpha induced secretion of proinflammatory cytokines and chemokines from intestinal epithelial cells. *Clin Nutr* 2004;23:1199-208.
34. Bouchama A, al Sedairy S, Siddiqui S, Shail E, Rezeig M. Elevated pyrogenic cytokines in heatstroke. *Chest* 1993;104:1498-502.
35. Sprague AH, Khalil RA. Inflammatory cytokines in vascular dysfunction and vascular disease. *Biochem Pharmacol* 2009;78:539-52.
36. Panossian A, Davtyan T, Gukassyan N, Gukasova G, Mamikonyan G, Gabrielian E, *et al.* Effect of andrographolide and Kan Jang – Fixed combination of extract SHA-10 and extract SHE-3–on proliferation of human lymphocytes, production of cytokines and immune activation markers in the whole blood cells culture. *Phytomedicine* 2002;9:598-605.
37. Nyakeriga AM, Williams TN, Marsh K, Wambua S, Perlmann H, Perlmann P, *et al.* Cytokine mRNA expression and iron status in children living in a malaria endemic area. *Scand J Immunol* 2005;61:370-5.
38. Good MF, Xu H, Wykes M, Engwerda CR. Development and regulation of cell-mediated immune responses to the blood stages of malaria: Implications for vaccine research. *Annu Rev Immunol* 2005;23:69-99.
39. Moormann AM. How might infant and paediatric immune responses influence malaria vaccine efficacy? *Parasite Immunol* 2009;31:547-59.
40. Cortés A, Mellombo M, Mueller I, Benet A, Reeder JC, Anders RF. Geographical structure of diversity and differences between symptomatic and asymptomatic infections for *Plasmodium falciparum* vaccine candidate AMA1. *Infect Immun* 2003;71:1416-26.
41. Nair M, Hinds MG, Coley AM, Hodder AN, Foley M, Anders RF, *et al.* Structure of domain III of the blood-stage malaria vaccine candidate, *Plasmodium falciparum* apical membrane antigen 1 (AMA1). *J Mol Biol* 2002;322:741-53.
42. Ashley EA, Dhorda M, Fairhurst RM, Amaratunga C, Lim P, Suon S, *et al.* Spread of artemisinin resistance in *Plasmodium falciparum* malaria. *N Engl J Med* 2014;371:411-23.
43. Ferreira MU, Hartl DL. *Plasmodium falciparum*: Worldwide sequence diversity and evolution of the malaria vaccine candidate merozoite surface protein-2 (MSP-2). *Exp Parasitol* 2007;115:32-40.
44. Polley SD, Tetteh KK, Lloyd JM, Akpogheneta OJ, Greenwood BM, Bojang KA, *et al.* *Plasmodium falciparum* merozoite surface protein 3 is a target of allele-specific immunity and alleles are maintained by natural selection. *J Infect Dis* 2007;195:279-87.