

5

ILORIN JOURNAL OF MANAGEMENT SCIENCES



ISSN 2408-655X

Published by:
FACULTY OF MANAGEMENT SCIENCES
UNIVERSITY OF ILORIN

e-mail: editorijms@hotmail.co.uk

www.unilorin.edu.ng

Volume 1, No. 2,
December 2014.

Table of CONTENTS

Stock splits and stock dividends: implications for bid ask spread components OKE, Lukman Adebayo, TIJANI, Oladipupo Muhrtala, MUSTAPHA, Abdulrasaq & AFOLABI, Hafsat Olatanwa	1
Effects of advertising on products sales volume in Nigeria Bottling Company Plc IMOUOKHOME, Ebun Omoniyi, AUN, Isaac Iortimbir & ABDUL, Falilat Ajoke.....	12
SMEs owners' Perception on the value of Accounting Information for economic decision making in Ilorin Metropolis, Nigeria MUSTAPHA, Abdulrasaq	24
Corporate profitability and liquidity: A study of Listed Nigerian Deposit Money Banks AKANET, Solomon	35
Finance, Education and Innovativeness as combined SMEs Growth Trajectories: Panel Data Analysis TUKUR, Mustapha Namakka & KAMBA, Kabir Muhammad	46
Corporate shareholdings structure and dividend payout ratio of Listed Chemical and Paints Companies in Nigeria GUGONG, Benjamin Kumai & MUSA, Adeiza Farouk...	61
Role of Information Technology in the management of Small and Medium Scale Enterprises in Ilorin Metropolis KADIRI, Ismaila Bolarinwa	72
Ethics in research with human Participants: Issues and Implications KURFI, Mustapha Hashim	79
Corporate Governance and Performance in the Banking and Food Industries in Nigeria. ADEWOLE, Clement	89
Audit partner preferences and risk assessment: Implications for Auditor Program Plans TIJANI, Oladipupo Muhrtala (ACCA), OKE, Lukman Adebayo (ACA) BADMUS, Dauda Abiola (FCA) & DAUDA, Folorunsho Quadri (ACA, FPFM London).....	102
Entrepreneurial Capital, Personal reputation and Performance of Small Service Firms KAMBA, Kabir Muhammad & TUKUR, Mustapha Namakka.....	118
An analysis of conventional concepts of Accounting from an Islamic Perspective MAHMOUD, Aisha Hamman.....	129
Students' usage of Technology mediated Platforms for Communication: Case Study of University of Ilorin Students MEJABI, O.V. and AJIBOYE, G.S.....	149
Spatio-temporal Patterns of Nitrate Concentration in Rainwater in Ilorin, Kwara State Nigeria. IROYE, K.A, Olanrewaju, R.M & OYELAKIN R.I.....	169

SPATIO-TEMPORAL PATTERNS OF NITRATE CONCENTRATION IN RAINWATER IN ILORIN, KWARA STATE NIGERIA.

By

Iroye, K.A, Olanrewaju, R.M and Oyelakin R.I
Department of Geography and Environmental Management
University of Ilorin, Nigeria.
Email: iroyekayode@yahoo.com

Abstract

This study examined the spatio-temporal pattern of nitrate concentration in rainwater in Ilorin, Kwara State, Nigeria. Thirty five rainwater samples were collected from each of the five different land-uses using sterilized 75 centiliters plastic bottles on event basis between the onset and cessation of rainfall (March 4th – October 24th, 2013) in the study area. The collected water samples were subsequently analyzed for nitrate concentration in University of Ilorin Chemistry laboratory using sodium salicylate (colorimetric) method. Result obtained indicates high variability in both spatial and temporal patterns of nitrate concentration in the study area. While the temporal pattern revealed high concentrations in the months of March, April, July and September; Oja-Oba, Asa Dam and Oke-Oyi areas of the city exhibits high degree of concentrations in that order spatially. Likely reasons for the observed pattern of concentrations were discussed while suggestions towards reducing the rate of pollution to biologically safe levels were presented.

Keywords: Nitrate, N-Nitroso, Pollution, Carcinogens, Land-use.

1.0 Introduction

The utilization of rainwater for domestic activities including cooking is a common practice in Africa (Olorunfemi and Arohunsoro, 2000) during the rainy season and has over the years been advocated, especially in less developed countries (NIRCCWSS, 1988; UNESCAP, 1989 ; Morgan, 1990). This activity which is commonly referred to as rain harvesting is being encouraged because of the failure of modern water supply system and increasing pollution of surface water. However, water from this source which is directly collected from the atmosphere is now being increasingly confronted with problem of quality induced by gases and particulate matter released into the earth's atmosphere as consequence of man's activities.

Among such atmospheric pollutant is nitrate, a pollutant which enter rainwater by the reaction of nitrogen oxides (NO_x species) with water to produce HNO_3 , which disassociate into H^+ and NO_3^- (MARYP, 1999).

Though NO_x species come from some natural sources such as lighting and volcanic gasses, greater amount of nitrates (more than 95%) come from anthropogenic sources such as automobiles and fossil fuel combustion in homes, vehicles and industries (Botkin *et al.* 1995 ;Long, 2003). According to Ubuoh, *et. al.* (2012), such sources could be point, non-point or mixed in nature. Pollutants once introduced into the atmosphere can be removed through mechanisms of dispersion, rainout or scavenging leading to rainwater contamination (Akhionbare, 2009). This is usually through dissolution of such gases or particles in the air or impaction into pre-existing liquid droplet (Carole *et al.*, 2002). And because such processes can either occur in a cloud or below it during rain, the levels of such gases or particles in resultant rainfall is thus greatly increased, hence the production of relatively strong nitric acid (HNO_3) in regions polluted with oxides of nitrogen.

Worldwide, there is a limit to the amount of nitrate that can be found in drinking water. The World Health Organization (2011) and the United State Environmental Protection Agency (2007) have both set a limit of 10mg/l. Nitrate in drinking water at levels above this limit poses a threat to man. When present in high concentrations in bottle-fed infants, they may give rise to a condition known as methemoglobinemia or blue-baby syndrome. The consequence of this is shortness of breath (cyanosis), blueness of skin and at times death (BGS, 2003). Its high concentration also poses potential health risk to pregnant women (Jennings and Sneed, 1996; Kelter, *et al.*, 1997; Kempster, *et al.*, 1997) and can also cause recurrent acute diarrhea (Gupta, *et al.*, 2001). Ingested nitrate from dietary sources and drinking water can be converted to nitrite and ultimately to N-nitroso compounds, many of which are known to be carcinogens (Sandor, *et. al.*, 2001; Weyer, *et al.*, 2006). In freshwater or estuarine systems close to land, nitrate can reach high levels that can potentially inhibit growth, impair the immune system and cause stress in some aquatic species and eventually cause death. In most cases, excess nitrate concentrations in aquatic systems results from primary sources such as rainfall and surface runoff from agricultural or landscaped areas that have received excess nitrate fertilizer. This is called eutrophication and can lead to algae blooms as well as water anoxia and dead zones. These blooms may cause other changes to ecosystem function, favouring some groups of organisms over others.

As a result of its dangerous effects when present in high concentration, nitrate forms a component of total dissolved a solid that is widely used as an indicator of water quality. Because of the aforementioned problems induced by high nitrate concentration in water and the fact that rainwater chemistry is highly variable both spatially and temporarily; there is need for investigation of this nature which is intended to shed more light on what the future hold in stock concerning the impact of rainwater pollution on both human and animal populations. Identification of pollution sources, transport characteristics and predicted concentrations according to Walker *et al.*, (2000) will no doubt lead to physical or chemical controls that can be used for the development of sound environmental management policies to prevent future damage to sensitive aquatic and terrestrial ecosystem.

2.0 The Study Area

Ilorin, the capital city of Kwara State Nigeria is the study area in this research (Fig I). The city which is located between latitudes $8^{\circ}24'$ and $8^{\circ}36'$ north of the equator and between longitudes $4^{\circ}10'$ and $4^{\circ}36'$ east of the Greenwich meridian covers an area of about 150km^2 . The city is located in the transitional zone between the forest and guinea savannah region of Nigeria.




Fig1 : Street Map of the Study Area Showing Data Collection Points

Source: Author's Field Work (2014)

The climate of Ilorin results from the influence of two trade winds prevailing over Nigeria. These are the Tropical Maritime Airmass which originates from the Atlantic Ocean and the Tropical Continental Airmass which originates from Shara desert. The Tropical Maritime Airmass is prevalent in the city between the months of March and October while the Tropical Continental Airmass is prevalent between November and February. These two wind systems produces two climatic seasons of wet and dry seasons. Annual rainfall in Ilorin is around 1,200mm with peak periods in July and September (Olaniran, 2002). Rainfall type which is conventional in nature exhibits high variability, both temporarily and spatially (Ajadi, 1996). Mean monthly temperature in Ilorin is generally high throughout the year, varying between 25° and 28°C (Oyegun, 1983). The average relative humidity in the city in wet season is between 75 and 80% while in dry season, it is about 65% (Tinuoye, 1990). Urbanization and industrialization process are fast taking place within and around the city (Olanrewaju, 2009) with their consequence effects on air pollution and rainwater contamination.

3.0 Methodology

This research work was based on data collected directly from the field. Rainwater samples were collected from five (5) locations within and around Ilorin metropolis based on landuse pattern. Table 1 shows the grid reference and altitude of the sampling points taken with hand-held Global Positioning System (GPS) receiver (Garmin GPS 12 Personal Navigator).

Table 1: Grid Reference and Altitude of Sampling Locations

Sampling Location	Landuse Type	Latitude	Longitude	Altitude (m)
Asa Dam	Industrial	08°27'	004°32'	327
Oja-Oba	Residential	08°30'	004°31'	320
Oke-Oyi	Agricultural	08°36'	004°46'	283
Ganmo	Rural	08°26'	004°36'	330
Oke-Odo	Commercial	08°28'	004°37'	352

Source: Authors' Fieldwork, 2013.

Rainwater samples were collected at each sampling points with a sterilized 75 centiliter plastic polyethylene bottles, filled to the brim and cocked to avoid an air column. In all, thirty-five (35) storm events were sampled in each of the five (5) landuse types between the periods of onset and cessation of rainy season (March 4th – October 24th, 2013) on event basis. The duration of collection however varied from a few minutes to about an hour depending on the intensity of the rainfall.

Rainfall collectors used in this investigation were placed on 2 meter platforms and fitted with sieves to prevent impurities from entering the water samples. The collected samples were subsequently analyzed for nitrate concentrations at University of Ilorin Chemistry laboratory using Sodium (Colorimetric) method. This analytical technique is preferred to the distillation method because it is simple and generates fast results. It can present result in the range of 0.2-20mg/l nitrate in water samples. Table 2 shows the analytical procedure followed in conducting the test.

Table 2: Analytical Procedure used in Testing for Nitrate Concentration

Method

Procedure

Sodium Salicylate

(Colorimetric) Technique

Fluorination (to remove interfering organic and inorganic substances): 20ml of sample (20) ml was added into the sample to give a concentration of 100mg/l. The pH of the samples is adjusted to between 11 and 11.5 with 40% NaOH. The concentration was raised with magnetic stirrer and allowed to stand for five minutes so that flocculated particles may settle. It was then filtered to discard the first portion of the filtrate.

Extraction and Drying: 2ml of the filtrate was accurately pipette into a 25ml centrifuge tube. 2ml of 1% sodium salicylate was added and vigorously shaken. The mixture was dried for at least thirty minutes in a drying oven at 105°C.

Colour Development: The sample is removed from oven, allowed to cool and 2ml of concentrated HCl, was added quickly, mixed well by swirling. It was allowed to stand for 10 – 15 minutes but checked occasionally to ensure development of an colour. When color, 10ml of water was added into the sample vial. The vial was shaken the pipette tip against the dish wall to avoid splashing that can lead to loss of part of the mixture. 10ml of sodium hydroxide/potassium hydroxide solution (10N/10N) was added into the concentration. If nitrate is present, a yellow colour develops immediately. It was then again and allows the solution to cool at room temperature for 10 minutes.

Calibration Standard: From the standard substance, 1ml = 2mg/l, 2.5ml = 2.5mg/l was pipette into 10ml bottles. 1ml of 1% sodium salicylate was added to each bottle. The standard was treated as the samples throughout the procedures above. A blank solution containing all reagents but without the sample, and the calibration substance were treated in the same manner.

Extrapolation from Calibration Standard: The absorbance of the standard solution was measured in a blank and a series of standards at 410nm, reading obtained at 410nm. Extrapolation was made from standard value.

Standard (mg/l)	1	2	4	6	8	10	12	14	16	18	20
Values	0.111	0.133	0.166	0.200	0.233	0.266	0.299	0.333	0.366	0.399	0.433

Source: Laboratory Manual, Dept. of Chemistry, University of Ilorin (2013)

4.0 Observations and Discussion

Table 3 present the results of the laboratory analysis on nitrate concentration in the sampled rainwater. The data revealed that nitrate concentration in rainwater in Ilorin exhibits high variability, both spatially and temporally in the city.

Table 3: Laboratory Result of Rainwater Samples

Sample Locations Dates	1 Ganmo	2 Oke-Odo	3 Oke-Oyi	4 Asa-Dam	5 Oja-Oba	Mean Value for Ilorin The Study Area
March 04, 2013	17.25	16.62	16.16	16.96	19.64	17.33
March 16, 2013	17.31	15.35	16.05	16.62	18.40	16.75
March 30, 2013	15.67	14.70	14.91	16.85	17.82	15.99
April 07, 2013	14.16	13.56	14.48	16.28	15.89	14.87
April 15, 2013	13.78	14.10	14.21	14.81	15.18	14.42
April 19, 2013	14.91	14.27	14.00	14.43	16.05	14.73
April 22, 2013	13.72	13.89	16.21	14.05	15.45	14.66
April 25, 2013	8.06	16.28	15.92	14.21	15.51	14.00
April 28, 2013	13.89	8.13	7.73	14.97	16.00	12.14
April 30, 2013	14.59	8.16	7.96	13.94	17.54	12.44
May 01, 2013	15.50	10.54	15.78	7.73	15.23	14.96
May 04, 2013	16.00	15.36	10.29	13.94	17.02	14.52
May 11, 2013	10.24	10.00	10.39	10.14	16.14	11.38
May 14, 2013	8.16	8.06	7.83	7.90	13.94	12.42
May 20, 2013	14.10	14.00	10.29	10.44	14.54	12.67
May 25, 2013	9.90	9.45	9.16	6.88	10.29	9.94
May 28, 2013	9.55	6.92	6.64	6.16	14.97	8.85
June 11, 2013	7.52	6.00	14.05	4.17	15.18	9.38
June 14, 2013	2.09	4.17	6.32	6.20	6.36	5.03
June 20, 2013	2.41	2.37	6.24	10.19	15.29	7.30
June 23, 2013	2.14	1.72	2.00	4.23	6.08	3.23
June 29, 2013	2.21	2.07	4.08	6.08	2.25	3.34
July 04, 2013	0.23	1.87	2.16	1.11	4.05	1.88
July 14, 2013	1.31	1.40	1.81	1.26	2.19	1.59
July 20, 2013	1.78	1.56	6.64	10.39	16.00	7.27
July 24, 2013	7.12	16.70	10.29	16.49	15.43	13.21
Sept 01, 2013	1.42	1.00	1.04	1.42	1.13	1.20
Sept 05, 2013	6.64	9.50	6.24	9.50	11.92	8.76
Sept 12, 2013	1.87	2.23	1.90	10.05	21.70	9.15
Sept 14, 2013	0.81	2.09	1.13	2.03	1.24	1.46
October 01, 2013	9.50	6.48	6.88	9.65	9.90	8.48
October 09, 2013	6.60	3.91	3.67	7.12	9.55	5.81
October 14, 2013	4.14	3.97	4.05	6.68	9.06	5.68
October 19, 2013	6.36	4.11	3.37	2.05	6.52	4.48
October 24, 2013	3.85	3.62	5.84	4.05	6.96	4.86
TOTAL	294.79	284.96	305.72	328.98	430.37	326.80
RANGE	17.08	15.70	15.12	15.85	20.57	15.87
MEAN	8.42	8.12	8.78	9.39	12.30	9.34
S.D	5.62	5.42	5.05	5.03	5.66	5.36
C.V	66.74	66.75	59.76	56.33	46.01	59.12

Note: Values and in ppm (parts per million)

Source: Authors' Fieldwork, 2013.

Nitrate concentration in rainwater in the study area varied between 0.23ppm observed in Ganmo on July 4th and 21.7 ppm observed in Oja-Oba on 12th of September. All the five (5) sampling points generally recorded high nitrate concentration values in the first 7 rainfall events which span a period of almost 7 weeks from the onset of rainy season in the study area. However, the high concentration values persisted till 11th and 17th weeks respectively in Asa-Dam and Oja-Oba areas of the city from the onset of rainy season. While the general high nitrate concentration in rainwater in the first seven weeks in the rainy season can be attributed to the build-up effect of nitrogen in the atmosphere during the dry season; the persistent high level concentration in Asa-dam and Oja-Oba for additional few weeks may not be unconnected with land use activities in these two areas.

While Asa-Dam is an industrial region, with presence of large sawmill where saw dust is always being burnt, Oja-Oba is urban residential area with high vehicular movement. The high rate of emission of gases from wood burning and combustion of fossil fuels in these two areas may have contributed to the persistent high nitrate concentration of rainwater in the areas. Not only that, the high nitrate concentration in rain water in these two sampling sites may also be linked to the closeness of Asa-Dam to Oja-Oba. Studies such as Morales, *et. al.*, 1998 and Russel, *et. al.* 1998 ; Walker, *et. al.* 2000; Sarah, 2006; Uboh, *et. al.* 2012 have linked degree of atmospheric pollution in a particular region to the direction and degree of movement of air masses. According to Carole, *et. al.* (2012) atmospheric pollutants may come from any direction, both local and distant areas following different pathways. Thus, Russel *et. al.*, (1998) classified storm trajectories and their associated pollutants into five groups viz: westerly, southerly easterly, northwesterly and southwesterly. According to their observation, the southwesterly trajectory was associated with high concentration of nitrate in their study area. Their findings thus indicate that source region information can be used to determine how much nitrogen is deposited in the atmosphere, what processes produce the nitrogen and where the nitrogen originated.

The relatively high mean nitrate concentration in rainwater at Oke-Oyi may be the result of intensive agricultural production in the area. Oke-Oyi houses plantation agricultural farm where nitrogen rich fertilizer is being used in improving soil fertility. Bocher (1995) observed that there are five fates for such nitrogen once applied to crop; according to him, it may be taken up by plant, stored in the soil, lost to groundwater, lost to runoff or lost to atmosphere where it mixes with rainwater. The generally high nitrate concentrations in rainwater in the study area however reduced in all the sampling points to low level in the months of July and August before picking up in the month of September when it rose to its peak of 21.05 and 21.13ppm in Oja-Oba and Asa-Dam respectively. By late October, the concentration of nitrate in rainwater had also significantly reduced with the cessation of rainfall in all the five sampling points in the study area (Fig. 2).




Fig 2: Temporal Variation in Nitrate Concentration Under Different Land-uses

Source: Authors' Fieldwork, 2014.

The sharp increase in level of nitrate concentration in rainwater observed in September may not be unconnected with the August break in rainfall in the study area. Absence of significant amount of rain in August in the study area might have aided the build-up of nitrogen in the atmosphere which thereafter is washed down by September rainfall. Figure 3 depicts rain water nitrate exceedance map for the study area.




Fig. 3: Nitrate Exceedance Map for the Study Area

Source: Authors' Fieldwork 2014.

Though this particular research did not examine the relationship between rainfall amount and nitrate concentration, studies such as Russel *et al* (1998) ; Walker *et al* (2000) and Dayan and Lamb (2003); have explained that the amount of precipitation is a factor affecting concentration of nitrogen analytes in atmospheric deposition. According to these studies, the term 'washout' is used in explaining the reason why there seems to be smaller concentrations of analytes when precipitation amounts are large.

This according to Sarah *et. al.*, (2006) is because high volume of water dilutes the concentration of the pollutant.

5.0 Conclusion and Recommendations

Nitrate concentration in rainwater in Ilorin exhibits spatio-temporal patterns. While temporal pattern reveals high concentration levels in the months of March, April, July and September; spatially, the concentrations were observed to be high under urban residential, industrial and agricultural land uses. This result presents great danger most especially for residents in high concentrations areas who may use such contaminated water for domestic activities, most especially for cooking. While ingested nitrate from dietary sources and drinking water by man can be converted to nitrite and ultimately to N-nitroso compounds, many of which are known to be carcinogens; high nitrate level in aquatic ecosystem potentially inhibit growth, causes stress and may result in death. Though excess nitrate concentrations in aquatic systems most times results from surface runoff from agricultural land that have received excess nitrate fertilizer, heavily polluted rainfall from atmospheric nitrogen can also impair aquatic systems leading to algae blooms, water anoxia and dead zones. Because nitrate cannot be completely removed from rainwater, efforts should be directed by government agencies responsible towards reducing its rate of pollution to biologically safe levels; this can be through:

- i. Legislation aimed at tackling the sources of pollution.
- ii. Public enlightenment programmes in various mass media on biomass burning, fossil fuel combustion and excessive application of fertilizer containing nitrogen on agricultural farmlands, and,
- iii. Development of necessary data base as currently being done in this study to identify and effectively manage the high risk areas.

References

- Ajadi, B.S. (1996); Pattern of Water Supply in Ilorin City, Nigeria. *An Unpublished B.Sc Project*, Department of Geography, University of Ilorin.
- Akhionbare, S.M.O (2009) *The Environment: Concepts, Issues and Control of Pollution* MC Publishers, Nnewi Nigeria 434pp.
- BGS, (2003); British Geological Survey; *Water Aid Fact Sheet*, London.
- Botkin, D.B. and Keller, E.A. (1995); *Environmental Science: Earth as a Living Planet*; John Wiley and Sons, Inc., New York; pp. 453
- Carole, B; Pierre, C; Raoul, C. and Enrique, B. (2002) Occurrence of Pesticides in the Atmosphere in France. *Agronomie* 22: 35-40.
- Dayan, U. and Lamb, D (2003) Meteorological Indicators of Summer Precipitation Chemistry in Central Pennysylvania. *Atmospheric Environment* 37: 1045-1055.
- Gupta S.K, Gupta R.C, Gupta A.B, Seth A.K, Bassin J.K, Gupta A, Sharma MI (2001) Recurrent Diarrhea in Children Living in Areas with High Levels of Nitrate in Drinking Water. *Archives of Environmental Health* July issue. Retrieved on February 18, 2014 from http://www.findarticles.com/p/articles/mi_m0907/is_4_56/ai_79124886
- Jennings G.D and Sneed R.E (1996); Nitrate in Drinking Water; North Carolina Cooperative Extension Service, *Publication No.* AG 473-4.

- Kelter P.B, Grundman J, Hage D.S, Carr J.D (1997); A Discussion of Water Pollution in the US and Mexico with High School Laboratory Activities for Analysis of Lead, Anthrazine and Nitrate. *J. Chem. Educ.* 74 (284): 1413-1418.
- Kempster P.L, Van Vliet H.R, Kuhn A. (1997); The Need for Guidelines to Bridge the Gap between Ideal Drinking-Water Quality and that Quality which is Practicably Available and Acceptable. *Water SA* 23 (40): 163-167.
- Long, M.S (2003) Atmospheric Deposition in Southeastern North Carolina and its Impact on the Cape Fear Estuary, Unpublished *M.Sc Dissertation* UNC Wilmington NC.
- Monitoring Acid Rain Youth Program (MARYP) "1999"; "Rainfall Chemistry." On-line: <http://www.trentu.ca/academic/biology/maryp/1996finalreport/rain.html> Accessed date: 14 July, 2013.
- Morales, J.A., Bifano, C., Escalona, A. (1998); Atmospheric Deposition of SO_4 , S and $(\text{NH}_4 + \text{NO}_3) - \text{N}$ at Two Rural Sites in the Western Maracaibo Lake Basin. *Atmospheric environment*. 32(437): 3051-3058.
- Morgan P. (1990) *Rainwater Harvesting in Rural Water Supplies and Sanitation*: Zimbabwe's BLAIR Research Laboratory, Macmillan Educ. London 225-234.
- Netherlands International Reference Centre for Community Water Supply and Sanitation (1988) *Rainwater Harvesting in Small Community Water Supplies*. In: Hofers, S.H (ed) *Technology of Small Water Supply Systems in Developing Countries*. John Wiley Publication pp. 59 – 73.
- Olaniran (2002) Rainfall Anomalies in Nigeria: The Contemporary Understanding 55th *Inaugural Lecture*, University Press, Ilorin. 66pp.
- Olanrewaju, R.M (2009) Climate Effect of Urbanization in a city of Developing Country: the Case Study of Ilorin, Kwara State, Nigeria *Ethiopian Journal of Environmental Studies and Management* 2 (2): 67-72.
- Olorunfemi J.F and Arohunsoro S.J (2000) Rainwater Harvesting for Rural Domestic Water Supply in Nigeria. *Geo-Studies Forum* 1 (1 and 2): 124-130.
- Oyegun, R.O (1983) *Water Resources Management in Kwara State Nigeria*. Matanmi Publishers, Ilorin.
- Russell, K.M, Galloway, J.N., Macko, S.A., Moody, J.L., and Scudlark, J.R. (1998); Sources of Nitrogen in Wet Deposition to the Chesapeake Bay Region; *Atmospheric Environment*. 32 (14/15): 2453-2465.
- Sarah, B.J Douglas, W.G., Michael, M.B and Joan, W. (2006) The Effects of Local Weather Patterns on Nitrate and Sulfate Rainwater Concentrations in Wilmington, North Caroline. *The North Carolina Geographer* 14: 29-38.
- Tinuoye A. (1990); "Climatic Condition in Dry and Wet Years at Ilorin"; *Unpublished B.Sc. Project Work*, University of Ilorin.
- Ubuoh E. A., Akhionbare, W.N., Akhionbare, S.M.O. and Ikhile, C.I. (2012); Evaluation of the Spatial Variation of Rainwater Quality in Parts of Akwa Ibom State of Nigeria Using Chloropleth Map. *American Based Research Journal*, 1 (2), 10 – 28.
- UNESCO (1989) Rainwater Harvesting Techniques: Prospects for their Application in Developing Countries. *Water Resources Series* 63:101-118.
- Walker, J.T., Ancja, V.P., and Dickey, D.A. (2000); Atmospheric Transport and Wet Deposition of Ammonium in North Carolina, *Atmospheric Environment*. 34 (310), 3407-3418.
- World Health Organization (2011) Nitrate and Nitrite in Drinking Water *Background Document for Development of WHO Guidelines for Drinking Water Quality* WHO/SDE/WSH/07.01/16/ Rev.