

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/326693362>

Growth pattern and condition factor of seven freshwater fish species from Jebba Lake, north-central Nigeria

Article · June 2018

DOI: 10.22034/iji.v5i2.272

CITATIONS

0

READS

2

7 authors, including:



Segun Olayinka Oladipo

Kwara State University

12 PUBLICATIONS **1** CITATION

[SEE PROFILE](#)



Lotanna Micah Nneji

Kunming Institute of Zoology CAS

12 PUBLICATIONS **6** CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Biodiversity Assesment of North-Central Fishes [View project](#)



Biodiversity assessment and conservation genetics of Nigerian freshwater fishes [View project](#)

Short Communication

Growth pattern and condition factor of seven freshwater fish species from Jebba Lake, north-central Nigeria

Segun Olayinka OLADIPO^{1,2*}, Lotanna M. NNEJI³, Abass Toba ANIFOWOSHE², Ifeanyi C. NNEJI⁴,
Oluyinka A. IYIOLA², Adeniyi C. ADEOLA³, Moshood K. MUSTAPHA²

¹Department of Biosciences and Biotechnology, Kwara State University, Malete, Kwara State, Nigeria.

²Department of Zoology, Faculty of Life Sciences, University of Ilorin, Ilorin, Kwara State, Nigeria.

³Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, China.

⁴Department of Biological Sciences, University of Abuja, Abuja, Nigeria.

*Email: olayinka_sgn@yahoo.co.uk

Abstract: Length-weight relationship and condition factor of seven freshwater fish species from Jebba Lake (Nigeria) were investigated. A total of 576 individuals were caught with hook and line, cast and gill nets from up and downstream sections of the lake. Sampling covered dry and raining seasons spanning across April to September 2017. The growth exponent b values of the studied species ranged between 2.23-3.89 and the condition factor between 0.47-1.61.

Keywords: *Barbus*, *Chrysichthys*, *Hydrocynus*, Length weight relationship, *Oreochromis*.

Citation: Oladipo, S.O.; Nneji, L.M.; Anifowoshe, A.T.; Nneji, I.C.; Iyiola, O.A., Adeola, A.C. & Mustapha, M.K. 2018. Growth pattern and condition factor of seven freshwater fish species from Jebba Lake, north-central Nigeria. Iranian Journal of Ichthyology 5(2): 167-172.

Introduction

Length-weight relationship (LWR) and condition factor (K) of fishes are fundamental morphometric parameters in fish ecology and research. They are valuable tools for investigating growth rate, age structure, feeding intensity and relative well-being of ichthyofaunal communities (Atama et al. 2013; Sossoukpe et al. 2013; Abu & Agarin 2016; Irom et al. 2017). Thus, an assessment of these parameters is useful for deducing inference on the general well-being of the fish communities (Ecoutin et al. 2005; Ayoade & Ikulala 2007).

Jebba Lake of Niger River Watershed, Jebba, Kwara State, Nigeria, is rich in fish biodiversity (Adelakun et al. 2017). To date, there are limited information on the LWRs and K of fish species of Jebba Lake. To advance knowledge on this, we examine the LWRs and K of some selected fish

species from the lake, and further compare the growth pattern and K of fishes from up and downstream sections of the lake.

Materials and Methods

Study area: Jebba Lake is located in North-Central Nigeria between latitudes 4°34'12"-4°43'48"E. This region is characterized mainly by two seasons- dry and rainy (Adelakun et al. 2017). The maximum length and width of the lake is about 134km and 24.1km, respectively, having surface area of 1270km² and catchment area of 1.6×10⁶km² (Obot 1989). Our study covered up and down stream sections (8.7km apart) of Jebba Lake notable for fishing.

Fish collection: We selected fish species based on its commercial importance and availability. In sum, our study includes seven commercially important fish

species namely *Auchenoglanis biscutatus* (Geoffroy Saint-Hilaire, 1809), *Barbus occidentalis* (Boulenger, 1911), *Lates niloticus* (Linne, 1762), *Citharinus citharus* (Geoffroy Saint-Hilaire, 1809), *Oreochromis niloticus* (Linne, 1758), *Chrysichthys nigrodigitatus* (Lacepede, 1803), and *Hydrocynus vittatus* (Castelnau, 1861). Samples were collected bimonthly from April to September 2017 spanning across dry and rainy seasons. For the collection of fish samples, three commercial fishing gears including gill net, hook and line, and cast net were used. Following Araoye (1999), the gears were set at the surface and bottom water. Our sampling of 576 individuals covered up- and downstream sections of the lake.

The selected specimens were preserved in a cooler with ice blocks and transported to the Laboratories in Department of Bioscience and Biotechnology, Kwara State University, Malete, Nigeria and Department of Zoology, University of Ilorin, Nigeria. Fish specimens were sorted and identified following fish identification keys of Olaosebikan and Raji (1998) and Idodo-Umeh (2003). Additional species identification and verification were sought from two trained taxonomists at the Department of Zoology, University of Ilorin, Nigeria.

Length-Weight Relationship and Fulton's Condition Factor (K): We measured the Total (TL) and Standard Lengths (SL) of each specimen to nearest 0.10cm using a measuring tape. We also assessed the body weight (BW) using a digital scale to nearest 0.10g. The equation: $W=aL^b$ was used to calculate length-weight relationship of the fishes (Bagenal 1978). Where W=the observed total weights of the fishes; L= the observed total length, *b* is the slope and *a* is the intercept on the length axis. Further, the equation above was linearised by a logarithmic transformation to give a straight-line relationship: $\text{Log } W=\text{Log } a+b\text{Log } L$. We plotted $\text{Log } 10W$ against $\text{Log } 10L$. We obtained the regression coefficient (*b*), and the intercept ($\text{Log } a$) on the Y axis.

The condition factor (*K*) was computed from the equation: $K=100W/L^3$, Where W= the observed total




Fig.1. Map of up and downstream sections of Jebba Lake showing the sampling points.

weight for each fish; L=the observed total length for each fish; *K*=the Condition Factor. The data was transformed into logarithm using Excel and the Log-Log graphs showing the regression analysis was performed using PAST (Hammer 2011).

Results and Discussion

The higher mean TL values were recorded in *H. vittatus*, *C. citharus* and *L. niloticus* collected from the upstream section, while in downstream, higher mean TL values were observed in *B. occidentalis*, *C. nigrodigitatus* and *A. biscutatus* (Table 1). Mean weight values of the fish specimens range between 72.90-276.70g (upstream) and 115.20-320.80g (downstream), respectively. The highest mean weight values were recorded in *L. niloticus* (upstream) and *C. citharus* (downstream). Compared to other fish species, *B. occidentalis* (upstream) and *L. niloticus* (downstream) had lower mean weight values (Table

Table 1. Morphometric, Length–Weight Relationships Parameters and Condition Factors (K) of Selected Species from Jebba Lake, North-Central Nigeria.

Species	Location	Sex	N	Length (min-max)	Length (mean±SD)	Weight (min-max)	Weight (mean±SD)	a	b	r ²	K
<i>Auchenoglanis biscutatus</i>	Upstream	M	15	16-36	23.50±1.20	333-400	149.40±23.40	-1.96	2.96	0.94	0.98
		F	22	16.5-34	22.50±1.20	39-450	154.40±19.20	-2.18	3.16	0.95	1.09
<i>Barbus occidentalis</i>	Downstream	M	18	14-36	24.10±1.00	35-430	198.30±15.20	-1.35	2.57	0.93	1.09
		F	20	15-36	25.10±1.00	40-480	206.50±31.20	-1.60	2.78	0.94	1.26
		M	22	19-26	21.70±0.50	40-130	74.40±7.30	-3.75	3.18	0.90	0.68
<i>Oreochromis niloticus</i>	Upstream	M	18	20-26	21.90±0.25	40-132	70.40±3.20	-3.19	3.76	0.89	0.69
		F	21	22-35	28.50±0.70	30-280	212.60±15.40	-1.36	2.91	0.90	0.73
		M	17	24-34	30.50±0.70	30-300	215.60±15.90	-1.73	2.75	0.96	0.79
<i>Chrysichthys nigrodigitatus</i>	Downstream	M	19	13-29	19.90±0.90	20-340	133.30±18.60	-2.18	3.26	0.90	1.48
		F	25	15-28	21.00±0.74	20-360	140.00±12.40	-2.56	3.54	0.96	1.32
		M	18	14-24	19.80±0.80	17-250	129.20±16.90	-3.04	3.89	0.98	1.32
<i>Hydrocynus vittatus</i>	Upstream	F	21	13-22	16.40±0.62	15-220	104.50±10.90	-2.25	3.35	0.98	1.61
		M	21	12.0-37	19.80±0.50	23-200	68.00±5.20	-1.97	2.86	0.95	0.67
		F	20	15.5-39	22.60±1.10	27-215	72.90±8.30	-1.79	2.75	0.93	0.72
<i>Citharinus citharus</i>	Downstream	M	17	19-33	23.00±0.25	60-305	128.40±12.50	-0.95	3.03	0.93	0.59
		F	28	21.5-37	28.40±1.10	60-344	168.60±17.80	-0.92	3.03	0.93	0.64
		M	13	20-38	26.50±1.05	55-275	100.00±7.40	-2.64	3.22	0.99	0.47
<i>Lates niloticus</i>	Upstream	F	30	22-38	27.20±1.10	50-275	111.00±16.40	-2.61	3.21	0.99	0.49
		M	22	14-37.5	25.10±2.30	20-268	119.60±14.10	-1.10	2.23	0.98	0.74
		F	31	14-40	24.00±1.25	24-260	146.60±11.20	-1.14	2.24	0.97	0.74
<i>Citharinus citharus</i>	Downstream	M	15	22.3-37	27.20±0.40	102-609	242.00±18.50	-3.39	3.89	0.93	0.89
		F	22	27.5-38	29.40±0.60	115-600	244.50±26.90	-2.38	3.27	0.91	1.02
		M	15	25-40.0	25.80±0.42	118-480	278.60±16.20	-1.14	2.48	0.93	1.27
<i>Lates niloticus</i>	Upstream	F	23	24-37.5	28.90±1.20	118-600	320.80±36.90	-1.67	2.84	0.97	1.29
		M	17	15-42	26.50±1.22	30-740	270.20±44.50	-1.74	2.79	0.99	0.95
		F	20	15-43	26.90±1.20	30-760	276.70±66.50	-1.67	2.76	0.99	0.99
<i>Lates niloticus</i>	Downstream	M	12	16-40	23.50±1.25	28-542	199.20±28.40	-2.31	3.12	0.99	0.72
		F	24	16-39	21.90±1.00	28-500	115.20±38.10	-2.02	2.93	0.99	0.76

1).

The condition factor (*K*) of species in this study ranged between 0.47-1.48. The difference in condition factor value of fish specimens collected from up- and downstream sections may be attributed to variation in living conditions such as feeding, habitat quality, climatic condition and fish density. Thus, condition factor of the fish is strongly influenced by biotic and abiotic conditions (Egbal et al. 2011). Thus, condition factor is a veritable tool for assessing the health status of the aquatic ecosystem (Abowei 2009; Ighwela et al. 2011).

Ninety-nine percent of the studied species have growth exponent *b* value ranging from 2-4 limit value. Specifically, the values of *b* varied from 2.48 in *C. citharus* to 3.89 in *O. niloticus* at the downstream section of the Jebba Lake. In the upstream, the values of *b* ranged from 2.75-3.89 with *C. nigrodigitatus* and *C. citharus* having the lowest and highest *b* values, respectively (Table 1). On our

estimates on the LWRs of all fish species from Jebba Lake, the observed growth pattern in *A. biscutatus* was similar to that of *A. occidentalis* from River Rima, North-western Nigeria (Shinkafi & Ipinjolu 2010), Akata lake, Benue State, Nigeria (Ikongbeh et al. 2013) and Southeastern Caspian Sea basin (Ghojoghi et al. 2017). The LWR of *O. niloticus* indicates growth in favour of body weight. This observation was similar to growth pattern in new Calabar River, Gbedikere Lake and Anambra River, Nigeria (Ezenwaji, 2004; Adeyemi et al. 2009; Olopade et al. 2018). In contrast, the allometric growth pattern in *C. nigrodigitatus* from Jebba Lake differed markedly from previously observed growth pattern in *C. nigrodigitatus* (Fagade & Adebisi 1979; Ikongbeh et al. 2015). Thus, this difference may imply that *C. nigrodigitatus* consists of heterogeneous species complex with variations in body weight and total length.

The growth patterns in *C. citharus* and

A. baremoze observed in the downstream of Jebba Lake correlated with the previous report on *C. citharus* and *A. baremoze* from Anambra River basin, Nigeria (Ezenwaji & Ezenwaji 2009). This may be attributed to shared environmental conditions and adaptive responses to the depth and size of the river system (Ezenwaji & Ezenwaji 2009). In comparison, the regression values of 3.20 and 2.30 observed in *H. vittatus* from up- and downstream sections of the dam, respectively, were higher than the value reported in *H. forskalii* in River Benue, Nigeria (Ogbe et al. 2008) and the variance may be attributed to environmental conditions in the region (King 1994).

In conclusion, our study provides basic information on the LWRs and condition factor (*K*) of seven commercially important freshwater fish species from the up- and downstream sections of Jebba Lake. Also, the results from this study contributes to the knowledge of fish populations in Jebba Lake. In addition, we recommend further study that would involve broader sampling across different life cycles of all fish species found across the lake, to examine the growth patterns across different life stages and seasons of the year. This will assist fish scientists and managers for future studies and conservation plans, particularly for highly exploited fish populations.

Acknowledgments

The authors thank the interpreter, R. Adeshike Aderinto and Jebba artisanal fishermen for their help in fish collection.

References

Abowei, F.N. & Hart, A.I. 2009. Some morphometric parameters of 10 finfish species from the lower Nun River, Niger Delta, Nigeria. *Russian Journal of Biological Science* 4(3): 282-288.

Abu, O.M.G. & Agarin, O.J. 2016. Length-Weight Relationship and Condition Factor of Silver Catfish (*Chrysichthys nigrodigitatus*) from the Lower Reaches of the New Calabar River Niger Delta.

International Journal of Innovative Studies in Aquatic Biology and Fisheries 2(4): 1-7.

Adelakun, K.M.; Mustapha, M.K.; Amali, R.P. & Mohammed, N. 2017. Seasonal Variation in Nutritional Quality of Catfish (*Clarias gariepinus*) from Upper Jebba Basin, Nigeria. *Journal of Nutrition & Food Science* 7: 5-9.

Adeyemi, S.O.; Akombu, P.M. & Toluhi, O.O. 2009. Diet and dietary habits Of *Labeo Senegalensis* in a tropical freshwater ecosystem continental. *Journal of Animal and Veterinary Research* 1: 25-30.

Araoye, P.A. 1999. Spatio-temporal distribution of the fish *Synodontis schall* (Teleostei: Mochokidae) in Asa lake, Ilorin, Nigeria. *Revista de Biología Tropical* 47(4): 1061-1066.

Atama, C.I.; Okeke, O.C.; Ekeh, F.N.; Ezenwaji, N.E.; Onah, I.E. Ivoke, N.; Onoja, U.S. & Eyo, J.E. 2013. Length-Weight Relationship and Condition Factor of Six Cichlid (Cichlidae: Perciformis) Species of Anambra River, Nigeria. *Journal of Fisheries and Aquaculture* 4(2): 82-86.

Ayoade, A.A. & Ikulala, A.O.O. 2007. Length weight relationship, condition factor and stomach contents of *Hemichromis bimaculatus*, *Sarotherodon melanotheron* and *Chromidotilapia guentheri* (Perciformes: Cichlidae) in Eleiyele Lake, Southwestern Nigeria. *Revista de Biología Tropical* 55(3-4): 969-977.

Bagenal, T.B. 1978. Aspects of fish fecundity. In: Gerking, S.D. (ed), *Ecology Of Fish Production*. Blackwell Scientific Publisher. Blackwell, Oxford. 519 p.

Egbal, O.A.; Mohammed, E.A. & Afra, A.A. 2011. Length-weight relationship and condition factor of six fish species in Atbara River and Khashim El-Girba Reservoir, Sudan. *International Journal of Agricultural Science* 3(1): 65-70.

Ecoutin, J.M. Albaret, J.J. & Trape, S. 2005. Length-weight relationships for fish populations of a relatively undisturbed tropical estuary: The Gambia Fish Research 72: 347-351.

Ezenwaji, H.M.G. 2004. Length-weight relationships of fishes from Anambra River south-western, Nigeria. *Animal Research International* 1(1): 1-6.

Ezenwaji, N.E. & Ezenwaji, H.M.G. 2009. Length-weight relationships and condition factor of

- Citharinus citharus* and *Alestes baremoze* from Anambra River basin, Nigeria. *Animal Research International* 6(3): 1107-1109.
- Fagade, S.O. & Adebisi, A.A. 1979. On the fecundity of *Chrysichthys nigrodigitatus* (Lacepede) of Asejire Dam, Oyo State, Nigeria. *Nigeria Journal of Natural Science* 1: 127-131.
- Ghojoghi, A.; Patimar, R.; Hojat Jafaryan, H. & Golzarianpour, K. 2017. Age and growth of *Alburnoides* cf. *tabarestanensis* (Teleostei: Cyprinidae) in the Zav Stream, Southeastern Caspian Sea basin. *Iranian Journal of Ichthyology* 4(4): 331-339.
- Hammer, O. 2011. PAST- Paleontological Statistics. Version 2.12 Reference manual. Natural History Museum, University of Oslo, Norway, Accessed at www.nhm.uio.no in March, 2018. 220 p.
- Idodo-Umeh, G. 2003. Freshwater fishes of Nigeria (taxonomy, ecological notes, diet and utilization). Idodo-Umeh Publisher Ltd. Benin-City, Nigeria. 232 p.
- Ighwela, A.; Ahmed, B. & Abol-Munafi, B. 2011. Condition factor as an indicator of growth and feeding intensity of Nile Tilapia fingerlings (*Oreochromis niloticus*) fed on different levels of Maltose. *American-Eurasian Journal of Agricultural and Environmental Science* 11(4): 559-563.
- Ikongbeh, O.A.; Ogbe, F.G. & Solomon, S.G. 2013. Length-Weight Relationship and Condition Factor of *Auchenoglanis occidentalis* (Valenciennes, 1775) from Lake Akata, Benue State, Nigeria. *IOSR Journal of Agriculture and Veterinary Science* 3(6): 11-17.
- Ikongbeh, O.A.; Ogbe, F.G.; Solomon, S.G. & Ataguba, G.A. 2015. Age, growth and mortality of *Chrysichthys nigrodigitatus* (Lacépède, 1803) from Lake Akata, Benue State, Nigeria. *Asian Journal of Conservation Biology* 4: 81-88.
- Irom, O.; Offem, B.O. & Keremah, R.I. 2017. Length-weight relationship, condition factor and gut content of *Chrysichthys furcatus* Gunther, 1864 (Bagridae) from Cross River at Ahaha. *Fisheries and Aquaculture Journal* 8: 2150-228.
- King, R.P. 1994. The biology of *Tilapia mariae* in a small tropical stream. Phd Thesis, University of Port Harcourt, Nigeria.
- Obot, E.A. 1989. The macrophytic flora of the draw-down area of Lake Kainji, Nigeria. *African Journal of Ecology* 27: 173-177.
- Ogbe, F.G. & Ataguba, G.A., 2008. Studies on the feeding habits and growth patterns and reproductive biology of *Malaptererus electricus*, (Gmelin, 1789) in Lower Benue River, Nigeria. *Biological Environment Science Journal Tropical* 5: 169-176.
- Olaosebikan, B.D. & Raji, A. 1998. Field Guide to Nigeria Freshwater Fishes. Federal College of Freshwater Fisheries Technology Publication. Decency printers, Ilorin, Nigeria. 110p.
- Olopade, O.A.; Dienye, H.E. & Eyekpegba, A. 2018. Length frequency distribution, length-weight relationship and condition factor of cichlid fishes (Teleostei: Cichlidae) from the New Calabar River, Nigeria. *Iranian Journal of Ichthyology* 5(1): 74-80.
- Shinkafi, B.A. & Ipinjolu, J.K. 2010. Morphometric relationships and relative condition factor of *Auchenoglanis occidentalis* (Cuvier and Valenciennes) from River Rima, North-western Nigeria. *Journal of Fishery International* 5(4): 61-66.
- Sossoukpe, E.; Nunoo, F.K.E.; Ofori-Danson, P.K.; Fiogbe, E.D. & Dankwa, H.R. 2013. Growth and mortality parameters of *P. senegalensis* and *P. typus* (Sciaenids) in the near shore waters of Benin and their implications for management and conservation. *Fishery Research* 137: 71-80.

یافته علمی کوتاه

الگوی رشد و ضریب وضعیت هفت گونه ماهی آب شیرین دریاچه جبا، شمال مرکزی نیجریه

سگون او. اولادیو^{۱,۲*}، لوتانا ام. نجی^۳، عباس توبا آنیفووش^۲، ایفینی سی. نجی^۴، اولوینکا ا. ایولا^۲، آدنی سی. آدیولا^۲، موشود کی. مصطفی^۲

^۱گروه علوم زیستی و زیست فناوری، دانشگاه ایالتی کوارا، مالت، ایالت کوارا، نیجریه.

^۲گروه جانورشناسی، دانشکده علوم زیستی، دانشگاه ایلورین، ایلورین، ایالت کوارا، نیجریه.

^۳انستیتو جانورشناسی کانمینگ، آکادمی علوم چین، کانمینگ ۶۵۰۲۲۳، چین.

^۴گروه علوم زیستی، دانشگاه آبوجا، آبوجا، ایالت، نیجریه.

چکیده: رابطه طول و وزن و ضریب وضعیت هفت گونه ماهی آب شیرین از دریاچه جبا (نیجریه) مورد بررسی قرار گرفت. جمعاً ۵۷۶ نمونه با قلاب، تور پرتابی و گوشگیر از بالادست و پایین دست دریاچه صید گردید. نمونه برداری در دو فصل خشک و بارانی از فروردین تا شهریور ۱۳۹۶ صورت گرفت. دامنه تغییرات توان رشد b بین ۲/۲۳-۳/۸۹ و ضریب وضعیت بین ۰/۴۷-۱/۶۱ بود. کلمات کلیدی: رابطه طول و وزن، *Oreochromis Hydrocynus*, *Chrysichthys Barbus*.