

Molecular diagnosis of the tick-borne pathogen *Anaplasma marginale* in cattle blood samples from Nigeria using qPCR

Nusirat Elelu¹ · Joana Ferrolho²  · Joana Couto² · Ana Domingos² · Mark C. Eisler¹

Received: 9 June 2016 / Accepted: 3 August 2016
© Springer International Publishing Switzerland 2016

Abstract Tick-borne diseases (TBDs) are some of the most important animal health and management problems in Africa, including Nigeria. This study aims to determine the prevalence of an important TBD, anaplasmosis, in a North-central region of Nigeria. Blood samples were collected from cattle and stored on Whatman FTA[®] cards. Information on village, age and sex associated with each cattle was also recorded. The packed red blood cell volume (PCV) for each blood sample was determined. After DNA extraction, pathogen presence was evaluated by TaqMan[®] based qPCR of which 75.9 % of the cattle tested positive for *Anaplasma marginale*. Statistical analysis revealed that the presence of *A. marginale* infection differed significantly between cattle age groups. However, there was no significant difference in the prevalence of this pathogen between the sexes or among cattle grouped by PCV level. Finally, using a highly sensitive molecular method our pioneer study contributes to the improvement of the current knowledge regarding tick-borne pathogens that seriously affect animal health in specific areas of Nigeria.

Keywords *Anaplasma marginale* · Tick-borne diseases · Tick-borne pathogens · qPCR · Kwara · Nigeria

✉ Joana Ferrolho
joana.ferrolho@ihmt.unl.pt

Nusirat Elelu
nusirat.elelu@bristol.ac.uk

Joana Couto
jmgtcouto@gmail.com

Ana Domingos
adomingos@ihmt.unl.pt

Mark C. Eisler
mark.eisler@bristol.ac.uk

¹ School of Veterinary Science, University of Bristol, Langford, Bristol BS40 5DU, UK

² Global Health and Tropical Medicine, Instituto de Higiene e Medicina Tropical, Rua da Junqueira 100, 1349-008 Lisbon, Portugal

Abbreviations

DNA	Deoxyribonucleic acid
EDTA	Ethylenediaminetetraacetic acid
LGA	Local Government Area
PCR	Polymerase chain reaction
PCV	Packed cell volume
qPCR	Real time PCR
rRNA	Ribosomal ribonucleic acid
TBDs	Tick-borne diseases

Introduction

The interest in tick-borne diseases (TBDs) has emerged during the past decades, especially from the veterinary point of view (Jongejan and Uilenberg 2004). It is well known that the prevalence of these diseases around the world is gradually increasing, demanding new treatment and preventive control measures (Dantas-Torres et al. 2012). In Africa, TBDs are one of the most important animal health and management problems, seriously affecting the development and productivity of the livestock industry (Young et al. 1988). It has been estimated that the annual cost of TBDs to the development and productivity of this industry is between US\$ 13.9–18.7 billion based on a worldwide cattle population of 1288 million (de Castro 1997). Other sources of financial losses that are more difficult to estimate also associated with TBDs include the downgrading of hides due to tick bites, secondary infections due to dermatophilosis and reduced fertility and abortion (Kivaria 2006). In addition, some TBDs are also zoonotic and fatal to man (Mehlhorn and Schein 1993). An important TBDs that pose a significant threat to cattle is bovine anaplasmosis whose transmission has been associated with ticks of the genus *Rhipicephalus* subgenus *Boophilus* (Aiello 1998; Bock et al. 2004). This disease represent a major constraint to livestock productivity in the tropical and sub-tropical regions of the world (Jongejan and Uilenberg 2004) and have been widely reported across Africa (Bell-Sakyi et al. 2004; Okuthe and Buyu 2006; Gachohi et al. 2010; Simuunza et al. 2011), including in Nigeria (Kamani et al. 2010; Reye et al. 2012). Bovine anaplasmosis is a tick-transmitted intraerythrocytic, rickettsial disease caused by *A. marginale* or *A. centrale* (Young et al. 1988). The disease is characterised by progressive anaemia due to destruction of erythrocytes by phagocytosis, resulting in icterus without haemoglobinemia and haemoglobinuria (Kocan et al. 2000). To the best of our knowledge, in Nigeria only a very few studies for the detection of tick-borne pathogens in cattle has used molecular methods and even less applied qPCR. The main objective of this study was to provide up-dated information using qPCR regarding the presence of *A. marginale* pathogens in blood samples collected from cattle of 11 villages of the Edu Local Government Area (LGA), in Kwara State, North-central Nigeria. A statistical analysis was also conducted to determine the effect of infection on cattle age, sex, levels of anaemia using packed cell volume (PCV). Our study constitutes the first report of the detection by qPCR of *A. marginale* in cattle blood samples in Edu LGA, Nigeria. Through a highly sensitive molecular method, such as the qPCR TaqMan® based assay we expect to contribute for current knowledge about the circulating tick-borne pathogens that seriously affect animal health in some areas of Nigeria and for a better planning of effective tick control measures.

Materials and methods

Study population

Edu LGA was selected as the study location because it has very large pastoralist settlements and is one of the largest cattle producing area in Kwara State, North-central Nigeria. A local informant identified 11 cattle producing villages (Bacita, Belle, Bokungi, Fanagun, Fedudangi, Gonandogo, Mokwagi, Ndabata, Ndachewoye, Tshonga and Yelwa) in Edu LGA (Fig. 1) and these formed the study population. Sixty-five households were visited in the 11 villages and between 3 and 4 cattle were sampled in each household. A total of 253 bovines were selected to collect individual blood samples between April and August 2013. During collection, the age and sex of each sampled animal was recorded.

Haematology

Five millilitres (ml) of blood were collected from the jugular vein of each animal with a syringe, transferred into EDTA specimen container and kept on ice during transport to the laboratory facilities. From each sample, approximately 100 µl was applied into Whatman® FTA® cards (Whatman, UK). The cards were air dried and stored at room temperature, according to the manufacturer's instructions, for downstream applications. From the remainder of the collected blood, the packed cell volume (PCV) was also determined using the micro haematocrit method (McInroy 1954).

DNA extraction

Extraction of DNA from blood stored on Whatman FTA cards was carried out following the method recommended for diagnosis of blood-borne infections (Ahmed et al. 2011), with some modifications. Briefly, a 3 mm diameter disc was cut from each blood sample stored on the Whatman FTA cards using a hand paper punch. To avoid cross contamination, five discs were punched from blank filter paper between each sample and included as negative controls. The excised discs were first washed twice for 15 min with Whatman FTA Purification Reagent (Whatman, UK) in order to remove haemoglobin, and then washed twice for 15 min with 100 µl of Tris/EDTA buffer to remove the purification reagent. The discs were then dried for 30 min at 37 °C. To elute the DNA, 60 µl of 5 % aqueous suspension of Chelex Resin (Bio-Rad, UK) was added to each disc and incubated at 90 °C for 30 min, as previously described (Becker et al. 2004). The eluted DNA was stored at −20 °C until used for the analysis.

Real time PCR

Real time PCR was conducted to determine the presence of *A. marginale* using primers and TaqMan fluorescence-labelled probes that specifically amplify a fragment of the *msp1b* gene of *A. marginale* (Carelli et al. 2007). PCR amplifications were carried out in a 7500 Fast Applied Biosystems thermocycler (Alfagene, Portugal) with the SensiFAST™ Probe Low-Rox kit (Bioline, Portugal). For the samples tested, a 20 µl reaction mix was prepared with 10 µl of SensiFAST Probe Low-Rox mix, 400 nM of reverse and forward primers, 100 nM of probe, 2 µl of DNA template and nuclease-free water up to the final volume. The qPCR machine was programmed for an initial denaturation stage at 95 °C for 5 min,

(a)




Fig. 1 Locations where the cattle blood samples were collected. **a** Kwara State showing the location of Edu Local Government Area, Nigeria. **b** Location of the 11 cattle producing villages in Edu Local Government Area of Kwara State. Maps were constructed using Arc Map software version 10.2.2

followed by a PCR stage of 45 cycles of 95 °C for 45 s and 60 °C for 1 min. Negative controls were prepared with no template and positive controls with *A. marginale* Jabotibabal strain. PCR efficiency ($90\% \leq \text{efficiency} \leq 115\%$) was determined generating




Fig. 1 continued

sequence-specific standard curves with fivefold serial dilutions of DNA from the positive controls. All reactions were loaded in triplicate into 96-well plates (Bioline, Portugal). The threshold was manually set at 0.05 for *A. marginale*. The data generated for each plate was analysed using the Applied Biosystems 7500 Fast 2.0.6 Software (Alfagene, Portugal). Samples with quantification cycle (C_q) values above 35 for *A. marginale* were considered negative for the presence of the pathogen.

Statistical analysis

At the time of sample collection, the age of and the sex of 253 animals were recorded. For the age classification, animals were considered weaners if they were between 1 and 2 years old and adults if they were older than 2 years old. Statistical analysis was conducted to determine the effect of *A. marginale* infection on cattle age, sex and PCV using Microsoft Excel (Microsoft Office Professional Plus 2013) and SPSS Statistical software (Version 23; IBM). Univariate analysis using χ^2 or Fisher's exact test (Plackett 1983) and Pearson's correlation (R) were carried out. The level of significance was set at 0.05. Animals with missing PCV values were excluded in the univariate analysis.

Results

Between April and August 2013, a total of 253 blood samples were collected from bovines of 11 cattle producing villages in Edu LGA that included Bacita, Belle, Bokungi, Fanagun, Fedudangi, Gonandogo, Mokwagi, Ndabata, Ndachewoye, Tshonga and Yelwa. Most of the samples were collected at Ndachewoye 64/253 (25.3 %), Fanagun 56/253 (22.1 %) and Gonandogo 40/253 (15.8 %).

Results of the descriptive statistics have shown that 11.1 % (28/253) were weaners, whereas 88.9 % (225/253) were adult animals. Regarding the sex of the animals, 45.5 % (115/253) were males and 54.5 % (138/253) were females.




Fig. 2 Villages of Edu LGA in Kwara State, North-central Nigeria where the blood samples were collected between April and August 2013. *Black* Number of samples collected from each village, with a total of 253 samples; *Grey* Number of positive samples for *Anaplasma marginale*

To determine the presence of *A. marginale* in the surveyed animals by qPCR, DNA was extracted from 253 blood samples stored on Whatman FTA cards. The specific primers and TaqMan fluorescence-labelled probes that amplify a fragment of the *msp1b* gene of *A. marginale* (Carelli et al. 2007) assured the specificity of the reaction. All the reaction plates were validated by the absence of amplification in the negative controls and by amplification with the positive controls of *A. marginale* Jaboticabal strain. The 5-serial dilution standard curves that were generated allowed to determine the efficiency of each reaction plate automatically with the Applied Biosystems 7500 Fast 2.0.6 Software (Stolovitzky and Cecchi 1996). The efficiencies of the plates were within the range 90–115 %, also validating the reaction. Regarding the detection of pathogens, for *A. marginale*, a total of 192 of the 253 blood samples were considered positive, which corresponds to an overall infection rate of 75.9 %. The village Belle presented 20/20 (100 %) of the samples infected with *A. marginale*, closely followed by Ndachewoye 59/64 (92.2 %), Fedudangi 10/11 (90.9 %), Bacita 6/7 (85.7 %) and Bokungi 10/12 (83.3 %). In the villages Gonandogo, Tshonga and Yelwa *A. marginale* was detected in 75 % of the collected samples. With a lower infection rate were the villages Ndabata, Mokwagi and Fanagun with 57.1, 50 and 48.2 %, respectively. The number of positive samples found in each village sampled is described in Fig. 2.

When considering the sex classification and the presence of infection, it was observed that the percentage of infected males and females was approximately the same. Eighty-six of the 115 males were positive (74.8 %), whilst 104 of the 138 females were found positive (75.4 %), the difference not being statistically significant ($p > 0.9$). Looking at the cattle age groups and the detection of infection, the prevalence of *A. marginale* was significantly higher ($p < 0.05$) in adult cattle 174/225 (78.4 %) when compared to weaners 16/28 (57.1 %).

The PCV of 233 animals was determined using the micro haematocrit method and grouped into five classes (10–20, 21–30, 31–40; 41–50 and 51–60 %). The results of the number of positive samples *per* PCV class are shown in Fig. 3. Twenty of the 253 cattle had missing PCV results and were excluded from the haematological analysis. It was




Fig. 3 Packed cell volume (PCV, %) determined using the micro haematocrit method. *Black* Number of animals according to the % of PCV; *Grey* Number of positive samples for *Anaplasma marginale*

observed that 66.7, 78.8, 78.2, 73.9, and 50 % of the animals that presented a PCV between 10–20, 21–30, 31–40, 41–50, and 51–60 % respectively were infected with *A. marginale*, the differences not being significant ($p > 0.8$). A weak negative correlation was observed between the presence of *A. marginale* infection and the values of PCV ($R = -0.026$).

Discussion

Having accurate disease diagnosis within animal populations is a necessary precursor for designing appropriate control strategies. To date, the prevalence of tick-borne pathogens in cattle from Nigeria has been assessed by parasitological examination of Giemsa stained blood smears (Kamani et al. 2010), and in ticks collected from cattle and dogs by molecular methods, such as conventional PCR (Kamani et al. 2011; Ogo et al. 2012). Although conventional PCR has been shown to be useful to determine the presence of *Babesia* and *Anaplasma* (Figuerola et al. 1993; Martins et al. 2008; Bilgiç et al. 2013; El-Ashker et al. 2015), more recently, qPCR has revealed to be more sensitive than the first for the detection of protozoan parasites in cattle (Criado-Fornelio et al. 2009). For the detection of anaplasmosis, some of the TaqMan based qPCR methods involve the amplification of species-specific genes such as the major surface proteins (*msp*) genes (Carelli et al. 2007; Bacanelli et al. 2014), and six *msp* (*msp* 1a, 1b, 2, 3, 4, 5) with the potential to be used in diagnostic assays have been identified on *A. marginale* (Kocan et al. 2003). The *msp1* gene complex (composed of *msp1a* and *msp1b*) has been shown to be a well-conserved gene in *A. marginale* (de la Fuente et al. 2001; Bowie et al. 2002), whose products act as adhesin during invasion of bovine erythrocytes (McGarey et al. 1994). In the present study, the TaqMan qPCR used were highly specific, and their sensitivity has been previously evaluated for *A. marginale* (Carelli et al. 2007). Probe-based qPCR methods are more expensive but, when compared to other PCR methods have multiple advantages, including the detection of very low infections, no requirement of post PCR steps, and they are a useful quantitative technique for chemotherapeutic and vaccine

evaluation (Monis et al. 2005; Bacanelli et al. 2014). They are also useful in processing several samples simultaneously for large scale epidemiological studies (Carelli et al. 2007). Our results have shown an overall prevalence of 75.9 % for *A. marginale*, being greater than 40 % in all the villages studied. This result suggests that the disease in the region of Kwara State is endemic and may have become endemically stable for *A. marginale*. This endemic stability for anaplasmosis was similarly reported in a previous study in Brazil (Bacanelli et al. 2014). Endemic stability is an epidemiological state, in which there is high haemoparasites-infected ticks but clinical disease is absent in the population (Jonsson et al. 2012). *Anaplasma marginale* was previously reported as the most prevalent (53.5 %) haemoparasite in ticks removed from Nigerian cattle (Reye et al. 2012) and the tick vector *Rhipicephalus* subgenus *Boophilus* is also the most prevalent in cattle in Nigeria (Lorusso et al. 2013). Furthermore, the high prevalence of anaplasmosis reported in this study may also suggest the detection of carrier or chronically infected animals with low parasitaemia. This is important in the epidemiology of anaplasmosis because these previously infected cattle can now serve as an infective reservoir for the disease. The principal source for infection with *Anaplasma* is a persistent carrier state that normally occurs after the animal recovers from the acute phase of the disease (Aiello 1998; Kocan et al. 2000; Bock et al. 2004). Acting as reservoirs for tick transmission, these animals contribute to the propagation of anaplasmosis to susceptible hosts, thereby increasing the number of newly infected animals. Moreover, carrier animals during a stressful condition or poor nutrition are at risk of succumbing with clinical disease.

The lack of significant difference in prevalence of anaplasmosis between sexes of cattle in this study is supported by similar findings from a past study in Sudan on haemoparasitic infection (Awad et al. 2011). In this study, adult cattle had a significantly higher prevalence ($p = 0.013$) of infection (78.4 %) compared to weaners (57.1 %). This finding is in agreement with past studies in Nigeria where this difference in prevalence between age groups (Kamani et al. 2010). A negative correlation might be expected between PCV and *A. marginale* infection because anaplasmosis is a haemoparasitic disease characterised by progressive anaemia due to destruction of erythrocytes by phagocytosis. The weak negative correlation observed here was consistent with this expectation but not statistically significant. A reduction in PCV in Nigerian livestock has been shown in infections with tick-borne parasites characterised by anaemia, including in *A. marginale* infection (Obi and Anosa 1980). This observation is in agreement with another study in Nigeria that reported a reduction in PCV due to haemoparasitic infections (Kamani et al. 2010). In a similar study carried out in Ghana, the PCV values in cattle studied were significantly higher in non-infected animals compared to those with haemoparasitic infection (Bell-Sakyi et al. 2004).

Conclusion

Ticks and TBDs are considered an emerging global threat both for humans and animals. The results of this study indicate that in the 11 villages of Edu LGA, in Nigeria, where the blood samples were collected *A. marginale* was detected with high prevalence rates. Also, it was confirmed that the qPCR TaqMan based assay is indeed sensitive and effective for the detection of tick transmitted haemoparasites, revealing more positive samples than previous studies carried out in Nigeria. The confirmation of the presence of *Anaplasma* in Nigeria is of extreme importance as this pathogen has great impact in animal health, influencing animal production. Finally, information about the prevalence of infection in

cattle is essential to provide information in order to develop future effective preventive and control strategies.

Acknowledgments The authors would like to acknowledge Rosângela Zacarias Machado (Universidade Estadual Paulista—Faculdade de Ciências Agrárias e Veterinárias, Brazil) for the *Anaplasma marginale* control. We would also like to thank the staff of the Department of Veterinary Services, Kwara State Ministry of Agriculture for providing field support and all the farmers that participated in the research.

Compliance with ethical standards

Conflict of interest The authors declare no competing personal or financial interests.

References

- Ahmed H, MacLeod E, Hide G, Welburn S, Picozzi K (2011) The best practice for preparation of samples from FTA(R)cards for diagnosis of blood borne infections using African trypanosomes as a model system. *Parasit Vectors* 4:68
- Aiello SE (1998) The Merck veterinary manual. Wiley, Hoboken
- Awad H, Antunes S, Galindo RC, do Rosário VE, de la Fuente J, Domingos A, El Hussein AM (2011) Prevalence and genetic diversity of *Babesia* and *Anaplasma* species in cattle in Sudan. *Vet Parasitol* 181:146–152
- Bacanelli GM, Ramos CAN, Araújo FR (2014) Molecular diagnosis of *Anaplasma marginale* in cattle: quantitative evaluation of a real-time PCR (Polymerase Chain Reaction) based on msp5 gene. *Pesqui Vet Bras* 34:29–33
- Becker S, Franco JR, Simarro PP, Stich A, Abel PM, Steverding D (2004) Real-time PCR for detection of *Trypanosoma brucei* in human blood samples. *Diagn Microbiol Infect Dis* 50:193–199
- Bell-Sakyi L, Koney EBM, Dogbey O, Walker AR (2004) Incidence and prevalence of tick-borne haemoparasites in domestic ruminants in Ghana. *Vet Parasitol* 124:25–42
- Bilgiç HB, Karagenc T, Simuunza M, Shiels B, Tait A, Eren H, Weir W (2013) Development of a multiplex PCR assay for simultaneous detection of *Theileria annulata*, *Babesia bovis* and *Anaplasma marginale* in cattle. *Exp Parasitol* 133:222–229
- Bock R, Jackson L, de Vos A, Jorgensen W (2004) Babesiosis of cattle. *Parasitology* 129:S247–S269
- Bowie MV, de la Fuente J, Kocan KM, Blouin EF, Barbet AF (2002) Conservation of major surface protein 1 genes of *Anaplasma marginale* during cyclic transmission between ticks and cattle. *Gene* 282:95–102
- Carelli G, Decaro N, Lorusso A, Elia G, Lorusso E, Mari V, Ceci L, Buonavoglia C (2007) Detection and quantification of *Anaplasma marginale* DNA in blood samples of cattle by real-time PCR. *Vet Microbiol* 124:107–114
- Criado-Fornelio A, Buling A, Asenzo G, Benitez D, Florin-Christensen M, Gonzalez-Oliva A, Henriques G, Silva M, Alongi A, Agnone A, Torina A, Madruga CR (2009) Development of fluorogenic probe-based PCR assays for the detection and quantification of bovine piroplasmids. *Vet Parasitol* 162:200–206
- Dantas-Torres F, Chomel BB, Otranto D (2012) Ticks and tick-borne diseases: a One Health perspective. *Trends Parasitol* 28:437–446
- de Castro JJ (1997) Sustainable tick and tickborne disease control in livestock improvement in developing countries. *Vet Parasitol* 71:77–97
- de la Fuente J, Garcia-Garcia JC, Blouin EF, Kocan KM (2001) Differential adhesion of major surface proteins 1a and 1b of the ehrlichial cattle pathogen *Anaplasma marginale* to bovine erythrocytes and tick cells. *Int J Parasitol* 31:145–153
- El-Ashker M, Hotzel H, Gwida M, El-Beskawy M, Silaghi C, Tomaso H (2015) Molecular biological identification of *Babesia*, *Theileria*, and *Anaplasma* species in cattle in Egypt using PCR assays, gene sequence analysis and a novel DNA microarray. *Vet Parasitol* 207:329–334
- Figuerola JV, Chievas LP, Johnson GS, Buening GM (1993) Multiplex polymerase chain reaction based assay for the detection of *Babesia bigemina*, *Babesia bovis* and *Anaplasma marginale* DNA in bovine blood. *Vet Parasitol* 50:69–81
- Gachohi JM, Ngumi PN, Kitale PM, Skilton RA (2010) Estimating seroprevalence and variation to four tick-borne infections and determination of associated risk factors in cattle under traditional mixed farming system in Mbeere District, Kenya. *Prev Vet Med* 95:208–223
- Jongejan F, Uilenberg G (2004) The global importance of ticks. *Parasitology* 129:S3–S14

- Jonsson NN, Bock RE, Jorgensen WK, Morton JM, Stear MJ (2012) Is endemic stability of tick-borne disease in cattle a useful concept? *Trends Parasitol* 28:85–89
- Kamani J, Sannusi A, Egwu OK, Dogo GI, Tanko TJ, Kemza S, Tafarki AE, Gbise DS (2010) Prevalence and significance of haemoparasitic infections of cattle in North-central, Nigeria. *Vet World* 3:445–448
- Kamani J, Jwander LD, Ubali Z (2011) Demonstration of vermicles of *Babesia* species in haemolymph smears of *Amblyomma variegatum* in Nigeria. *J Adv Vet Res* 1:1–3
- Kivaria FM (2006) Estimated direct economic costs associated with tick-borne diseases on cattle in Tanzania. *Trop Anim Health Prod* 38:291–299
- Kocan KM, Blouin EF, Barbet AF (2000) Anaplasmosis control: past, present, and future. *Ann N Y Acad Sci* 916:501–509
- Kocan KM, de la Fuente J, Guglielmone AA, Meléndez RD (2003) Antigens and alternatives for control of *Anaplasma marginale* infection in cattle. *Clin Microbiol Rev* 16:698–712
- Lorusso V, Picozzi K, de Bronsvort B, Majekodunmi A, Dongkum C, Balak G, Igweh A, Welburn S (2013) Ixodid ticks of traditionally managed cattle in central Nigeria: where *Rhipicephalus* (*Boophilus*) *microplus* does not dare (yet?). *Parasit Vectors* 6:171
- Martins TM, Pedro OC, Caldeira RA, do Rosário VE, Neves L, Domingos A (2008) Detection of bovine babesiosis in Mozambique by a novel seminested hot-start PCR method. *Vet Parasitol* 153:225–230
- McGarey DJ, Barbet AF, Palmer GH, McGuire TC, Allred DR (1994) Putative adhesins of *Anaplasma marginale*: major surface polypeptides 1a and 1b. *Infect Immun* 62:4594–4601
- McInroy RA (1954) A micro-haematocrit for determining the packed cell volume and haemoglobin concentration on capillary blood. *J Clin Pathol* 7:32–36
- Mehlhorn H, Schein E (1993) The piroplasms: “A long story in short” or “Robert Koch has seen it”. *Eur J Protistol* 29:279–293
- Monis PT, Giglio S, Keegan AR, Andrew Thompson RC (2005) Emerging technologies for the detection and genetic characterization of protozoan parasites. *Trends Parasitol* 21:340–346
- Obi TU, Anosa VO (1980) Haematological studies on domestic animals in Nigeria. *Zentralblatt für Veterinärmedizin Reihe B* 27:789–797
- Ogo NI, de Mera IG, Galindo RC, Okubanjo OO, Inuwa HM, Agbede RI, Torina A, Alongi A, Vicente J, Gortazar C, de la Fuente J (2012) Molecular identification of tick-borne pathogens in Nigerian ticks. *Vet Parasitol* 187:572–577
- Okuthe OS, Buyu GE (2006) Prevalence and incidence of tick-borne diseases in smallholder farming systems in the western-Kenya highlands. *Vet Parasitol* 141:307–312
- Plackett RL (1983) Karl Pearson and the Chi squared test. *Int Stat Rev Revue Internationale de Statistique* 51:59–72
- Reye AL, Arinola OG, Hübschen JM, Muller CP (2012) Pathogen prevalence in ticks collected from the vegetation and livestock in Nigeria. *Appl Environ Microbiol* 78:2562–2568
- Simuunza M, Weir W, Courcier E, Tait A, Shiels B (2011) Epidemiological analysis of tick-borne diseases in Zambia. *Vet Parasitol* 175:331–342
- Stolovitzky G, Cecchi G (1996) Efficiency of DNA replication in the polymerase chain reaction. *Proc Natl Acad Sci* 93:12947–12952
- Young AS, Grocock CM, Kariuki DP (1988) Integrated control of ticks and tick-borne diseases of cattle in Africa. *Parasitology* 96:403–432