




[Submit Your Article](#)

Volume 15, Number 2, Year 2024

[Home](#) [Issue](#)



Do You Want To Upload Your Article?
Submit Now & Make Your Online Presence

[Submit Now](#)



 Mary Oluwatobiloba Adedigba , Adeshina Oloruntoba Adekeye , Ephraim Samuel Jen , Kehinde S. Olaniyi , Oluwatosin Olalekan Ogedengbe

[Lycopersicon esculentum Attenuates Oxidative Stress and Hormonal Dysregulation in Letrozole-induced Polycystic Ovarian Syndrome Rat Model](#)

Polycystic ovarian syndrome (PCOS) is the primary culprit behind anovulatory infertility and is one of the most prevalent endocrine disorders affecting women of childbearing age. The aim of the study was to explore the impact of Lycopersicon esculentum isolate on PCOS induced by letrozole in fem...

[View Abstract](#)

 Abdulrahim Halimat Amin, Fatoki John Olabode , Oyewo Emmanuel Bukoye , Nafiu Abdulrazaq Bidemi , Omeiza Noah Adavize, Adekunle Adeniran Sanmi

[Dyslipidemic Prospective of Levonorgestrel in the Selected Central Compartments of the Liver, Kidney and Heart](#)

Levonorgestrel, a dedicated emergency contraceptive, is prone to abuse by women of reproductive age group. We evaluated the modulatory effects of varying and repeated doses of the drug on lipid profiles in female rats. Twenty adult female Wistar rats were randomized into four groups. The normal ...

[View Abstract](#)

 Bolaji Israel Jayeoba , Musa Habib Modibbo , Ahmad Ahmad Gaya , Adam Jibril Nma

[Assessment of Celiac Trunk and Hepatic Artery Dimensions among Hausa Population in Northern Nigeria](#)

Anatomical variations in the celiac trunk and hepatic artery are critical for clinical practice, particularly in regions with unique population characteristics. This study investigated the relationship between celiac trunk, hepatic artery dimensions and their sex differences among Hausa subjects...

[View Abstract](#)

 Sarah I. David , Barnabas Danborno , Zainab M. Bauchi

[Anticancer Effect of Aqueous and Ethanol Extracts of Pepper Fruit \(Dennettia tripetala\) on the Morphology of MCF-7 Breast Cancer Cell Line](#)

[View Abstract](#)

Cancer is a multifactorial disease caused by combined genetic and non-genetic changes. Breast cancer being the second leading cause of death in women. A large percentage of known anticancer medicines are derived from various plant species. Dennettia tripetala has been found to contain vitamins, ...

 Kudighe P. Udoh , Akpan U. Ekanem , Utomobong U. Akpan , Eno-Obong I. Bassey , Ekaette P. Akpan

Costus afer Stem Juice Extract Enhances the Expression of HAS3 during the Wound Healing Process

Among its various applications, the stem juice of Costus afer has gained attention in the scientific community for its potential in tissue repair and regeneration. This study was carried out to assess the expression of hyaluronic acid synthase 3 (HAS3) in wounds treated with C. afer stem juice e...

[View Abstract](#)

 Ehi-Omosun Mabel Bilu and Samuel Victor

Effects of Aqueous Leaf Extract of Psidium guajava on Formaldehyde-induced Interstitial Pneumonitis in Adult Wistar Rats

The most common cause of lung diseases among people with formaldehyde-related occupations is formaldehyde poisoning which has significant morbidity and mortality if left untreated. This study was aimed at investigating the effects of aqueous leaf extract of Psidium guajava on formaldehyde-induced...

[View Abstract](#)

 Onyilo PO, Enaohwo TM, Omonigho EM

Neurobehavioural and Histomorphologic Study of Mice Hippocampus Following Exposure to Rohypnol

Rohypnol (Flunitrazepam), a highly lipophilic benzodiazepine derivative, is commonly used in treating seizures and sleep disorders. This study aimed to evaluate the histomorphological changes in the hippocampus and the associated neurobehavioral effects on spatial memory in mice following exposu...

[View Abstract](#)

 Ukpai C. J. , Eweoya O. , Abue A. , Okolo R. U , Onwih E.

Evaluating the Prevalence of Elongated Styloid Process in Dry Human Skulls

Styloid process is a bony projection from the inferior part of the petrous temporal bone of the skull. The research aimed to study the styloid process structure and evaluate the prevalence of elongated styloid processes in dry skulls collated by the Department of Anatomical Sciences University o...

[View Abstract](#)

 Yusha'u Z, Agbon AN, Danladi J

Morphological Differences in the Cerebrum of Adult and Juvenile Thryonomys swinderianus (African Grasscutter)

The African grasscutter (AGC; Thryonomys swinderianus) has been observed to possess unique physiological and structural features enabling them to survive in challenging environments. This behavior displayed by AGCs across different age groups could be attributed to a well-developed CNS where M1 ...

[View Abstract](#)

 Koko Ottoh Arikpo , Barnabas Danborno , James A. Timbuak , M. Tanko , Monday Nwankwo

Sexual Dimorphism in Anthropometric Characteristics of Children from Cross River State, Nigeria

Human sexual dimorphism is an outcome of a survival strategy, a balancing of the need for high degree of biological variation within species with the need for a narrow range of variation in the female, who is physically structured for the perinatal support of infant. This study aimed to evaluate...

[View Abstract](#)

 Abdulfattah Moisule Ganga , Maryam Isa Sharif , Lawan Hassan Adamu , Rabi Inuwa Fage

Relationship between Cephalic Anthropometry and Somatotype among Adolescents of Hausa Ethnic Group in Kano Metropolis, Nigeria

Anthropometric studies are very important areas for craniofacial surgery and syndromology. Surgeons use these measurements to understand the unique features of a patient's face, head and body, helping them tailor their surgical approaches for optimal results. It aids in achieving aesthetic and f...

[View Abstract](#)

👤 Elvis T Godam , Wilson Francisca Cheyachi , Mesole Samuel Bolaji , Musa Abraham Sunday

Neuroprotective Effects of Phoenix dactylifera L. and Melatonin on Aluminum Chloride-induced Neurotoxicity in the Cerebellum of Adult Male Wistar Rats

Exposure to aluminium is associated with neurotoxicity, affecting various parts of the nervous system. The study investigated the neuroprotective potential of ethanolic fruit extract of Phoenix dactylifera (EFPD) and melatonin against aluminium chloride (AlCl₃)-induced neurotoxicity in male adul...

[View Abstract](#)

👤 Adana Misturah Yetunde, Towaju Alaba Magrette, Alade Oluwadamilola Esther, Oluwasegun Blessing Olaitan, Adewale David Adebayo, Onigbolabi Oluwafemi Gideon, Ajao Moyosore Salihu

Comparative Analysis of the Protective Potentials of Nigella sativa and Lepidium meyenii on Alcohol-induced Testicular Toxicity in Adolescent Wistar Rats

Alcohol, a psychoactive drug, is soluble in both water and lipids, due to which it can diffuse to all tissues and affect the normal functioning of the body. Gonadal toxicity is reported as one of the side effects of its long-term consumption. This study examined the possible comparable protectiv...

[View Abstract](#)

👤 Enobong Obong and Olatunde Owoeye

Lasianthera africana Aqueous Extract Protects the Cerebral Cortex and CA1 Hippocampal Region against Ischemic-reperfusion Insult by Bilateral Common Carotid Artery Occlusion in Adult Male Wistar Rats

Animal models of cerebral ischaemia make a significant contribution to the knowledge of the pathogenesis of stroke and its methods of management. Many treatments have failed to reduce neuronal injury in cerebral ischaemia, prompting the exploration of new treatment models. The purpose of this st...

[View Abstract](#)

👤 Kehinde Oluwaseyi Adeniji, Olakayode Olaolu Ogundoyin

Morphometric and Histopathological Evaluation of the Testes and Accessory Sex Organs of Male Wistar Pups Following Perinatal Exposure to Chronic Cigarette Smoking

Exposure to chronic cigarette smoking during pregnancy into infancy results in numerous morphological and behavioral consequences in infants, children, and adolescents. Cigarette smoke is a reproductive toxicant, demonstrating a strong relationship between smoking and impaired fertility. This st...

[View Abstract](#)

👤 Happiness O. Inyang, Uchenna K. Ezemagu , Stephen O. Odey , Omamuyovwi M. Ijomone

Manganese Exposure Induces Mild Behavioral and Histological Deficits in Wistar Rats: Ameliorative Effects of D-ribose-L-cysteine

Heavy metal exposure is a global health problem and their presence in our environment has been greatly increased by industrial activities over the past century. Manganese is an essential trace metal in the body. However, exposure to elevated levels is neurotoxic, leading to manganism, a Parkinso...

[View Abstract](#)

👤 Nasidi A, Mustapha M, Sabo HL, Abdullahi I, Babankanwa AH, Ibrahim T, Ladan MB, Idris M, Bello SA, Ahmed Abdullahi UA, Shaaba JB

Association between Handedness, Digit Ratio and Personality Traits among Adult Hausa Ethnic Group in Jigawa State, North Western Nigeria

This study explored the relationship between handedness, 2D:4D ratio, and personality traits among the Hausa ethnic group in Jigawa State, Nigeria. Data from 500 participants aged 18-30 were analysed using a cross-sectional design and simple random sampling, confirming Hausa ethnicity and Nigeri...

[View Abstract](#)

👤 Kareem SB, Muyideen MK, Ajiboye AA, Imam A, Ibrahim A, Alabi AS, Ajao MS

Sex Determination Using Tali Bones

The talus is an important bone used in forensic and archaeological cases for the determination of the sex of human remains because it is a preservational favored bone of the body. The study is designed to take the measurements and dimensions of dry tali to determine the sex as well as the sides ...

[View Abstract](#)

👤 Anyanimeh J. Ukoh, Ejiro Ederademu, Affiong E. Ekanem, Mokutima A. Eluwa

Effects of Combined Aqueous Seed Extract of *Monodora myristica* and *Xylopia aethiopica* on the Histology of Pancreas of Wistar Rats

Monodora myristica and *Xylopia aethiopica* have been commonly used as spices for decades, both in culinary and medicinal contexts. This study aimed to assess the effect of aqueous extract of *Monodora myristica* and *Xylopia aethiopica* on the histology of the pancreas of adult albino Wistar rats. Tw...

[View Abstract](#)

👤 Aliyu Sabo Aliyu , Nana-Asmau Hassan , Awan Stephen Kasham , Ahmad Ahmad Gaya , Maryam Isa Sharif , Abdullahi Gudaji , Rilwanu Bello , Ali Bala Umar

Histological Effects of Aqueous *Allium sativum* Bulb Extract on Monosodium Glutamate-induced Hepatotoxicity in Guinea Pigs

Monosodium glutamate (MSG) is among the most common food additives in the developing world and is commonly used as a flavor enhancer. It contains glutamic acid, one of the naturally occurring amino acids that the body transforms into glutamate. *Allium sativum* (Garlic) has tremendous pharmacologic...

[View Abstract](#)

👤 Misturah Yetunde Adana, Edidiong Nnamso Akang, Offor Ugochukwu, Simeon Eche , Edwin Coleridge Stephen Naidu , Onyemaechi Okpara Azu

Naringenin Modulates Sertoli Cell Dysfunction and Altered SCF/c-kit Ligand System in the Setting of Combination Antiretroviral Therapy

Combination Antiretroviral Therapy (cART) has been shown in previous studies to induce histopathological changes in the testes. An essential function of the c-kit signalling system is to control cell survival, proliferation, differentiation, and death. Consequently, certain male infertility may ...

[View Abstract](#)

👤 Anthony Bol Kon Giet, Olatunji Sunday Yinka, Sylvia Santos

Case Report: A Large Cadaveric Simple Hepatic Cyst

Simple hepatic cysts are the most frequent forms of liver cysts, often filled with clear or yellow fluid. They are usually asymptomatic and are typically diagnosed in adulthood. Their prevalence however varies across populations. Ultrasound and serological tests are the primary diagnostic tools ...

[View Abstract](#)

👤 Kareem SB, Eweoya OO, Akinola O, Imam A, Ibrahim A, Arogundade TT, Adana MY, Alabi AS, Ajao M.S

Anthropometric Analysis of Craniofacial Indices in Students of Yoruba Lineage: Implications for Clinical Medicine, Forensic Anthropology, and Ethnic Diversity

Population-specific facial anthropometry is essential for cultural esteem, preservation of ethnicity, and advances in reconstructive medicine. This study focuses on craniofacial indices in students of Yoruba lineage. Using a questionnaire, biodata were collected from 2109 participants, aged 18 t...

[View Abstract](#)

👤 Ogunlade B

Therapeutic Response of Sulforaphane on Oxidative Stress and Cognitive Impairment in Aluminum-Induced Alzheimer-type Hippocampal Neurodegeneration in Adult Wistar Rats

The accumulation of Aluminum in the brain is believed to be involved in the pathophysiology of neurodegenerative diseases including Alzheimer's disease. This study evaluated the neuroprotective activities of sulforaphane (SFN) on aluminum chloride (AlCl₃)-induced hippocampal degeneration in Wist...

[View Abstract](#)

👤 Obioma, Ogbonnaya, Chikera, Samuel Ibe, Ekele, Ikpegbu

Comparative Postnatal Determination of Absolute Brain Weight and Size using Body Morphometric Parameters in the African Giant Rat (*Cricetomys gambianus*)

One neuroanatomical measure found to allometrically scale the body is brain's size. Species with larger brains relative to body are better complex information processors. We compared absolute brain weight (ABW), generated regression formulae and determined intelligence level in each age group of ...

[View Abstract](#)

👤 Felix E. Ehidiāmhēn , Stanley E. Ogbata , Godson O. Eze , Martins A. Nnoli , Olushola O. Jegede , Cornelius C. Chukwuegbo , Modupeola O. Samaila , Lateef A. Odukoya

Histologic Variants of Cutaneous Squamous Cell Carcinoma Diagnosed in a Tertiary Health Facility in Southeast Nigeri

Cutaneous squamous cell carcinomas are tumours originating from cutaneous keratinocytes and they are among the most commonly encountered skin cancers. This study analysed the various histologic variants of cutaneous Squamous Cell Carcinoma (cSCC) diagnosed in the hospital. The study was a descri...

[View Abstract](#)

👤 Adekunle V.O, Ofusori D.A

Morphometric Effects of β , γ -Carotene- 3, 3'-diol on Indomethacin- induced Gastric Ulcer in Adult Male Wistar Rats

Stomach ulcer is a disease that causes the erosion of the epithelial lining of the gastric mucosa. This study aims to determine the ameliorative effects of β , γ - carotene- 3, 3'-diol following Indomethacin-induced gastric ulcer in Wistar rats. Thirty adult male Wistar rats (7 weeks old, 160 – 165...

[View Abstract](#)

👤 Musa Ibrahim Kurawa

Detrimental Effects of Mosquito Coil Smoke Exposure on Memory and Hippocampal Morphology

Mosquito coil smoke is widely used to repel mosquitoes and prevent malaria in many malaria-endemic regions. While there is considerable awareness of the toxicity of carbon monoxide (CO), exposure to CO from less obvious sources like MC smoke is often overlooked. This study investigates the neuro...

[View Abstract](#)

👤 Olatunji Sunday Yinka and Nwoha Polycarp

Cannabidiol Modulates Hippocampal Neuronal Morphology and Attenuates Reactive Astrogliosis in Animal Model of Epilepsy

Intractable form of epilepsy is a major challenge worldwide, and despite an increasing number of medical therapies in the treatment of epilepsy, about one in three patients continues to experience seizures. This study evaluated the effects of cannabidiol on neuronal morphology and astrocytes in ...

[View Abstract](#)

👤 Olakunle Olutoye Osinubi, Peter Chuka Ozegbe, Oluwasanmi Olayinka Aina

Histological, Histochemical and Immunohistochemical Characterization of the Testis of Double-spurred Francolin (*Francolinus bicalcaratus*) in a Reproductive Cycle

The double-spurred Francolin is a feral bird and remains undomesticated with very little known about its reproductive capability. This study aimed to know the structural integrity of its testis. Birds (n=5), were obtained during the dry and rainy seasons from their natural habitat in Kano, stabi...

[View Abstract](#)

Search for articles..

ADVERTISEMENT 270 X 270



**Connect With Your Peers
on Issues in Medicine**

Consectetur adipisicing elit sedo
amod tempor incididunt.

[Join In Now](#)



ADVERTISEMENT 270 X 270



Connect With Your Peers on Issues in Medicine

Consectetur adipisicing elit sedo
amod tempor incididunt.

[Join In Now](#)



**ANATOMICAL
SOCIETY OF
NIGERIA**

The Journal of Anatomical Sciences is the official journal of the Anatomical Society of Nigeria and publishes original research articles dealing with morphological sciences in all vertebrates. [Read More](#)



Quick links

[Submit Article](#)

[Editorial Policies](#)

[Renew Subscription](#)

[Get Help and Support](#)

Get In Touch

 Department of Anatomy, Faculty of Basic Medical Sciences,
University of Ilorin, PMB 1515, Ilorin

 (+234) 703 050 5707

 (+234) 706 743 4502

 chiefeditor.asnjournal@gmail.com

 sendpapertojas@gmail.com

 journalofanatomicalsciences@gmail.com



The Journal of Anatomical Sciences

Email: journalofanatomicalsciences@gmail.com

J. Anat Sci 15(2)

Submitted August 13th, 2024

Accepted August 29th, 2024

Published September 30th, 2024

Comparative Analysis of the Protective Potentials of *Nigella sativa* and *Lepidium meyenii* on Alcohol-induced Testicular Toxicity in Adolescent Wistar Rats

¹Adana Misturah Yetunde, ¹Towoju Alaba Magrette, ¹Alade Oluwadamilola Esther, ¹Oluwasegun Blessing Olaitan, ²Adewale David Adebayo, ¹Onigbolabi Oluwafemi Gideon, ¹Ajao Moyosore Salihu

¹Department of Anatomy, Faculty of Basic Medical Sciences, College of Health Sciences, University of Ilorin, Ilorin

²Department of Anatomy, ECWA College of Nursing Sciences, Egbe, Kogi State, Nigeria

Corresponding Author: Adana M. Y.

Email: adana.my@unilorin.edu.ng

ABSTRACT

Alcohol, a psychoactive drug, is soluble in both water and lipids, due to which it can diffuse to all tissues and affect the normal functioning of the body. Gonadal toxicity is reported as one of the side effects of its long-term consumption. This study examined the possible comparable protective potentials of *Lepidium meyenii* (LM) and *Nigella sativa* oil (NSO) on the biological integrities of the testes after exposure to excessive alcohol. Thirty-six adolescent Wistar rats (60–150 g) were randomly divided into six groups and treated orally for 56 consecutive days. Group treatments included normal saline for the control group, 40% alcohol, LM, NSO, LM+alcohol, and NSO+alcohol. On the 57th day, samples were collected to assess reproductive hormones, sperm analysis, testicular histology and proliferating cell nuclear antigen (PCNA) expression. The results revealed that excessive alcohol consumption affected the structural integrities of the testes by depleting the mature sperm cell population and actively dividing PCNA immune-reactive cells. The treatment with *Lepidium meyenii* does not show any significant protective effects on alcohol-induced structural distortion of rats' testes. However, NSO promises to be effective in protecting against alcohol-induced changes.

Keywords: testes, alcohol, germinal epithelium, *Lepidium meyenii*, *Nigella sativa*

INTRODUCTION

Infertility is a disease of the male or female reproductive system defined by the failure to achieve a pregnancy after 12 months or more of regular unprotected sexual intercourse¹. Estimates suggest that between 48 million couples and 186 million individuals live with infertility globally¹. In the male reproductive system, infertility is most commonly caused by problems in the ejection of semen, absence or low levels of sperm, or abnormal shape (morphology) and movement (motility) of the sperm². However, this incidence varies from one region to the other, with the highest reports in the infertility belt of Africa, which includes Nigeria^{3,4}. Unfortunately, about 50% of these infertility cases are male factor-related⁵.

Alcohol has been reported as one of the causative factors of male infertility. Clinically, the most important endocrine consequences of long-term alcohol use are its effects on the gonads as it affects the synthesis of testosterone, consequently reducing sperm parameters⁶ and nuclear maturity as well as the DNA integrity of spermatozoa^{7,8}. Many men with infertility consume alcohol. The big problem in the management of male factor infertility associated with alcohol consumption is the addictive nature and the rate of relapse among participants⁹.

In this era of advanced biomedical technology, medical management of male infertility includes in-vitro fertilization (IVF), sperm freezing, donation techniques, and the use of pro-fertility drugs such as clomiphene citrate. In developing countries, medicinal

plants such as *Nigella sativa* and other traditional fertility supplements (such as maca) are used as alternatives to modern therapies ¹⁰.

Nigella sativa is a dicotyledonous of the Ranunculaceae family which is an amazing herb with a rich historical and religious background ^{11,12} found in southern Europe, northern Africa, and Asia Minor. Historically, it has been recorded that *Nigella sativa* seeds were prescribed by ancient Egyptian and Greek physicians to treat headaches, nasal congestion, toothache and intestinal worms, diuretics, and to increase milk production. The seeds of *Nigella sativa* have long been used in the Middle and Far East as a traditional medicine for a wide range of illnesses ¹³.

Lepidium meyenii (LM), popularly known as maca, is a Peruvian hypocotyl found in the central Andes. Traditionally, its roots have been used by people residing in the high altitudes of the Andean subregion as a nutrient, energizer, aphrodisiac and for its fertility-enhancing properties ¹⁴. Several studies have reported that Maca root preparations improved sexual function in healthy adult males ¹⁵ and in males with drug-induced sexual dysfunction ¹⁶ or erectile dysfunction ¹⁷. In rodents, there are reports that administration of LM enhances spermatogenesis, fertility and sexual behavior, likely due to its phytoesterol or phytoestrogen content ^{18,19} and its androgen-like effects ¹⁷. LM contains several bioactive compounds (maca nutrients, macaridine, glucosinolates, maca alkaloid, macamides and macaene), which have also been suggested to be responsible for their beneficial effects on sexual function ^{18,19}. The goal of this study is to compare the protective potentials of *Nigella sativa* and *Lepidium meyenii* on alcohol-induced testicular toxicity in adolescent Wistar rats

MATERIALS AND METHODS

Animal care

Thirty-six adolescent male Wistar rats weighing 60-150 g were collected from the University of Ilorin Biological Garden. They were kept under standard laboratory conditions (12 hour light/dark cycles at 25-28°C) in the animal house facility of the Faculty of Basic Medical Sciences, University of Ilorin, Ilorin. They were fed daily with pelletized grower feed containing 15% crude protein, 7% fat, 10% crude fibre, 1% calcium, 0.35% phosphorus, and 2.55 kcal/kg of metabolized energy as indicated. Portable water was provided, and a hygienic environment was ensured. They were cared for and used following the standard laboratory animal care and use guide.

Materials used

Alcohol (40%) was purchased from a local shop at Agbade in Ilorin. *Nigella sativa* oil and *Lepidium meyenii* (1000 mg) were purchased in the state. This research was approved by the University of Ilorin ethical review committee (UERC) (UERC/ASN/2019/856). The study was carried out upon the recommendation of the College of Health Sciences ethical review committee in compliance with the Institutional Animal Care and Use Committee (IACUC).

Experimental design

Thirty-six 8-week-old male Wistar rats were randomly placed into six groups of six rats each. They were subjected to an adaptation period of 14 days in the animal facility before the commencement of the experiment. The six groups include Group DW, which received distilled water; Alcohol group, received 3.2 ml/kg of alcohol; LM group received 100 mg/kg of *Lepidium meyenii*; NSO group received 0.5 ml/kg *Nigella sativa* oil; LM + alcohol group received 100 mg/kg *Lepidium meyenii* and 3.2 ml/kg of 40% alcohol; and NSO + alcohol received 0.5 ml/kg of NSO and 3.2 ml/kg of alcohol. All administrations were via the oral route. The rats were weighed weekly for dosage adjustments, and the experiment lasted 56 days.

Experimental termination and sample collection

The rats were euthanized a day after the treatments under anesthesia by administration of intraperitoneal ketamine, 100 mg/kg. Blood was collected for hormonal assay, while the testes were excised, weighed, fixed in Bouin's fluid and subsequently processed for histological and immunohistochemical analysis. Also, the cauda epididymis of each rat was excised and gently minced in 2 ml of normal saline. The resulting suspension was used for the determination of semen analysis. A drop of the suspension on the glass slide was viewed under the light microscope. The percentage of motile spermatozoa was derived by subtracting the number of still spermatozoa from the total sperm count according to the methods of Tjioe and Oentoeng ²⁰. Spermatozoa exhibiting any motion after the dilution were considered motile. Sperm concentration was determined by immobilizing cells in 3% saline. This was then thoroughly mixed and assessed using a Makler Counting Chamber (Sef- Medical Instruments Ltd Haifa) according to the method of Eliasson. ²¹ A sperm suspension of 10 µl was placed onto the Makler Chamber stage and counted for the number of sperms under a light microscope (Olympus).

Reproductive hormone assay

The serum reproductive hormones i.e. follicle-stimulating hormone (FSH), testosterone and Luteinizing hormone (LH) levels were assayed using a commercially available kit (IBL) and the manufacturer's instructions. The kit employs a competition-based solid-phase enzyme-linked immunosorbent assay (ELISA). The samples were plated in duplicate, and the concentrations of testosterone (catalogue no. CSB- E05100r), FSH (catalogue no. CSB- E12654r) and LH (catalogue no. CSB- E06869r) were determined using a standard curve with 5 standard values. Using a microplate reader ²², the optical density of the samples and testosterone, FSH and LH standards were quantified at 450 nm.

Histological and morphometric studies

For histopathological examination, fixed testicular tissues were dissected, processed through graded ethanol and embedded in paraffin. Microtome sections at 5 µm were obtained and stained with hematoxylin and eosin using a microtome (micron HM 315 microtome). The slices were examined and images were captured using an Olympus light microscope (Olympus BX51) and a camera (Olympus E330, Olympus Optical Co. Ltd.). Adopting a systematic random sampling, a total of eleven seminiferous tubules were carefully chosen for each rat, and Image J software was used to estimate and measure the lumen and total seminiferous tubule diameters. ^{23,24}

Proliferative nuclear cell antigen (PCNA) studies

The PCNA study was done using immunohistochemical studies. Five blocks per testis for the PCNA study were immunostained in each group. The carefully chosen paraffin blocks for immunohistochemical studies were stained with monoclonal antibody PCNA (LabVision, avidin-biotin-peroxidase method). Deparaffinized sections were rehydrated with graded alcohols after being deparaffinized in xylol. They were then immersed in 0.01 M citrate buffer (pH 6) and microwaved (Gibson's oven) on high for 20 min before cooling for at least 20 min at room temperature. Endogenous peroxidase activity was inhibited by a 10-minute incubation in phosphate buffer saline with 3% hydrogen peroxide (PBS, 10 mM Na₂PO₄, 140 mM NaCl, pH 7.2). The sections were rinsed thrice for 2 minutes each before beginning the immunostaining

process ²⁵. Histostainplus kit (Zymed) immunohistochemical staining was conducted using the streptavidin-biotin method, which contains 10% nonimmune serum and secondary antibody (biotinylated) and Streptavidin-peroxidase. Sections were rinsed multiple times with PBS before being treated with blocking serum for 10 min to inhibit unspecific binding. After removing the excess blocking serum, the histological sections were coated with primary antibody PCNA in dilution (1:50) for one hour at room temperature. For 10 min, the secondary antibody was utilized as a second layer. Then, the sections were washed for 10 min with PBS and streptavidin-peroxidase complex. Following that, the sections were rinsed with PBS, and the peroxidase signal was generated in phosphate buffer saline containing 0.05% diaminobenzidine (DAB) and 0.01% hydrogen peroxide (H₂O₂). Before drying and mounting per mount, the sections were faintly counter-stained in hematoxylin. Negative control slides were created by replacing the primary antibody with PBS in identical areas. In the same run, slides with normal positive reactivity tissues of the antibody were also employed. Nuclear staining was observed in brown PCNA expression instances ²⁵.

Statistical analysis

Statistical data were analyzed using GraphPad Prism 8.4.2 and SPSS version 20 (Statistical Package for the Social Sciences, version 20.0). Results were presented as mean ± SEM, and all statistical comparisons were performed using one-way analysis of variance (ANOVA). The test of significance was also performed to test any significant differences between different treatments (the level of significance was set at P < 0.05). The post-hoc test used to determine the test of significance was Turkey. Cell counting was done using Image J.

RESULTS

Alcohol-induced emaciation in rats

The weighting of rats revealed significant physical changes in the weight of the animals across the different groups. The rats in the Alcohol group experienced a substantial difference in weight gain when compared to groups NS, LM, and NSO. Groups LM + alcohol and NSO + alcohol displayed a significant increase in weight when compared to Alcohol. There was no significant difference in the relative testicular weight (see Table 1).

Table 1: Body weight changes in experimental animals

Groups	IBW	FBW	WC	TW	RTW
NS	81.33±2.12	137.20±2.24	55.87±3.22	2.40±0.13	1.74±0.07
Alcohol	124.50±5.63	173.33±13.03	48.83±7.70*	2.06±0.27	1.23±0.20
LM	124.00±6.57	202.83±13.50	78.83±16.11* ^α	2.57±0.12	1.28±0.06
NSO	127.50±6.72	208.00±13.41	80.50±9.93* ^α	2.54±0.16	1.22±0.04
LM+Alcohol	125.17±6.96	206.00±12.71	80.83±12.07* ^α	2.82±0.13	1.37±0.05
NSO+Alcohol	123.83±7.87	188.40±15.50	64.57±9.14	2.58±0.11	1.40±0.08

IBW=Initial body weight, FBW=Final body weight, weight changes, TW=Testicular weight, RTW=Relative testicular weight. * indicates a significant difference when compared to DW. ^α indicates a significant difference when compared to Alcohol.

***Nigella sativa* oil prevented semen composite depletion in alcohol-induced testicular toxicity**

Repeated exposure to excessive consumption of 40% Alcohol caused significant ($p < 0.05$) reductions in the sperm population and a drop in their motility when

compared to the control, LM + alcohol and NSO + alcohol-treated rats (Figure 1). The sperm concentration and motility were significantly increased ($p < 0.05$) in NSO and NSO + alcohol when compared to the rats only exposed to excessive alcohol (Figure 1).




Figure 1: Semen analysis in rats

Rats exposed to normal saline (NS), 40% alcohol (Alcohol), *Lepidium meyenii* (LM), *Nigella sativa* oil (NSO), *Lepidium meyenii* + 40% alcohol (LM + alcohol), *Nigella sativa* oil + 40% alcohol (NSO + alcohol) respectively. Bars indicate mean ± SD, (*) and (***) represent statistically significant differences ($p < 0.05$ and $p < 0.001$ respectively) compared to DW; ^{ααα} indicate statistical difference ($p < 0.001$) between the groups and the Alcohol group.

Differential hormonal responses following administration of alcohol, *Nigella sativa* oil and *Lepidium meyenii*

The testosterone levels of the rats exposed to alcohol were significantly higher when compared to the control. Treatment with NSO + alcohol stimulated increased testosterone levels, while treatment with

LM + alcohol was observed to reduce testosterone levels (Figure 2). The NSO group had testosterone levels comparable to that of the control. In contrast to the effects of the NSO on testosterone concentration, NSO and Alcohol significantly increased the luteinizing hormone concentration in treated rats (Figure 4), while the FSH concentrations were not significantly different in all groups (Figure 3).




Figure 2: Serum testosterone in rats

Normal saline (NS), 40% alcohol (Alcohol), *Lepidium meyenii* (LM), *Nigella sativa* oil (NSO), *Lepidium meyenii* + 40% alcohol (LM + alcohol), *Nigella sativa* oil + 40% alcohol (NSO + alcohol) respectively. Bars indicate mean $\pm$ SD




Figure 3: Serum Follicle Stimulating Hormone in rats

Normal saline (NS), 40% alcohol (Alcohol), *Lepidium meyenii* (LM), *Nigella sativa* oil (NSO), *Lepidium meyenii* + 40% alcohol (LM + alcohol), *Nigella sativa* oil + 40% alcohol (NSO + alcohol) respectively. Bars indicate mean $\pm$ SD




Figure 4: Serum Luteinizing Hormone in rats

Normal saline (NS), 40% alcohol (Alcohol), *Lepidium meyenii* (LM), *Nigella sativa* oil (NSO), *Lepidium meyenii* + 40% alcohol (LM + alcohol), *Nigella sativa* oil + 40% alcohol (NSO + alcohol) respectively. Groups Alcohol, LM + alcohol and NSO + alcohol displaced altered serum LH. Bars indicate mean $\pm$ SEM, α represents $p < 0.05$

Immunohistochemical studies of proliferative nuclear cell antigen

In Figure 5 below, early germ cells along the basement membrane are deeply stained (black arrow), whereas

several spermatocytes show less intense (orange arrows) staining. Sertoli cells (green) and Leydig cells (red arrow) stain negative for PCNA (Original magnification X100).




Figure 5: Photomicrograph of the testes showing the Proliferative cell nuclear antibodies

The control group and experimental groups. (Control group (NS), 40% Alcohol (Alcohol), *Lepidium meyenii* (LM), *Nigella sativa* oil (NSO), *Lepidium meyenii* and 40% alcohol (LM + alcohol), *Nigella sativa* oil and 40% alcohol (NSO + alcohol). Green arrow (secondary spermatocyte), Orange arrow (primary spermatocyte), Blue arrow (seminiferous tubule), Red arrow (Leydig cells). Depletion of PCNA-stained cells in Alcohol, NSO, LM + alcohol and NSO + alcohol groups.

Table 2: Population of PCNA-stained cells

Animal Groups	Cell Counts
NS	1702 ±132.3
Alcohol	995 ± 377.0*
LM	1887 ±249.7 ^a
NSO	1074 ± 173.5
LM + Alcohol	1555 ±270.7 ^a
NSO + Alcohol	1256 ±161.3 ^a

*indicates a significant difference compared to NS

^a indicates a significant difference compared to Alcohol

Histological observations

Observations from the basic histology using light microscopy revealed normal seminiferous tubules in

all groups, with a normal progression of the cells of the germinal epithelium. However, there was a widened interstitium in the NSO group, but the sizes of the seminiferous tubules were maintained compared to the control (NS) group. See Figure 6.




Figure 6: Representative photomicrographs of H&E stained rats' testis

Normal saline (NS), 40% alcohol (Alcohol), *Lepidium meyenii* (LM), *Nigella sativa* oil (NSO), *Lepidium meyenii* + 40% alcohol (LM + alcohol), *Nigella sativa* oil + 40% alcohol (NSO + alcohol) respectively. a- Lumen, b- Seminiferous tubule, c- Primary spermatocyte, d- Interstitium/interstitial space, e- Spermatogonia. H&E stain, ×100

DISCUSSION

Sperm cell development begins with the differentiation of spermatogonia in the seminiferous tubules, which then change into mature spermatozoa after a sequence of events. Spermatogenesis must proceed unobstructed and undisturbed to create viable spermatozoa with reproductive potential. Due to the sensitivity and delicate features of sperm cells, an insult to the organism, particularly when directed to the reproductive organs, such as the testes, could distort spermatogenesis and impair reproductive abilities²⁶. The structural state and functionality of the testis were evaluated in this research work to determine the toxic effect of alcohol and comparable protective roles of *Lepidium meyenii* and NSO in the reproductive parameters of male Wistar rats.

Previously, testicular atrophy has been noted as the consequence of chronic alcohol usage in humans as well as in experimental animal models, and alcoholism has been linked with reduced testicular volume and size²⁷. While many researchers have observed atrophy in testicular weight, some have reported an increase²⁸. However, the report from this study reports that neither alcohol, *Lepidium meyenii*, nor NSO-treated rats displayed an appreciable change in the relative testicular weight. This is similar to a recent finding where no change in weight was observed in both ethanol and *Nigella sativa*-treated animals²⁹. It was experiential that treatment with alcohol did not affect the relative weights of the reproductive organs. Results from previous studies revealed decreased testicular weight and a reduced organ coefficient as an indication of alcohol-induced testicular atrophy³⁰. A reduced organ coefficient might be due to increased apoptosis in the respective organs as increased apoptosis is associated with weight loss in mice testis³⁰. Moreover, NSO restored the alcohol-induced reduction in body weight and organ coefficient.

Testosterone level is a good indication of the reproductive functioning in males. Exposure to alcohol in this study is associated with an increased level of testosterone. Similarly, prior studies have shown that modest consumption of alcohol increases serum testosterone levels, particularly in short-term exposures^{31,32}, as observed in the present study. This has been attributed to the increased hepatic enzyme activity during such exposures³³. However, in humans and animal models, during chronic exposures to excessive alcohol, the production of testosterone drops^{34,35}.

The serum luteinizing hormone levels displayed reverse effects, as they dropped with alcohol exposure. On the other hand, alcohol, *Lepidium meyenii*, and NSO did not affect the follicle-

stimulating hormone levels. This is similar to the findings of other researchers^{35,36}.

From the result of this study, sperm count and sperm motility in rats treated with 40% alcohol were reduced. The co-administration of NSO with alcohol markedly reversed the adverse effect induced by alcoholism to within control levels and this probably results from the antioxidant activities of its bioactive components. However, treatment with *Nigella sativa* reversed the toxicity caused by alcohol. This is similar to the results from previous studies²⁹.

The semen parameters are a measure of both endocrine and testicular health. It also has a direct relationship with the reproductive capabilities of males. Insults to the testes are easily picked by an assessment of the sperm concentration, motility and percentage of sperms with normal morphology. Previous studies have also shown that changes in sperm parameters might be due to the direct effect of alcohol on testicular tissue which led to impaired sperm parameters³⁷. Moreover, the decrease in sperm count in alcohol-treated rats might be due to decreased testosterone levels observed in chronic use³⁴ and increased oxidative stress. Additionally, reduced sperm motility might be a result of increased head-tail breakage in alcohol-treated rats as explained by Anderson *et al*³⁸. The improvement in sperm parameters observed with concomitant use of NSO might be attributed to its anti-oxidative properties³⁹. However, results from this study show that the co-administration of alcohol with *Lepidium meyenii* has not proven to be beneficial.

Alcohol has been found to increase the epithelial barrier permeability and affect tight junction-associated protein expression⁴⁰. This might be a reason behind the profound destruction of seminiferous tubules, disorganized lining of seminiferous tubules and sloughing of germ cells. The histopathological examinations of testes showed that Alcohol had a deleterious effect on the germinal epithelium and co-administration with NSO was unable to alleviate the adverse effects induced by Alcohol. Data from this study supports those obtained by Hala who reported that pre-treatment of male rats with *Nigella sativa*, and celery oils for 4 weeks produced a protective effect against testicular injury induced by sodium valproate⁴¹. This effect was manifested by increased weight of the testis, improved semen quality and quantity, elevated serum testosterone level, decreased lipid peroxidation in the testis, and alleviation of degenerative changes in the testes of rats given sodium valproate. The mechanism for this protective effect of NSO against the toxic effects of alcohol can be attributed to its free radical scavenging activity and increased antioxidant enzymes in rats³⁹.

The role of proliferating cell nuclear antigen (PCNA) in the testes is well established. It plays an essential role in nucleic acid metabolism, cell replication and repair. The expression of PCNA in the Alcohol-treated groups was diminished. This is suggestive of reduced cellular replication and damage to the actively dividing portion of the germinal epithelium lined along the basement membrane of the seminiferous tubules – spermatogenic cells and the primary spermatocytes. Damage to the mitotic germinal epithelium is also indicative of DNA insults. While some researchers believe that the expression of PCNA indicates a good response to impaired spermatogenesis⁴², others suggest that it is an indication of well-regulated spermatogenesis⁴³.

Nigella sativa oil has been proven to display a wide range of favourable biological activities, the most prominent being antioxidant and anti-inflammatory agents. Co-administration with Alcohol was shown to limit these structural and biochemical changes in the gonads. Many researchers have reported gonadal protection against insults following testicular exposure to different varieties of insults^{44,45}.

Conclusion

Besides being an important public and social issue, alcohol consumption can also significantly impact male reproduction by inducing testicular damage evidenced by abnormal sperm parameters. However, treatment with *Nigella sativa* oil protected against alcohol-induced testicular toxicity. This result shows that *Nigella sativa* oil could be a promising supplement for preventing male infertility induced by excessive alcohol consumption.

Acknowledgements

The authors appreciate the contributions of the technical staff of the Department of Anatomy, University of Ilorin.

REFERENCES

- WHO. Infertility. 2020. WHO fact sheet. Accessed 2024. <https://www.who.int/news-room/fact-sheets/detail/infertility>
- Oehninger S, Kruger TF. Sperm morphology and its disorders in the context of infertility. *F&S Reviews*. 2021;2(1):75-92. <https://doi.org/10.1016/j.xfnr.2020.09.002>
- Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the globe. *Reproductive Biology and Endocrinology*. 2015;13(1):37. doi:10.1186/s12958-015-0032-1
- Bowa K, Kachimba JS, Mapulanga V, Chalwe M, Manda E. Male infertility from the developing nation perspective. *Male Infertility: Contemporary Clinical Approaches, Andrology, ART and Antioxidants*. 2020:855-63.
- Okonofua FE, Ntoimo LF, Omonkhua A, Ayodeji O, Olafusi C, Unuabonah E, et al. Causes and risk factors for male infertility: a scoping review of published studies. *International Journal of General Medicine*. 2022:5985-97. <https://doi.org/10.2147%2FIJGM.S363959>
- Finelli R, Mottola F, Agarwal A. Impact of alcohol consumption on male fertility potential: a narrative review. *International Journal of Environmental Research and Public Health*. 2021;19(1):328. <https://doi.org/10.3390%2FIjerp19010328>
- Talebi AR, Sarcheshmeh AA, Khalili MA, Tabibnejad N. Effects of ethanol consumption on chromatin condensation and DNA integrity of epididymal spermatozoa in the rat. *Alcohol*. 2011;45(4):403-9. <https://doi.org/10.1016/j.alcohol.2010.10.005>
- Akang EN, Oremosu AA, Osinubi AA, James AB, Biose IJ, Dike SI, et al. Alcohol-induced male infertility: Is sperm DNA fragmentation a causative. *J. Exp. Clin. Anat*. 2017;16(1):53. http://dx.doi.org/10.4103/jeca.jeca_14_16
- Guliyev C, Ince-Guliyev E, Ögel K. Predictors of relapse to Alcohol and Substance Use: are there any differences between 3 and 12 months after Inpatient Treatment? *Journal of psychoactive drugs*. 2022;54(4):358-67. <https://doi.org/10.1080/02791072.2021.1976887>
- Abarikwu SO, Onuah CL, Singh SK. Plants in the management of male infertility. *Andrologia*. 2020; 52(3):e13509. <https://doi.org/10.1111/and.13509>
- Dabeer S, Rather MA, Rasool S, Rehman MU, Alshahrani S, Jahan S, et al. History and traditional uses of black seeds (*Nigella sativa*). *Black seeds (Nigella sativa)* 2022; 1-28. Elsevier. <https://doi.org/10.1016/b978-0-12-824462-3.00016-0>
- Ahmad MF, Ahmad FA, Ashraf SA, Saad HH, Wahab S, Khan MI et al. An updated knowledge of Black seed (*Nigella sativa* Linn.): Review of phytochemical constituents and pharmacological properties. *Journal of Herbal Medicine* 2021;25:100404. <https://doi.org/10.1016/j.hermed.2020.100404>
- Majeed A, Muhammad Z, Ahmad H, Hayat SS, Inayat N, Siyyar S. *Nigella sativa* L.: Uses in traditional and contemporary medicines—An overview. *Acta Ecologica Sinica*. 2021;41(4):253-8. <https://doi.org/10.1016/j.chnaes.2020.02.001>
- Gonzales GF, Gonzales C, Gonzales-Castañeda C. *Lepidium meyenii* (Maca): a plant from the highlands of Peru—from tradition to science. *Forschende Komplementärmedizin/Research in Complementary Medicine*. 2009;16(6):373-80. <https://doi.org/10.1159/000264618>

15. Shin D, Jeon SH, Piao J, Park HJ, Tian WJ, Moon DG et al. Efficacy and safety of maca (*Lepidium meyenii*) in patients with symptoms of late-onset hypogonadism: a randomized, double-blind, placebo-controlled clinical trial. *The World Journal of Men's Health*. 2023;41(3):692. <https://doi.org/10.5534/wjmh.220112>
16. Winter J, Curtis K, Hu B, Clayton AH. Sexual dysfunction with major depressive disorder and antidepressant treatments: impact, assessment, and management. *Expert Opin Drug Saf*. 2022;21(7):913-930. <https://doi.org/10.1080/14740338.2022.2049753>
17. Lee HW, Lee MS, Kil KJ. Maca (*L. meyenii*) for erectile dysfunction: a systematic review and meta-analysis. *Journal of Men's Health*. 2023;19(1):1-6.
18. Chen R, Wei J, Gao Y. A review of the study of active components and their pharmacology value in *Lepidium meyenii* (Maca). *Phytotherapy Research*. 2021;35(12):6706-19. <https://doi.org/10.1002/ptr.7257>
19. Wang Y, Wang Y, McNeil B, Harvey LM. Maca: An Andean crop with multi-pharmacological functions. *Food Research International*. 2007;40(7):783-92. <https://doi.org/10.1016/j.foodres.2007.02.005>
20. Tjioe D, Oentoeng SJF. The viscosity of human semen and the percentage of motile spermatozoa. *Fertility and Sterility*. 1968;19(4):562-565. [https://doi.org/10.1016/s0015-0282\(16\)36728-0](https://doi.org/10.1016/s0015-0282(16)36728-0)
21. Makler A. A new chamber for rapid sperm count and motility estimation. *Fertility and Sterility*. 1978;30(3):313-318. [https://doi.org/10.1016/s0015-0282\(16\)43518-1](https://doi.org/10.1016/s0015-0282(16)43518-1)
22. Joelsson D, Moravec P, Troutman M, Pigeon J, DePhillips P. Optimizing ELISAs for precision and robustness using laboratory automation and statistical design of experiments. *Journal of Immunological Methods*. 2008;337(1):35-41. <https://doi.org/10.1016/j.jim.2008.05.012>
23. Adana MY, Akang EN, Peter AI, Jegede AI, Naidu EC, Tiloke C, et al. Naringenin attenuates highly active antiretroviral therapy-induced sperm DNA fragmentations and testicular toxicity in Sprague-Dawley rats. *Andrology*. 2018;6(1):166-75. <https://doi.org/10.1111/andr.12439>
24. Howard V, Reed M. Unbiased stereology: three-dimensional measurement in microscopy. Second ed. Liverpool: Garland Science, Taylor & Francis Group; 2004.
25. Foley JF, Dietrich DR, Swenberg JA, Maronpot RR. Detection and evaluation of proliferating cell nuclear antigen (PCNA) in rat tissue by an improved immunohistochemical procedure. *Journal of Histotechnology*. 1991;14(4):237-241. <http://dx.doi.org/10.1179/his.1991.14.4.237>
26. Bhartiya D, Kaushik A. Testicular stem cell dysfunction due to environmental insults could be responsible for deteriorating reproductive health of men. *Reproductive Sciences*. 2021;28(3):649-58. <https://doi.org/10.1007/s43032-020-00411-3>
27. Pozza C, Kanakis G, Carlomagno F, Lemma A, Pofi R, Tenuta M, et al. Testicular ultrasound score: a new proposal for a scoring system to predict testicular function. *Andrology*. 2020;8(5):1051-63. <https://doi.org/10.1111%2Fandr.12822>
28. Handelsman DJ, Staraj S. Testicular size: the effects of aging, malnutrition, and illness. *Journal of Andrology*. 1985;6(3):144-51. <https://doi.org/10.1002/j.1939-4640.1985.tb00830.x>
29. Sanpinit S, Wetchakul P, Chonsut P, Ngamdokmai N, Ahmad AR, Warinhomhoun S. Repeated 28-Day Oral Toxicological Study and Gastroprotective Effects of *Nigella sativa* L. Oil (Shuhada) against Ethanol-Induced Gastric Mucosal Injury in Rats. *Nutrients*. 2023;15(6):1532. <https://doi.org/10.3390/nu15061532>
30. Shaha C, Tripathi R, Mishra DP. Male germ cell apoptosis: regulation and biology. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2010;365(1546):1501-15. <https://doi.org/10.1098%2Frstb.2009.0124>
31. Smith SJ, Lopresti AL, Fairchild TJ. The effects of alcohol on testosterone synthesis in men: a review. *Expert Review of Endocrinology & Metabolism*. 2023;18(2):155-66. <https://doi.org/10.1080/17446651.2023.2184797>
32. Sarkola T, Eriksson CP. Testosterone increases in men after a low dose of alcohol. *Alcoholism: Clinical and Experimental Research*. 2003;27(4):682-5. <https://doi.org/10.1097/01.alc.0000060526.43976.68>
33. Zhang M, Yin D, Kong F. The changes of serum testosterone level and hepatic microsome enzyme activity of crucian carp (*Carassius carassius*) exposed to a sublethal dosage of pentachlorophenol. *Ecotoxicology and Environmental Safety*. 2008;71(2):384-9. <https://doi.org/10.1016/j.ecoenv.2007.10.014>
34. Kulkarni SR, Ravindra KP, Dhume CY, Rataboli P, Rodrigues E. Levels of plasma testosterone, antioxidants and oxidative stress in alcoholic patients attending the de-addiction centre. *Biology and Medicine*. 2009;1(4):11-20.
35. Oremosu AA, Akang EN. Impact of alcohol on male reproductive hormones, oxidative stress and semen parameters in Sprague-Dawley rats. *Middle East Fertility Society Journal*. 2015;20(2):114-8. <https://doi.org/10.1016/j.mefs.2014.07.001>
36. Ren JC, Banan A, Keshavarzian A, Zhu Q, LaPaglia N, McNulty J, et al. Exposure to ethanol induces oxidative damage in the pituitary gland.

- Alcohol. 2005;35(2):91-101.
<https://doi.org/10.1016/j.alcohol.2005.03.005>
37. Ricci E, Al Beitawi S, Cipriani S, Candiani M, Chiaffarino F, Viganò P, et al. Semen quality and alcohol intake: a systematic review and meta-analysis. *Reproductive Biomedicine Online*. 2017;34(1):38-47.
<https://doi.org/10.1016/j.rbmo.2016.09.012>
 38. Anderson RA, JR, Willis BR, Oswald C, Zaneveld LJD. Partial reversal of ethanol of ethanol-induced male reproductive pathology following abstinence. *Alcohol and Alcoholism*. 1985;20(3):273-286.
 39. Alenzi FQ, Altamimi MA, Kujan O, Tarakji B, Tamimi W, Bagader O, et al. Antioxidant properties of *Nigella sativa*. *J Mol Genet Med*. 2013;7(3):1-5. <http://dx.doi.org/10.4172/1747-0862.1000077>
 40. Wang Y, Tong J, Chang B, Wang B, Zhang D, Wang B. Effects of alcohol on intestinal epithelial barrier permeability and expression of tight junction-associated proteins. *Molecular Medicine Reports*. 2014;9(6):2352-6.
<https://doi.org/10.3892/mmr.2014.2126>
 41. Hala MA. Protective effect of *Nigella sativa*, linseed and celery oils against testicular toxicity induced by sodium valproate in male rats. *Journal of American Science*. 2011;7:687.
 42. Zhao WP, Wang HW, Liu J, Tan PP, Luo XL, Zhu SQ et al. Positive PCNA and Ki-67 expression in the testis correlates with spermatogenesis dysfunction in fluoride-treated rats. *Biological Trace Element Research*. 2018;186:489-97.
<https://doi.org/10.1007/s12011-018-1338-6>
 43. Fadlalla MB, Wei Q, Fedail JS, Mehfooz A, Mao D, Shi F. Effects of hyper-and hypothyroidism on the development and proliferation of testicular cells in prepubertal rats. *Animal Science Journal*. 2017; 88(12):1943-54.
<https://doi.org/10.1111/asj.12883>
 44. Sayed DA. The protective effect of *Nigella sativa* oil against reproductive toxicity, hormonal alterations, and oxidative damage induced by Sertraline in male rats. *Egyptian Journal of Experimental Biology (Zoology)*. 2019;15(1).
<http://dx.doi.org/10.5455/egyseb.20190325113108>
 45. Mosbah R, Yousef MI, Maranghi F, Mantovani A. Protective role of *Nigella sativa* oil against reproductive toxicity, hormonal alterations, and oxidative damage induced by chlorpyrifos in male rats. *Toxicology and Industrial Health*. 2016;32(7):1266-77.
<https://doi.org/10.1177/0748233714554675>